

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 February 2002 (14.02.2002)


PCT

(10) International Publication Number
WO 02/12341 A2

- (51) International Patent Classification⁷: **C07K 14/705** (74) Agents: PARENT, Annette, S. et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, Eighth Floor, San Francisco, CA 94111-3834 (US).
- (21) International Application Number: PCT/US01/24283
- (22) International Filing Date: 3 August 2001 (03.08.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/632,507 3 August 2000 (03.08.2000) US
- (71) Applicants (for all designated States except US):
CORIXA CORPORATION [US/US]; 1124 Columbia St., Suite 200, Seattle, WA 98104 (US). **SMITHKLINE BEECHAM BIOLOGICALS S.A.** [BE/BE]; 89, Rue de L'Institut, B-B-1330 Rixensart (BE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **CHEEVER, Martin, A.** [US/US]; 6210 SE 22nd, Mercer Island, WA 98040 (US). **GHEYSEN, Dirk** [US/BE]; 89, Rue de L'Institut, B-B-1330 Rixensart (BE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AT (utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, CZ (utility model), DE, DE (utility model), DK, DK (utility model), DM, DZ, EC, EE, EE (utility model), ES, FI, FI (utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: HER-2/NEU FUSION PROTEINS



(57) Abstract: The present invention is generally directed to Her-2/neu fusion proteins, nucleic acid molecules encoding Her-2/neu fusion proteins, viral vectors expressing Her-2/neu fusion proteins, and pharmaceutical compositions (e.g., vaccines) comprising the Her-2/neu fusion proteins and/or nucleic acid molecules encoding the Her-2/neu fusion proteins. The present invention is also directed to methods of treating or preventing cancer by eliciting or enhancing an immune response to the Her-2/neu protein, including for uses in the treatment of malignancies associated with the Her-2/neu oncogene.

BEST AVAILABLE COPY

WO 02/12341 A2

WO 02/12341 A2



Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

HER-2/NEU FUSION PROTEINS

CROSS-REFERENCES TO RELATED APPLICATIONS

The present application claims the benefit of U.S. patent application No.
5 09/632,507, filed August 3, 2000, herein incorporated by reference in its entirety.

The present application is also related to U.S. patent application No.
09/493,480, filed January 28, 2000; and U.S. provisional patent application No.
60/117,976, filed January 29, 1999, herein incorporated by reference in their entirety..

10 STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

Not applicable.

BACKGROUND OF THE INVENTION

15 Despite enormous investments of financial and human resources, cancer
remains one of the major causes of death. For example, cancer is the leading cause of
death in women between the ages of 35 and 74. Breast cancer is the most common
malignancy in women and the incidence for developing breast cancer is on the rise. It is
estimated that one in nine women will be diagnosed with the disease. Standard
20 approaches to cure breast cancer have centered around a combination of surgery,
radiation and chemotherapy. These approaches have resulted in some dramatic successes
in certain malignancies. However, breast cancer is most often incurable, when diagnosed
beyond a certain stage. Alternative approaches to early diagnosis and therapy are
necessary.

25 A common characteristic of malignancies is uncontrolled cell growth.
Cancer cells appear to undergo a process of transformation from the normal phenotype to
a malignant phenotype capable of autonomous growth. Amplification and overexpression
of somatic cell genes is considered to be a common primary event that results in the
transformation of normal cells to malignant cells. The malignant phenotypic
30 characteristics encoded by the oncogenic genes are passed on during cell division to the
progeny of the transformed cells.

At least forty oncogenes operative in malignant cells and responsible for,
or associated with, transformation have been identified. These oncogenes have been

classified into different groups based on the putative function or location of their gene products, such as the protein expressed by the oncogene.

Oncogenes are believed to be essential for certain aspects of normal cellular physiology. In this regard, the Her-2/neu oncogene appears to be a member of the tyrosine kinase family of receptor-like glycoproteins, and shares a high degree of identity with the epidermal growth factor receptor (EGFR). Her-2/neu presumably plays a role in cell growth and/or differentiation. Her-2/neu appears to induce malignancies through quantitative mechanisms that result from increased or deregulated expression of an essentially normal gene product.

The p185 glycoprotein is the protein product of the Her-2/neu oncogene. The Her-2/neu gene is amplified and p185 is overexpressed in a variety of cancers including breast, ovarian, colon, lung and prostate cancer. p185 is related to malignant transformation, and is found in 50-60% of ductal *in situ* carcinomas, in 20-40% of invasive breast cancers, and in a substantial fraction of adenocarcinomas arising in the ovaries, prostate, colon and lung. Her-2/neu expression is intimately associated not only with malignant phenotype, but also with the aggressiveness of the malignancy. Her-2/neu overexpression is correlated with a poor prognosis in both breast and ovarian cancers.

p185 is a transmembrane protein with a predicted relative molecular mass of 185 kD that is about 1255 amino acids in length. p185 has an extracellular domain (ECD) of about 645 amino acids with at least a 40% identity to EGFR, a highly hydrophobic transmembrane domain, and a carboxy terminal intracellular domain (ICD) of about 580 amino acids with at least an 80% identity to EGFR.

There is a need for anti-cancer vaccines that can target a malignancy with which the Her-2/neu oncogene is associated, and for compositions and methods that can elicit and enhance an immune response to the Her-2/neu gene. The present invention is directed to these, as well as other, important ends.

SUMMARY OF THE INVENTION

The present invention provides Her-2/neu p185 fusion proteins, nucleic acid molecules that encode Her-2/neu fusion proteins, and viral vectors that comprise polynucleotide sequences encoding Her-2/neu fusion proteins, for uses that include the immunization of warm-blooded animals against malignancies with which the Her-2/neu oncogene is associated. In one embodiment, the fusion proteins of the present invention

further comprise a TcP0 motif. In another embodiment, the fusion proteins of the invention further comprise an NS1 antigen or an immunogenic fragment thereof, preferably attached to the N-terminus of the fusion protein. Fusion proteins or nucleic acid molecules according to the invention may be present in compositions that include a pharmaceutically acceptable carrier or diluent, *e.g.*, an oil-in water emulsion, and optionally include one or more additional active ingredients, such as an immunostimulatory substance, *e.g.*, SBAS-2, SBAS-4, 3D-MPL, QS21, or a combination of 3D-MPL and QS21. The compositions of the invention are useful as, and can be in the form of vaccines. The fusion proteins, nucleic acid molecules, viral vectors, pharmaceutical compositions and/or vaccines may be administered on a one-time basis (*e.g.*, for an individual with an elevated risk of acquiring or reacquiring a malignancy or when a malignancy is suspected) or on a periodic basis (*e.g.*, for an individual with an elevated risk of acquiring or reacquiring a malignancy or when a malignancy is suspected). Compounds or compositions of the present invention are useful in treating one or more existing tumors, or preventing tumor occurrence or reoccurrence, in warm-blooded animals including humans.

The present invention also provides methods for inhibiting or preventing the development of a cancer in a patient by eliciting or enhancing an immune response to the Her-2/neu protein, comprising administering to a patient a pharmaceutical composition or vaccine as recited above. The patient may be afflicted with, *e.g.*, breast, ovarian, colon, lung or prostate cancer, in which case the methods provide treatment for the disease, or a patient considered at risk for such a disease may be treated prophylactically. In one embodiment, the administration of the pharmaceutical composition or vaccine comprises transfecting cells of a warm-blooded animal *ex vivo* with a nucleic acid molecule of the invention or infecting cells of a warm-blooded animal *ex vivo* with a viral vector comprising a nucleic acid molecule of the invention, and subsequently delivering the transfected or infected cells to the warm-blooded animal. The viral vector is preferably an adenovirus or an alphavirus.

The present invention further provides, within other aspects, methods for removing tumor cells from a biological sample, comprising contacting a biological sample with T cells that specifically react with a Her-2/neu fusion protein, wherein the step of contacting the sample is performed under conditions and for a time sufficient to permit the removal of cells expressing the protein from the sample. Within related

aspects, methods are provided for inhibiting the development of a cancer in a patient, comprising administering to a patient a biological sample treated as described *supra*.

In another embodiment, methods are provided for stimulating and/or expanding T cells specific for a Her-2/neu fusion protein, comprising contacting T cells with one or more of: (i) a fusion protein as described above; (ii) a polynucleotide encoding such a fusion protein; and/or (iii) an antigen presenting cell that expresses such a fusion protein; under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells. Isolated T cell populations comprising T cells prepared as described above are also provided. Within further aspects, the present invention provides methods for inhibiting the development of a cancer in a patient, comprising administering to a patient an effective amount of a T cell population as described above.

In yet another embodiment, the present invention further provides methods for inhibiting the development of a cancer in a patient, comprising the steps of: (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with one or more of: (i) a Her-2/neu fusion protein; (ii) a polynucleotide encoding such a fusion protein; and (iii) an antigen-presenting cell that expresses such a fusion protein; and (b) administering to the patient an effective amount of the proliferated T cells, thereby inhibiting the development of a cancer in the patient. Proliferated cells may, but need not, be cloned prior to administration to the patient.

Finally, the invention provides a method of making a Her-2/neu fusion protein, the method comprising the steps of (a) introducing into a cell an expression vector comprising a polynucleotide encoding a Her-2/neu fusion protein; (b) culturing the transfected cell; and (c) purifying the expressed protein. In a preferred embodiment, the cell is a CHO or a BHK cell. In another preferred embodiment, the cell is cultured in suspension, under serum-free conditions. In yet another embodiment the expressed protein is purified by a two-step procedure, comprising an anion exchange chromatography on Q sepharose High Performance Columns, and a hydrophobic chromatography on Phenyl Sepharose 6 Fast Flow low substitution.

These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows a map of the pFLAGCMV-1/ICD expression plasmid, which has a size of 6.7 kb.

Figure 2 shows a map of the pFLAGCMV-1/KD expression plasmid, which has a size of 5.7 kb.

Figure 3 shows a map of the pFLAGCMV-1/PD expression plasmid, which has a size of 5.7 kb.

Figure 4 shows that the Her-2/neu phosphorylation domain was secreted into the culture media, and that the Her-2/neu intracellular domain and Her-2/neu kinase domain were not secreted into the culture media, as described in Example 3.

Figure 5 shows a map of the pcDNA3.1/hygro/ECD-PD expression vector, which has a size of 8.3 kb.

Figure 6 illustrates the results of the expression of the ECD-PD fusion protein in HEK-293 and CHO cells, and shows that the fusion protein was secreted into the culture media, as described in Example 4.

Figure 7 shows the full length amino acid sequence of the human Her-2/neu protein (SEQ ID NO:1).

Figure 8 shows the full length amino acid sequence of the rat Her-2/neu protein (SEQ ID NO:2). The kinase domain spans the region from amino acid 721 to amino acid 998, inclusively.

Figure 9 shows the amino acid sequence of the extracellular domain of the human Her-2/neu protein (SEQ ID NO:3).

Figure 10 shows the amino acid sequence of the phosphorylation domain (PD) of the human Her-2/neu protein (SEQ ID NO:4).

Figure 11 shows the amino acid sequence of a preferred portion of the phosphorylation domain (APD) of the human Her-2/neu protein (SEQ ID NO:5).

Figure 12 shows the amino acid sequence of a fusion protein comprising the extracellular domain (ECD) and the phosphorylation domain (PD) of the human Her-2/neu protein (SEQ ID NO:6).

Figure 13 shows the amino acid sequence of a fusion protein comprising the extracellular domain (ECD) and a preferred portion of the phosphorylation domain (APD) of the human Her-2/neu protein (SEQ ID NO:7).

Figure 14 shows the amino acid sequence of the extracellular domain (ECD) of the rat Her-2/neu protein (SEQ ID NO:8).

Figure 15 shows the full length nucleotide sequence (SEQ ID NO:9) of a DNA molecule encoding the human Her-2/neu protein and the deduced amino acid sequence of the encoded protein. This full length nucleotide sequence is described in WO 96/30514.

Figure 16 shows the full length nucleotide sequence (SEQ ID NO:10) of a DNA molecule encoding the rat Her-2/neu protein. This full length nucleotide sequence is described by Bargmann *et al.* (1986) *Nature* 319:226-30, and GENBANK/X03362.

Figure 17 illustrates the results of an ELISA assay for Herceptin binding to different ECD-PD fusion proteins produced either in mammalian cells or in *E. coli*. The fusion proteins produced in *E. coli* are in frame with a C- or N-terminal 6 x histidine tag (noted C-His tag and N-His tag, respectively).

Figure 18 shows a comparison of Her-2/neu ECD-PD fusion protein expression in CHO-K1, grown in suspension under serum-free conditions, and in *Pichia* cells.

Figure 19 shows the nucleotide sequence of a DNA molecule encoding the mouse Her-2/neu protein (SEQ ID NO:11).

Figure 20 shows the amino acid sequence of the mouse Her-2/neu protein (SEQ ID NO:14).

Figure 21 shows the results of a Western blot analysis of the expression of the mouse Neu (mNeu), mouse delta Neu (Delta mNeu), mouse ECD-PD (mECDPD) and mouse ECD-PDTcP0 (mECD-PD-TcP0) recombinant proteins in HEK293 lysates. The cNeu3 anti-PD antibody was used for detection of the proteins. Delta mouse Neu has a mutated kinase domain, while mECD-PD-TcP0 contains a 6-mer peptide encoding a ribosomal P-peptide epitope.

Figure 22 shows the amino acid sequence of the mouse ECD-PD fusion protein (SEQ ID NO:27).

Figure 23 shows the nucleotide sequence of a cDNA molecule encoding the mouse ECD-PD fusion protein (SEQ ID NO:28).

Figure 24 shows the amino acid sequence of the mouse ECD-PD-TcP0 fusion protein (SEQ ID NO:29).

Figure 25 shows the nucleotide sequence of a cDNA molecule encoding the mouse ECD-PD-TcP0 fusion protein (SEQ ID NO:30).

Figure 26 shows the amino acid sequence of the TcP0 peptide (SEQ ID NO:31).

Figure 27 shows a nucleotide sequence encoding the TcP0 peptide (SEQ ID NO:32).

5

DESCRIPTION OF THE SPECIFIC EMBODIMENTS

I. INTRODUCTION

The present invention is directed to compounds and compositions capable of modulating, preferably eliciting or enhancing, immunity to the protein product of Her-2/neu oncogene expression, including for malignancies in a warm-blooded animal where an amplified Her-2/neu gene with a malignancy does not require that the protein expression product of the gene be present on the tumor. For example, overexpression of the gene may be involved with initiation and early stages of tumor formation, but the protein expression may subsequently be reduced or absent. The present invention may be used to elicit or enhance an effective immune response to convert a Her-2/neu positive tumor to Her-2/neu negative, in addition to preventing the establishment of Her-2/neu positive tumors and provoking the regression of existing Her-2/neu positive tumors.

The following abbreviations are used throughout the specification: "ECD" refers to the extracellular domain, "ICD" refers to the intracellular domain, "PD" refers to the phosphorylation domain (*i.e.*, the domain that is phosphorylated) that is within the intracellular domain, "ΔPD" refers to a fragment of the phosphorylation domain that is within the phosphorylation domain, and "KD" refers to the kinase domain that is within the intracellular domain. "Delta Her-2/neu" refers to a Her-2/neu protein containing an amino acid change in PD that eliminates the kinase activity. For example, the nucleotide sequence encoding the mouse Delta Her-2/neu protein (Delta mNeu) comprises an A to T nucleotide change at position 2261 of the cDNA encoding the mouse Her-2/neu protein that results in a Lysine to a Methionine amino acid change. The product of expression of the Her-2/neu gene is referred to herein as the "Her-2/neu protein," also known and referred to as "p185" or "c-erbB2."

II. DEFINITIONS

The "Her-2/neu ECD-ICD fusion protein," also referred to herein as "ECD-ICD" or "ECD-ICD fusion protein," refers to a fusion protein (or fragments thereof) comprising the extracellular domain (or fragments thereof) and the intracellular domain (or fragments thereof) of the Her-2/neu protein. As used herein, the ECD-ICD fusion protein does not include a substantial portion of the Her-2/neu transmembrane domain, and preferably does not include any of the Her-2/neu transmembrane domain.

The "Her-2/neu ECD-PD fusion protein," also referred to as "ECD-PD" or "ECD-PD fusion protein," or the "Her-2/neu ECD-ΔPD fusion protein," also referred to as "ECD-ΔPD" or "ECD-ΔPD fusion protein," refer to fusion proteins (or fragments thereof) comprising the extracellular domain (or fragments thereof) and phosphorylation domain (or fragments thereof, *e.g.*, the ΔPD or the PD of the Delta Her-2/neu protein) of the Her-2/neu protein. The ECD-PD and ECD-ΔPD fusion proteins do not include a substantial portion of the Her-2/neu transmembrane domain, and preferably do not include any of the Her-2/neu transmembrane domain.

The terms "Her-2/neu ECD-ICD fusion protein" and "Her-2/neu ECD-PD fusion protein" and their related terms are also understood to refer to fragments thereof, homologs thereof and functional equivalents thereof (collectively referred to as "variants"), such as those in which one or more amino acids is inserted, deleted or replaced by other amino acid(s) or non-amino acid(s) which, in preferred embodiments of the invention, either (i) increase the elicitation or enhancement of an immune response as compared to the Her-2/neu protein, or (ii) do not substantially affect elicitation or enhancement of an immune response as compared to the Her-2/neu protein (*e.g.*, a variant that stimulates a response by helper T cells or cytotoxic T cells or that stimulates the production of antibodies). Specific, non-limiting, examples of variants including exemplary fragments, homologs and functional equivalents of the Her-2/neu ECD-ICD fusion protein and Her-2/neu ECD-PD fusion protein are described in more detail herein. Variants can be "substantially identical" or "substantially similar" to a fusion protein comprising native polypeptide components, and retain the ability to stimulate an immune response.

A "fusion protein" refers to a protein having at least two polypeptides covalently linked, in which one polypeptide comes from one protein sequence or domain and the other polypeptide comes from another protein sequence or domain. The

polypeptides can be linked either directly or via a covalent linker, *e.g.*, an amino acid linker, such as a polyglycine linker, or another type of chemical linker, *e.g.*, a carbohydrate linker, a lipid linker, a fatty acid linker, a polyether linker, *e.g.*, PEG, *etc.* (see, *e.g.*, Hermanson (1996) *Bioconjugate techniques*). The polypeptides forming the fusion protein are typically linked C-terminus to N-terminus, although they can also be linked C-terminus to C-terminus, N-terminus to N-terminus, or N-terminus to C-terminus. The polypeptides of the fusion protein can be in any order. The term "fusion protein" also refers to conservatively modified variants, polymorphic variants, alleles, mutant, subsequences and interspecies homologues of the polypeptides that make up the fusion protein. Fusion proteins may be produced by covalently linking a chain of amino acids from one protein sequence to a chain of amino acids from another protein sequence, *e.g.*, by preparing a recombinant polynucleotide contiguously encoding the fusion protein. Fusion proteins can comprise 2, 3, 4 or more different chains of amino acids from the same or different species. The different chains of amino acids in a fusion protein may be directly spliced together or may be indirectly spliced together via a chemical linking group or an amino acid linking group. The fusion protein may optionally comprise other components, as described in more detail herein.

The term "protein" is used herein interchangeably with "polypeptide" and "peptide."

"Nucleic acid" refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs).

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer *et al.* (1991) *Nucleic Acid Res.* 19:5081; Ohtsuka *et al.* (1985) *J. Biol. Chem.* 260:2605-2608; Rossolini *et al.* (1994) *Mol. Cell.*

Probes 8:91-98). The term nucleic acid is used interchangeably with gene, cDNA, mRNA, oligonucleotide, and polynucleotide.

5 A polynucleotide sequence comprising a fusion protein of the invention hybridizes under stringent conditions to each of the nucleotide sequences encoding each individual polypeptide of the fusion protein. The polynucleotide sequences encoding the individual polypeptides of the fusion polypeptide therefore include conservatively modified variants, polymorphic variants, alleles, mutants, subsequences, and interspecies homologs.

10 "Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino
15 acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 25% sequence identity.
20 Alternatively, percent identity can be any integer from 25% to 100%. More preferred embodiments include at least: 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% or higher, compared to a reference sequence using the programs described herein, preferably BLAST using standard parameters, as described below. One of skill will recognize that these values can be appropriately
25 adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like. "Substantial identity" of amino acid sequences for these purposes normally means sequence identity of at least 40%. Preferred percent identity of polypeptides can be any integer from 40% to 100%. More preferred embodiments
30 include at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99%. Polypeptides which are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having

similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

Optimal alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) *Add. APL. Math.* 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. U.S.A.* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

A preferred example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul *et al.* (1977) *Nuc. Acids Res.* 25:3389-3402 and Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-410, respectively. BLAST and BLAST 2.0 are used, with the parameters described herein, to determine percent sequence identity for the nucleic acids and proteins of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as

defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments (B) of 50, expectation (E) of 10, M=5, N=4, and a comparison of both strands.

Another indication that nucleotide sequences are substantially identical is
5 if two molecules hybridize to each other, or to a third nucleic acid, under moderately, and preferably highly, stringent conditions. Stringent conditions are sequence dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic*
10 *Probes*, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent conditions are selected to be about 5-10°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions
15 will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For
20 selective or specific hybridization, a positive signal is at least two times background, preferably 10 times background hybridization.

Exemplary stringent hybridization conditions can be as following: 50% formamide, 5X SSC, and 1% SDS, incubating at 42°C, or, 5X SSC, 1% SDS, incubating at 65°C, with wash in 0.2X SSC, and 0.1% SDS at 65°C.

25 For the purpose of the invention, suitable "moderately stringent conditions" include, for example, prewashing in a solution of 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridizing at 50°C-65°C, 5X SSC overnight, followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC (containing 0.1% SDS). Such hybridizing DNA sequences are also within the scope of this invention.

30 "Proliferation of T cells," as described herein, includes the multiplication of T cells as well as the stimulation of T cells leading to multiplication, i.e., the initiation of events leading to mitosis and mitosis itself. Methods for detecting proliferation of T cells are discussed below.

III. FUSION PROTEINS OF THE INVENTION

A. Intra- and Extracellular Domains of the Her-2/neu Protein

The Her-2/neu protein was selected as a target for anti-cancer vaccines based on observations of Her-2/neu as described above. One of the major obstacles to this approach is the difficulty of isolating a sufficient amount of Her-2/neu protein. One attempt to address this problem was to express the ECD and ICD separately in mammalian cells. The ECD expressed at a high level as a secreted protein, *i.e.*, about 20 mg of the ECD protein were purified from one liter of mouse cell culture. However, the level of expression of ICD was low, *i.e.*, only about 0.2 mg of the ICD protein were purified from one liter of HEK-293 cultured cells. In addition, the resulting ICD protein was very labile in cell lysate, creating numerous unexpected problems for purification of useful quantities.

As discussed above, Her-2/neu is an oncogenic self protein, and immunological tolerance to self proteins can dampen the immune response. The level of immunological tolerance to different portions of any particular protein may depend upon whether the expressed portion of the protein resides within or without the cell membrane. The ECD resides on the cell surface and is shed. By contrast, the ICD and portions thereof reside inside the cell and are not shed. The ECD readily comes in contact with the body's immune system, whereas the ICD and portions thereof are relatively sequestered from the body's immune system. As a result, the level of immunologic tolerance to ECD is greater than that to the ICD portion of the Her-2/neu protein. Thus, for vaccines according to the present invention, ICD protein and ICD peptides, and variants thereof, including PD proteins and PD peptides, induce a relatively greater level of immune response than ECD protein and ECD peptides.

Although the ICD and its variants are more immunogenic than the ECD and its variants, antibodies to ECD and its variants are beneficial and possibly desirable. The ECD resides at the cell surface whereas the ICD and portions thereof are not secreted and are sequestered inside the cell. Thus, antibody responses to the ECD can have a greater therapeutic benefit, and accordingly are preferred according to the invention. The ECD by itself is not very immunogenic. Since the ICD (including the PD and Δ PD) is more immunogenic than the ECD, an ECD-ICD fusion protein and/or an ECD-PD fusion protein are more immunogenic than the ECD alone. The ECD-ICD and/or the ECD-PD

fusion proteins are expected to be more effective for inducing antibodies to the ECD than the ECD alone, and are a preferred embodiment of the invention.

5 In the present invention, the ECD or its variants is combined, linked or fused (either directly or indirectly) with the ICD or its variants, preferably with the PD or its variants. The ECD provides the structural conformation for inducing antibodies that react with Her-2/neu protein at the cell surface, while the ICD or PD increases the immunogenicity of the ECD. The combination is surprisingly more effective for inducing an immune response to the ECD than the ECD alone.

10 The ECD, or portions thereof, can be combined with the ICD or its variants, including portions of the ICD or with the PD or its variants, including portions of the PD (*e.g.*, the Δ PD or the PD of the Delta Her-2/neu protein). The ECD of the present invention is preferably a human, a rat or a mouse ECD. The human ECD is set forth in Fig. 9 and as SEQ ID NO:3. The rat ECD is set forth in Fig. 14 and as SEQ ID NO:8.

15 The ICD of the present invention is preferably a human, a rat or a mouse ICD. The human ICD is set forth in Fig. 7 and SEQ ID NO:1 as inclusively spanning the region of Lys 676 to Val 1255. The rat ICD is set forth in Fig. 8 and SEQ ID NO:2 as inclusively spanning the region of Lys 677 to Val 1256.

20 The PD of the present invention is preferably a human, a rat or a mouse PD. The human PD is set forth in Fig. 10 and as SEQ ID NO:4. The human PD may be the human Δ PD, which is set forth in Fig. 11 and as SEQ ID NO:5. The rat PD is shown in Fig. 8 and SEQ ID NO:2 as inclusively spanning the region of Gln 991 to Val 1256. The rat PD may be the rat Δ PD, which is shown in Fig. 8 and SEQ ID NO:2 as inclusively spanning the region of Gln 991 to Arg 1049.

25 In one embodiment, a human ECD can be fused with either (i) a human ICD or a rat ICD or (ii) a human PD or Δ PD, or a rat PD or Δ PD. In another embodiment, a rat ECD can be fused with either (i) a human ICD or a rat ICD or (ii) a human PD or Δ PD, or a rat PD or Δ PD. In yet another embodiment, a mouse ECD can be fused with a mouse PD.

30 The Her-2/neu PD is 268 amino acids in length, is intracellular, and can be phosphorylated by protein tyrosine kinases. This region shares no identity with the corresponding part of other tyrosine kinase receptors. Thus, the specificity and uniqueness of this domain makes it particularly preferred for use as a tumor vaccine.

However, the expression of this domain alone in bacteria and mammalian cells is problematic. For example, the resultant PD protein is very labile and is not appropriate for large scale production. In one embodiment, this invention has solved such problems by fusing all or part of the intracellular domain or the phosphorylation domain to all or part of the Her-2/neu extracellular domain. The ECD-ICD fusion proteins and the ECD-PD fusion proteins of the invention are soluble, are secreted and are stable in culture media. This system can provide large quantities of intracellular domain or phosphorylation domain protein for cancer vaccine development, preferably breast cancer vaccine development, but will be useful for vaccines against any cancer characterized by Her-2/neu expression. In addition to allowing increased expression of the intracellular domain or phosphorylation domain, or variants thereof, as a fusion protein with the extracellular domain or its variants, the ECD-ICD and ECD-PD fusion proteins provide for an improved vaccine formulation.

The PD was secreted by introducing a secretion signal sequence preceding the N-terminus of the PD to yield a soluble, secreted recombinant protein. The secretion process is preferred since the recombinant proteins accumulate in the culture media. Because the protein is not associated with the intracellular proteins, proteolysis is limited. The protein can thus be purified more easily and economically.

As described in Example 3, the pFLAGCMV-1 expression plasmid (Kodak) was used to determine which region of the Her-2/neu intracellular domain was able to be secreted. The proteins were expressed as fusion proteins with a preprotrypsin secretion signal and FLAG-Tag at their N-terminus. HEK-293 cells were transfected with such constructs, and cell and culture media were assayed for FLAG-Tag fusion proteins by western blot with FLAG-Tag M2 antibody as probe. The results in Fig. 4 demonstrate that neither full length ICD nor KD was secreted, but that the PD was soluble and was secreted and detected in the culture media. The results indicate that the full length structure ICD or KD protein did not result in secretion, that is in passage of the protein through the cell membrane.

As described in Example 4, since the ECD has a secretion signal sequence and can be expressed well as a secreted protein, the ECD was used as a fusion partner for PD. The ECD-PD fusion protein was expressed in HEK-293 cells. The secretion of soluble ECD-PD fusion protein was determined by ELISA assay with Her-2/neu ECD-specific antibodies, followed by western blot with Her-2/neu PD-specific antibodies. As

shown in Fig. 6, the soluble ECD-PD expressed in HEK-293 and was secreted into the culture medium.

B. Immunogenicity of the Fusion Proteins of the Invention

In a preferred embodiment, the present invention is directed to a fusion protein based on particular portions (*e.g.*, Her-2/neu ECD-ICD fusion protein or Her-2/neu ECD-PD fusion protein) of the protein expression product of the Her-2/neu gene, which is capable of eliciting an antibody response and can be recognized by thymus-dependent lymphocytes ("T cells"). Accordingly, the autochthonous immune T cell response can be used prophylactically or to treat malignancies in which Her-2/neu is or has been overexpressed. In another aspect, the present invention is directed to the use of nucleic acid molecules directing the expression of such ECD-ICD fusion proteins or ECD-PD fusion proteins, or their variants, alone or in a viral vector for immunization.

In general, CD4⁺ T cell populations are considered to function as helpers or inducers through the release of lymphokines when stimulated by a specific antigen; however, a subset of CD4⁺ cells can act as cytotoxic T lymphocytes (CTL). Similarly, CD8⁺ T cells are considered to function by directly lysing antigenic targets; however, under a variety of circumstances they can secrete lymphokines to provide helper or DTH function. Despite the potential of overlapping function, the phenotypic CD4 and CD8 markers are linked to the recognition of peptides bound to class I or class II MHC antigens. The recognition of an antigen in the context of class I or class II MHC mandates that CD4⁺ and CD8⁺ T cells respond to different antigens or to the same antigen presented under different circumstances. The binding of immunogenic peptides to class II MHC antigens most commonly occurs for antigens ingested by antigen presenting cells.

As disclosed within the present invention, an ECD-ICD fusion protein or an ECD-PD fusion protein of the protein expression product of the Her-2/neu oncogene is recognized by T cells. Circulating Her-2/neu ECD-ICD fusion protein or Her-2/neu ECD-PD fusion protein is degraded to peptide fragments. Peptide fragments from the ECD-ICD fusion protein or ECD-PD fusion protein bind to major histocompatibility complex (MHC) antigens. By display of a peptide bound to MHC antigen on the cell surface and recognition by host T cells of the combination of peptide plus self MHC antigen, the Her-2/neu ECD-ICD or the Her-2/neu ECD-PD fusion protein (including that expressed on a malignant cell) are immunogenic to T cells. The exquisite specificity of

the T cell receptor enables individual T cells to discriminate between protein fragments which differ by a single amino acid residue.

During the immune response to a peptide fragment from an ECD-ICD or an ECD-PD fusion protein, T cells expressing a T cell receptor with high affinity binding of the peptide-MHC complex binds to the peptide-MHC complex and thereby become activated and induced to proliferate. In the first encounter with a peptide, small numbers of immune T cells will secrete lymphokines, proliferate and differentiate into effector and memory T cells. The primary immune response will occur *in vivo* but has been difficult to detect *in vitro*. A subsequent encounter with the same antigen by the memory T cells will lead to a faster and more intense immune response. The secondary response will occur either *in vivo* or *in vitro*. The *in vitro* response is easily gauged by measuring the degree of proliferation, the degree of cytokine production, or the generation of cytolytic activity of the T cell population re-exposed to the antigen. Substantial proliferation of the T cell population in response to a particular antigen is considered to be indicative of prior exposure to or priming with the antigen.

C. Fusion Proteins of the Invention

In one embodiment, the compounds of the present invention comprise Her-2/neu ECD-ICD fusion proteins or variants, or polynucleotides that encode such ECD-ICD fusion proteins. Preferably, the nucleic acid molecules are DNA molecules. In the Her-2/neu ECD-ICD fusion proteins of the invention, the ECD and ICD polypeptide components can be directly fused or fused via a linker, *e.g.*, an amino acid linker or another type of chemical linker. In a preferred embodiment, the ECD-ICD fusion proteins of the present invention comprise all or a portion of the Her-2/neu ECD fused directly to all or a portion of the Her-2/neu ICD.

In additional embodiments, the size of the ECD in the ECD-ICD fusion protein can be altered by sequentially removing anywhere from 1 to about 100 amino acids from the carboxyl terminus of the ECD, preferably about 100 amino acids. Similarly, the size of the ICD in the ECD-ICD fusion protein can be altered by sequentially removing anywhere from 1 to about 100 amino acids from the N-terminus and/or the C-terminus of the ICD. The resulting variant forms may be selected on the basis of their antigenicity and/or immunogenicity using appropriate screening methods as described in the literature and herein, for use according to the invention.

In another embodiment, the compounds of the present invention comprise Her-2/neu ECD-PD fusion proteins, or variants, or nucleic acid molecules that encode such ECD-PD fusion proteins. In one embodiment, the Her-2/neu ECD is fused to the Her-2/neu Δ PD. Preferably, the nucleic acid molecules are DNA molecules. In the Her-2/neu ECD-PD fusion proteins of the invention, the ECD and PD or Δ PD polypeptide components can be directly fused or fused via a linker, *e.g.*, a peptide linker. In a preferred embodiment, the ECD-PD fusion protein of the present invention comprises the Her-2/neu ECD fused directly to the Her-2/neu PD or to the Her-2/neu Δ PD. Here and throughout the specification, a preferred embodiment of the fusion proteins of the invention is the Her-2/neu PD fusion protein.

In another embodiment, the size of the ECD in the ECD-PD fusion protein is altered by sequentially removing anywhere from 1 to about 100 amino acids from the carboxyl terminus of the ECD, preferably about 100 amino acids. Similarly, the size of the PD in the ECD-PD fusion protein can be altered by sequentially removing amino acids from the N-terminus of the PD. Again, the preferred embodiment is PD. Other variant forms may be selected on the basis of their antigenicity and/or immunogenicity using appropriate screening methods as described in the literature and herein, for use according to the invention.

Table 1 below shows that removing 100 amino acids from the carboxyl terminus of the ECD does not have any impact on the expression level and stability of the ECD-PD fusion protein.

Table 1: ECD_{truncated}-PD Summary

Clone I.D.	area of deletion (bp/aa)	# of bp	# of aa	relative expression
A5	1655-1882 / 552-628	228	76	***
B4	1660-1866 / 554-622	207	69	**
B9	1595-1891 / 532-631	297	99	***
C2	1681-1902 / 561-634	222	74	*
C7	1612-1902 / 538-634	291	97	*****
F10	1634-1951 / 545-651	318	106	*****
ECD-PD WT	-	-	-	*

Variants of the ECD-ICD fusion protein and the ECD-PD fusion protein also include various structural forms of the native ECD-ICD fusion protein and ECD-PD fusion protein, respectively. Due to the presence of ionizable amino and carboxyl groups, for example, a Her-2/neu ECD-ICD or ECD-PD fusion protein can be in the form of an
5 acidic or basic salt, or may be in a neutral form. Individual amino acid residues may also be modified by oxidation or reduction.

Other variants within the scope of the invention include ECD-ICD fusion proteins or ECD-PD fusion proteins in which the primary amino acid structure native Her-2/neu ECD-ICD protein or native Her-2/neu ECD-PD protein, respectively, is
10 modified by forming covalent or aggregative conjugates with other peptides or polypeptides, or chemical moieties such as glycosyl groups, lipids, phosphate, acetyl groups and the like. Covalent derivatives may be prepared, for example, by linking particular functional groups to amino acid side chains or at the N- or C-terminus. In some embodiments, the fusion proteins of the invention further comprise a TcP0 peptide
15 motif (SEQ ID NO:31).

The present invention also includes Her-2/neu ECD-ICD fusion proteins and Her-2/neu ECD-PD fusion proteins with or without glycosylation. ECD-ICD fusion proteins and ECD-PD fusion proteins expressed in yeast or mammalian expression systems may be similar to, or slightly different in molecular weight and glycosylation
20 pattern from, the native molecules, depending upon the expression system. Expression of DNA encoding polypeptides in bacteria such as *E. coli* typically provides non-glycosylated molecules. N-glycosylation sites of eukaryotic proteins are characterized by the amino acid triplet Asn-A₁-Z, wherein A₁ is any amino acid except Pro, and Z is Ser or Thr. Variants of Her-2/neu ECD-ICD or ECD-PD fusion proteins having inactivated N-
25 glycosylation sites can be produced by techniques known to one skilled in the art, such as oligonucleotide synthesis and ligation or site-specific mutagenesis techniques, and are within the scope of the invention. Alternatively, N-linked glycosylation sites can be added to a Her-2/neu ECD-ICD or ECD-PD fusion protein.

The ECD-ICD fusion proteins of the present invention, which will be
30 understood to include variants, include any possible combination between human and non-human polypeptides. Non-human polypeptides comprise polypeptides from any mammal, such as, *e.g.*, rat, mouse, guinea pig, horse, cow, pig, sheep, dog, *etc.* In one embodiment, the ECD-ICD fusion proteins include:

(i) human ECD - human ICD fusion proteins, such as those formed by linking the human ECD of Fig. 9 (SEQ ID NO:3) with the human ICD, which is the amino acid sequence inclusively spanning Lys 676 to Val 1255, as shown in Fig. 7 (SEQ ID NO:1), with or without a chemical and/or amino acid linking group, and variants thereof;

(ii) rat ECD - rat ICD fusion proteins, such as those formed by linking the rat ECD of Fig. 14 (SEQ ID NO:8) with the rat ICD, which is the amino acid sequence inclusively spanning Lys 677 to Val 1256, as shown in Fig. 8 (SEQ ID NO:2), with or without a chemical and/or amino acid linking group, and variants thereof;

(iii) human ECD - rat ICD fusion proteins, such as those formed by linking the human ECD shown in Fig. 9 (SEQ ID NO:3) with the rat ICD, which is the amino acid sequence inclusively spanning Lys 677 to Val 1256, as shown in Fig. 8 (SEQ ID NO:2), with or without a chemical and/or amino acid linking group, and variants thereof; and

(iv) rat ECD - human ICD fusion proteins, such as those formed by linking the rat ECD, as shown in Fig. 14 (SEQ ID NO:8), with the human ICD, which is the amino acid sequence inclusively spanning Lys 676 to Val 1255, as shown in Fig. 7 (SEQ ID NO:1), with or without a chemical and/or amino acid linking group, and variants thereof.

Any variants of the ECD-ICD fusion proteins of the present invention are included as embodiments of the present invention. In one embodiment, such variants are substantially identical or substantially similar to the native Her-2/neu ECD-ICD protein and retain the ability to stimulate an immune response. Human DNA sequences that encode the ECD protein are shown, for example, in Fig. 15 (SEQ ID NO:9) as inclusively spanning nucleotide 1 to nucleotide 1959. Human DNA sequences that encode the ICD protein are shown, for example, in Fig. 15 (SEQ ID NO:9) as inclusively spanning nucleotide 2026 to nucleotide 3765. The effect of any sequence modification on the ability of a Her-2/neu ECD-ICD protein to produce an immune response may be readily determined, for example, by analyzing the ability of the mutated Her-2/neu ECD-ICD protein to induce a T cell response using, for example, the methods described herein, or by analyzing the ability of the mutated Her-2/neu ECD-ICD protein to produce antibodies.

The ECD-PD fusion proteins of the present invention, which will be understood to include variants, include any possible combination between human and

non-human polypeptides. Non-human polypeptides comprise, *e.g.*, rat, mouse, guinea pig, horse, cow, pig, sheep, dog, *etc.* In one embodiment, the ECD-PD fusion proteins include:

- (i) human ECD - human PD fusion proteins, such as shown in Fig. 12
5 (SEQ ID NO:6) and variants thereof, including fusion proteins formed by linking the human ECD of Fig. 9 (SEQ ID NO:3) with the human PD of Fig. 10 (SEQ ID NO:4) with or without a chemical and/or amino acid linking group, and variants thereof;
- (ii) rat ECD - rat PD fusion proteins, such as those formed by linking the
10 rat ECD of Fig. 14 (SEQ ID NO:8) with the rat PD, which is the amino acid sequence inclusively spanning Gln 991 to Val 1256, as shown in Fig. 8 (SEQ ID NO:2), with or without a chemical and/or amino acid linking group, and variants thereof;
- (iii) human ECD - rat PD fusion proteins, such as those formed by linking
15 the human ECD shown in Fig. 9 (SEQ ID NO:3) with the rat PD, which is the amino acid sequence inclusively spanning Gln 991 to Val 1256, as shown in Fig. 8 (SEQ ID NO:2), with or without a chemical and/or amino acid linking group, and variants thereof;
- (iv) rat ECD - human PD fusion proteins, such as those formed by linking
the rat ECD, as shown in Fig. 14 (SEQ ID NO:8), with the human PD, as shown in Fig. 10 (SEQ ID NO:4), with or without a chemical and/or amino acid linking group, and variants thereof.
- (v) mouse ECD - mouse PD fusion proteins, such as those shown in Fig.
20 22 (SEQ ID NO:27).

Any variants of the ECD-PD fusion proteins of the present invention are included as embodiments of the present invention. In one embodiment, such variants are substantially identical or substantially similar to the native Her-2/neu ECD-PD protein
25 and retain the ability to stimulate an immune response. Human DNA sequences that encode the ECD protein are shown, for example, in Fig. 15 (SEQ ID NO:9) as inclusively spanning nucleotide 1 to nucleotide 1959. Human DNA sequences that encode the PD protein are shown, for example, in Fig. 15 (SEQ ID NO:9) as inclusively spanning nucleotide 2968 to nucleotide 3765. Mouse DNA sequences that encode the mouse ECD-
30 PD fusion proteins are shown in Fig. 23 (SEQ ID NO:28). The effect of any sequence modification on the ability of a Her-2/neu ECD-PD protein to produce an immune response may be readily determined, for example, by analyzing the ability of the mutated Her-2/neu ECD-PD protein to induce a T cell response using, for example, the methods

described herein, or by analyzing the ability of the mutated Her-2/neu ECD-PD protein to produce antibodies.

In another embodiment, the ECD-PD fusion proteins are ECD-APD fusion proteins of the present invention, which will be understood to include variants, including any possible combination between human and non-human polypeptides. Non-human polypeptides comprise, *e.g.*, rat, mouse, guinea pig, horse, cow, pig, sheep, dog, *etc.* In one embodiment, the ECD-APD fusion proteins include:

(i) human ECD - human APD fusion proteins, such as shown in Fig. 13 (SEQ ID NO:7) and variants thereof, including fusion proteins formed by linking the human ECD of Fig. 9 (SEQ ID NO:3) with the human APD of Fig. 11 (SEQ ID NO:5) with or without a chemical and/or amino acid linking group, and variants thereof;

(ii) rat ECD - rat APD fusion proteins, such as those formed by linking the rat ECD of Fig. 14 (SEQ ID NO:8) with the rat APD, which is the amino acid sequence inclusively spanning Gln 991 to Arg 1049, as shown in Fig. 8 (SEQ ID NO:2), with or without a chemical and/or amino acid linking group, and variants thereof;

(iii) human ECD - rat APD fusion proteins, such as those formed by linking the human ECD shown in Fig. 9 (SEQ ID NO:3) with the rat APD, which is the amino acid sequence inclusively spanning Gln 991 to Arg 1049, as shown in Fig. 8 (SEQ ID NO:2), with or without a chemical and/or amino acid linking group, and variants thereof; and

(iv) rat ECD - human APD fusion proteins, such as those formed by linking the rat ECD, as shown in Fig. 14 (SEQ ID NO:8), with the human APD, as shown in Fig. 11 (SEQ ID NO:5), with or without a chemical and/or amino acid linking group, and variants thereof.

Any variants of the ECD-APD fusion proteins of the present invention are included as embodiments of the present invention. In one embodiment, such variants are substantially identical or substantially similar to the native Her-2/neu ECD-APD protein and retain the ability to stimulate an immune response. Human DNA sequences that encode the ECD protein are shown, for example, in Fig. 15 (SEQ ID NO:9) as inclusively spanning nucleotide 1 to nucleotide 1959. Human DNA sequences that encode the APD protein are shown, for example, in Fig. 15 (SEQ ID NO:9) as inclusively spanning nucleotide 2968 to nucleotide 3144. The effect of any sequence modification on the ability of a Her-2/neu ECD-APD protein to produce an immune response may be readily

determined, for example, by analyzing the ability of the mutated Her-2/neu ECD-APD protein to induce a T cell response using, for example, the methods described herein, or by analyzing the ability of the mutated Her-2/neu ECD-APD protein to produce antibodies.

5 In a preferred embodiment, the Her-2/neu ECD-PD fusion proteins of the invention are ECD-PD fusion proteins.

 Within certain specific embodiments, the fusion proteins of the invention may comprise a fusion partner, such as, *e.g.*, an immunological fusion partner or an expression enhancer. A fusion partner may, for example, assist in providing T helper
10 epitopes (an immunological fusion partner), preferably T helper epitopes recognized by humans, or may assist in expressing the fusion protein (an expression enhancer) at higher yields than the recombinant fusion protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the fusion protein or to enable the fusion
15 protein to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags, which facilitate the purification of the fusion protein of interest.

 Also provided are fusion proteins that comprise a fusion polypeptide as described herein together with an unrelated immunogenic protein. Preferably, the immunogenic protein is capable of eliciting a recall response. Examples of such proteins
20 include tetanus, tuberculosis and hepatitis proteins (*see, e.g.*, Stoute *et al.* (1997) *New Engl. J. Med.* 336:86-91).

 In other embodiments, an immunological fusion partner is derived from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza* B (WO 91/18926). Preferably, a protein D derivative comprises approximately the first
25 third of the protein (*e.g.*, the first N-terminal 100-110 amino acids), and the protein D derivative may be lipidated. Within certain preferred embodiments, the first 109 residues of a Lipoprotein D fusion partner are included on the N-terminus to provide the fusion protein with additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal
30 presentation of the fusion protein to antigen presenting cells. Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin), or an immunogenic portion thereof (*see, e.g.*, WO 99/40188 and WO 93/04175). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper

epitopes may be used. In one embodiment, the fusion proteins of the present invention further comprise the NS1 fusion partner or an immunogenic fragment thereof, preferably attached to the N-terminal region of the fusion protein. The fusion proteins comprising the NS1 partner are preferably recombinantly expressed in BHK cells, as described *infra*.

5 In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292 (1986)). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone.

10 The C-terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA fragment at the amino terminus has been described (*see, Biotechnology* 10:795-798 (1992)). Within a preferred

15 embodiment, a repeat portion of LYTA may be incorporated into a fusion protein. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

In a preferred embodiment, the fusion proteins of the invention further comprise a TcP0 peptide motif, which enhances the immunogenicity of the fusion protein.

20 TcP0 is a 6 amino acid peptide from a ribosomal P protein that has been shown to induce epitope spreading. The TcP0 motif can be attached to the fusion protein at different sites, but is preferably fused to the C-terminus of the fusion protein. The amino acid sequence of TcP0 is shown in Fig. 26 (SEQ ID NO:31), and the nucleic acid sequence encoding TcP0 is shown in Fig. 27 (SEQ ID NO:32). Recombinant nucleic acids encoding a fusion

25 protein comprising the TcP0 motif can be obtained using standard techniques or as described *infra*. Recombinant nucleic acids are constructed so that, preferably, the TcP0 polynucleotide sequence is located 3' to the fusion protein of interest.

In another preferred embodiment, the fusion proteins of the present invention further comprise a fusion partner. Preferred fusion partners include, but are not

30 limited to, *e.g.*, Ra12 or LeIF. In particular, the invention provides materials and methods for using a Ra12 or a LeIF sequence as a fusion partner to facilitate the stable and high yield expression of recombinant fusion polypeptides or break tolerance.

Ra12 is a 14 kD C-terminal fragment of the *M. tuberculosis* MTB32A coding sequence, which is expressed at high levels on its own and remains soluble

throughout the purification process. LeIF is a *Leishmania* antigen which is homologous to the eukaryotic ribosomal protein eIF and which is capable of stimulating a Th1 and/or CTL immune response (*see, e.g.*, U.S. Patent Nos. 5,876,966, 5,876,735, and 5,879,687). The present invention utilizes these properties of the Ra12 and LeIF polypeptides and provides recombinant nucleic acid molecules, expression vectors, host cells, and methods for stable and high yield expression of fusion polypeptides comprising, in addition to the Her-2/neu fusion polypeptides, a Ra12 or LeIF polypeptide. Recombinant nucleic acids, which encode a fusion polypeptide comprising a Ra12 or LeIF polypeptide and a fusion protein of interest, can be readily constructed by conventional genetic engineering techniques. Recombinant nucleic acids are constructed so that, preferably, the fusion partner polynucleotide sequence is located 5' to a selected fusion protein sequence. It may also be appropriate to place an fusion partner polynucleotide sequence 3' to the polynucleotide sequence of the fusion protein of interest or to insert the polynucleotide sequence of the fusion protein into a site within an fusion partner polynucleotide sequence. In addition, any suitable polynucleotide that encodes a fusion partner or a portion or other variant thereof as described herein can be used in constructing recombinant fusion nucleic acids of the present invention.

Nucleic acids encoding the fusion partner polypeptides of this invention can be prepared by any suitable method known in the art. Exemplary methods include cloning and restriction of appropriate sequences or direct chemical synthesis by methods such as the phosphotriester method of Narang *et al.* (1979) *Meth. Enzymol.* 68:90-99; the phosphodiester method of Brown *et al.* (1979) *Meth. Enzymol.* 68:109-151; the diethylphosphoramidite method of Beaucage *et al.* (1981) *Tetra. Lett.* 22:1859-1862; and the solid support method of U.S. Patent No. 4,458,066.

Recombinant nucleic acids that encode a fusion polypeptide comprising a fusion partner polypeptide and a selected fusion protein can be prepared using any methods known in the art. As described above, recombinant nucleic acids are constructed so that, preferably, the fusion partner polynucleotide sequence is located 5' to the polynucleotide sequence encoding the fusion protein of interest. The fusion partner and fusion protein polynucleotide sequences can also be modified to facilitate their fusion and subsequent expression.

The recombinant nucleic acids can further comprise other nucleotide sequences such as sequences that encode affinity tags to facilitate protein purification protocol.

D. Variants of the Fusion Proteins of the Invention

CD4⁺ T cells generally recognize antigens that have been external to the tumor cells. By contrast, under normal circumstances, binding of peptides to class I MHC occurs only for proteins present in the cytosol and synthesized by the target itself, proteins in the external environment are excluded. An exception to this is the binding of exogenous peptides with a precise class I binding motif which are present outside the cell in high concentration. Thus, CD4⁺ and CD8⁺ T cells have broadly different functions and tend to recognize different antigens as a reflection of where the antigens normally reside.

Another way to make amino acid substitutions to produce variants of the present invention is to identify and replace amino acids in T cell motifs with potential to bind to class II MHC molecules (for CD4⁺ T cell response) or class I MHC molecules (for CD8⁺ T cell response). Peptide segments (of a Her-2/neu ECD-ICD or ECD-PD fusion protein) with a motif with theoretical potential to bind to class II MHC molecules may be identified by computer analysis. For example, a protein sequence analysis package, *T Sites*, that incorporates several computer algorithms designed to distinguish potential sites for T cell recognition can be used (Feller *et al.* (1991) *Nature* 349:720-721). Two searching algorithms are used: (1) the AMPHI algorithm described by Margalit (Feller *et al.* (1991) *Nature* 349:720-721; Margalit *et al.* (1987) *J. Immunol.* 138:2213-2229) identifies epitope motifs according to alpha-helical periodicity and amphipathicity; (2) the Rothbard and Taylor algorithm identifies epitope motifs according to charge and polarity pattern (Rothbard *et al.* (1988) *EMBO J.* 7:93-100). Segments with both motifs are most appropriate for binding to class II MHC molecules. CD8⁺ T cells recognize peptides bound to class I MHC molecules. Parker *et al.* (1994) *J. Immunol.* 152:163 have determined that peptides binding to particular MHC molecules share discernible sequence motifs. A peptide motif for binding in the groove of HLA-A2.1 has been defined by Edman degradation of peptides stripped from HLA-A2.1 molecules of a cultured cell line (Table 2, from Falk *et al.* (1991) *Nature* 351:290-296). The method identified the typical or average HLA-A2.1 binding peptide as being 9 amino acids in length with dominant anchor residues occurring at positions 2 (L) and 9 (V). Commonly occurring strong binding residues have been identified at positions 2 (M), 4 (E,K), 6 (V), and 8 (K). The identified motif represents the average of many binding peptides.

Table 2: The HLA-A2.1 Restricted Motif

	Amino Acid Position									Point Assignment
	1	2	3	4	5	6	7	8	9	
Dominant Binding Anchor Residue		L							V	+3
Strong Binding Residue		M		E		V		K		+2
Weak Binding Residue	I		A	G	I	I	A	E	L	+1
	L		Y	P	K	L	Y	S		
	F		F	D	Y	T	H			
	K		P	T	N					
	M		M		G					
	Y		S		V					
					H					

The derived peptide motif as currently defined is not particularly stringent. Some HLA-A2.1 binding peptides do not contain both dominant anchor residues and the amino acids flanking the dominant anchor residues play major roles in allowing or disallowing binding. Not every peptide with the current described binding motif will bind, and some peptides without the motif will bind. However, the current motif is valid enough to allow identification of some peptides capable of binding. Of note, all MHC molecules and respective motifs place 6 amino acids between the dominant anchor amino acids at residues 2 and 9.

Following identification of peptide motifs within Her-2/neu ECD-ICD or ECD-PD fusion proteins, amino acid substitutions can be made conservatively or non-conservatively. The latter type of substitutions are intended to produce an improved ECD-ICD or ECD-PD fusion protein or polypeptide that is more potent and/or more broadly cross-reactive. An example of a more potent protein or peptide is one that binds with higher affinity to the same MHC molecule as the natural protein or polypeptide, without affecting recognition by T cells specific for natural protein or polypeptide. An example of a polypeptide with broader cross-reactivity is one that induces more broadly cross-reactive immune responses (*i.e.*, binds to a greater range of MHC molecules) than

natural polypeptide. Similarly, one or more amino acids residing between peptide motifs and having a spacer function (*e.g.*, do not interact with a MHC molecule or T cell receptor) can be substituted conservatively or non-conservatively. It will be evident to one of ordinary skill in the art that polypeptides containing one or more amino acid
5 substitutions can be tested for beneficial or adverse immunological interactions by a variety of assays, including those described herein for the ability to stimulate T cell recognition.

Variants within the scope of this invention can also, or alternatively, contain other modifications, including the deletion or addition of amino acids, that have
10 minimal influence on the desired immunological properties of the polypeptide, as described *supra*. It will be appreciated by one of ordinary skill in the art that truncated forms or non-native extended forms of a Her-2/neu ECD-ICD or ECD-PD fusion protein can be used, provided the desired immunological properties are at least roughly equivalent to that of full length, native Her-2/neu ECD-ICD or ECD-PD fusion protein.

15 Cysteine residues may be deleted or replaced with other amino acids to prevent formation of incorrect intramolecular disulfide bridges upon renaturation. Other approaches to mutagenesis involve modification of adjacent dibasic amino acid residues to enhance expression in yeast systems in which KBX2 protease activity is present.

IV. PREPARING THE FUSION PROTEINS OF THE INVENTION

20 A. Polynucleotides Encoding Fusion Proteins

In one embodiment, the present invention relates to isolated or purified polynucleotides that encode the Her-2/neu fusion proteins. In accordance with the invention, any nucleotide sequence which encodes the amino acid sequence of the fusion protein of interest can be used to generate recombinant molecules which direct the
25 expression of the fusion protein.

In order to clone full-length coding sequences or homologous variants to generate the Her-2/neu fusion polynucleotides, labeled DNA probes designed from any portion of the Her-2/neu nucleotide sequences or their complements may be used to screen a genomic or cDNA library, to identify the coding sequence of each individual
30 component of the fusion protein. The Her-2/neu nucleotide sequences can be from any suitable mammal, *e.g.*, human, rat, mouse, horse, cow, pig, sheep, dog, *etc.*

In one embodiment, the Her-2/neu sequence is from a human, a rat, or a mouse. The nucleotide sequence of the mouse Her-2/neu gene is shown in Figure 19

(SEQ ID NO:11). Mouse Her-2/neu can also be amplified from mouse brain RNA using the following primers: 5' primer: CCATGGAGCTGGCGGCCTGGTGCCGTTG (SEQ ID NO:12) and 3' primer: GGCCTTCTGGTTCATACTGGCACATCCAGGC (SEQ ID NO:13). The mouse Her-2/neu amino acid sequence is shown in Figure 20 (SEQ ID NO:14). A variant amino acid sequence for mouse Her-2/neu is described in Nagata *et al.* (1997) *J. Immunol.* 159:1336-1343.

Such clones may be isolated by screening an appropriate expression library for clones that express a full length Her-2/neu protein. The library preparation and screen may generally be performed using methods known to one of ordinary skill in the art, such as methods described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring Harbor, NY (1989). Briefly, a bacteriophage expression library may be plated and transferred to filters. The filters may then be incubated with a detection reagent. In the context of this invention, a "detection reagent" is any compound capable of binding to the Her-2/neu protein, which may then be detected by any of a variety of means known to one of ordinary skill in the art. Typical detection reagents contain a "binding agent," such as Protein A, Protein G, IgG or a lectin, coupled to a reporter group. Preferred reporter groups include enzymes, substrates, cofactors, inhibitors, dyes, radionuclides, luminescent groups, fluorescent groups and biotin. More preferably, the reporter group is horseradish peroxidase, which may be detected by incubation with a substrate such as tetramethylbenzidine or 2,2'-azino-di-3-ethylbenz-thiazoline sulfonic acid. Plaques containing genomic or cDNA sequences that express Her-2/neu protein are isolated and purified by techniques known to one of ordinary skill in the art. Appropriate methods may be found, for example, in Sambrook *et al.*, *supra*.

Isolation of coding sequences may also be carried out by the polymerase chain reaction (PCR) using two degenerate oligonucleotide primer pools designed on the basis of the coding sequences disclosed herein. The desired nucleic acids can also be cloned using other well known amplification techniques. Examples of protocols sufficient to direct persons of skill through *in vitro* amplification methods, including PCR, ligase chain reaction (LCR), Q β -replicase amplification and other RNA polymerase mediated techniques are found in Sambrook *et al.*, *supra*, and Ausubel *et al.* *Current Protocols in Molecular Biology* (1994), as well as in U.S. Patent No. 4,683,202; *PCR Protocols A Guide to Methods and Applications* (Innis *et al.* eds. 1990); Arnheim & Levinson *C&EN* pp. 36-47 (October 1, 1990); *The Journal of NIH Research* 3:81-94

(1991); Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:1173; Guatelli *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:1874; Lomell *et al.* (1989) *J. Clin. Chem.* 35:1826; Landegren *et al.* (1988) *Science* 241:1077-1080; Van Brunt (1990) *Biotechnology* 8:291-294; Wu *et al.* (1989) *Gene* 4:560; and Barringer *et al.* (1990) *Gene* 89:117. Improved
5 methods of cloning *in vitro* amplified nucleic acids are described in U.S. Patent No. 5,426,039. Suitable primers for use in the amplification of the nucleic acids of the invention can be designed based on the sequences provided herein.

In accordance with the invention, a polynucleotide of the invention which encodes a fusion protein, fragment thereof, or functional equivalent thereof may be used
10 to generate recombinant nucleic acid molecules that direct the expression of the fusion protein, fragment thereof, or functional equivalent thereof, in appropriate host cells. The fusion polypeptide products encoded by such polynucleotides may be altered by molecular manipulation of the coding sequence.

Due to the inherent degeneracy of the genetic code, other DNA sequences
15 which encode substantially the same or a functionally equivalent amino acid sequence, may be used in the practice of the invention for the expression of the fusion polypeptides. Such DNA sequences include those which are capable of hybridizing to the coding sequences or their complements disclosed herein under low, moderate or high stringency conditions as described herein.

Altered nucleotide sequences which may be used in accordance with the invention include deletions, additions or substitutions of different nucleotide residues resulting in a sequence that encodes the same or a functionally equivalent gene product. The gene product itself may contain deletions, additions or substitutions of amino acid residues, which result in a silent change thus producing a functionally equivalent
20 antigenic epitope. Such conservative amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine, histidine and arginine; amino acids with uncharged polar head groups having
25 similar hydrophilicity values include the following: glycine, asparagine, glutamine, serine, threonine and tyrosine; and amino acids with nonpolar head groups include alanine, valine, isoleucine, leucine, phenylalanine, proline, methionine and tryptophan.
30

The nucleotide sequences of the invention may be engineered in order to alter the fusion protein coding sequence for a variety of ends, including but not limited to,

alterations which modify processing and expression of the gene product. For example, mutations may be introduced using techniques which are well known in the art, *e.g.*, to insert or delete restriction sites, to alter glycosylation patterns, phosphorylation, to create and/or destroy translation, initiation, and/or termination sequences, or to create variations in coding regions, to facilitate further *in vitro* modification, *etc.* One of skill will recognize many ways of generating alterations in a given nucleic acid construct. Such well-known methods include, *e.g.*, site-directed mutagenesis, PCR amplification using degenerate oligonucleotides, exposure of cells containing the nucleic acid to chemical mutagenic agents or radiation, chemical synthesis of a desired oligonucleotide (*e.g.*, in conjunction with ligation and/or cloning to generate large nucleic acids) and other well-known techniques (*see, e.g.*, Gilman *et al.* (1979) *Gene* 8:81-97; Hutchinson *et al.* (1978) *J. Biol. Chem.* 253:6551; Roberts *et al.* (1987) *Nature* 328: 731-734). Preferably, the manipulations do not destroy immunogenicity of the fusion polypeptides.

In one embodiment of the invention, the coding sequence of a fusion protein may be synthesized in whole or in part, using chemical methods well known in the art (*see, e.g.*, Caruthers *et al.* (1980) *Nuc. Acids Res. Symp. Ser.* 7:215-233; Crea *et al.* (1980) *Nuc. Acids Res.* 9(10):2331; Matteucci *et al.* (1980) *Tetrahedron Letter* 21:719 (1980); and Chow *et al.* (1981) *Nuc. Acids Res.* 9(12):2807-2817).

B. Polypeptide Synthesis

Alternatively, the fusion polypeptide itself can be produced using chemical methods to synthesize an amino acid sequence in whole or in part. For example, peptides can be synthesized by solid phase techniques such as, *e.g.*, the Merrifield solid phase synthesis method, in which amino acids are sequentially added to a growing chain of amino acids (*see*, Merrifield (1963) *J. Am. Chem. Soc.* 85:2149-2146). Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Perkin Elmer Biosystems, Inc. (Foster City, CA), and may generally be operated according to the manufacturer's instructions. The synthesized peptides can then be cleaved from the resin and purified, *e.g.*, by preparative high performance liquid chromatography (*see*, Creighton, *Proteins Structures and Molecular Principles*, pp. 50-60 (1983)). The composition of the synthetic fusion polypeptides may be confirmed by amino acid analysis or sequencing (*e.g.*, the Edman degradation procedure; *see*, Creighton, *Proteins, Structures and Molecular Principles*, pp. 34-49 (1983)).

In addition, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the sequence. Non-classical amino acids include, but are not limited to, the D-isomers of the common amino acids, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, γ -Abu, ϵ -Ahx, 6-amino
5 hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, C α -methyl amino acids, N α -methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D
10 (dextrorotary) or L (levorotary).

C. Linking Groups

In another embodiment, the polypeptides of the fusion protein, e.g., ECD and ICD or ECD and PD, are joined via a linking group. The linking group can be a chemical crosslinking agent, including, for example, succinimidyl-(N-maleimidomethyl)-
15 cyclohexane-1-carboxylate (SMCC). The linking group can also be an additional amino acid sequence(s), including, for example, a polyglycine linking group.

In a specific embodiment, the coding sequences of each polypeptide in the fusion protein are directly joined at their amino- or carboxy-terminus via a peptide bond in any order.

20 Alternatively, an amino acid linker sequence may be employed to separate the first and second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such an amino acid linker sequence is incorporated into the fusion protein using standard techniques well known in the art. Suitable peptide linker sequences may be chosen based on the following factors:
25 (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues. Other near neutral amino acids, such as Thr and Ala, may also be used
30 in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea *et al.* (1985) *Gene* 40:39-46; Murphy *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:8258-8262; and in U.S. Patent Nos. 4,935,233 and 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in

length. Linker sequences are not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

Other chemical linkers include carbohydrate linkers, lipid linkers, fatty acid linkers, polyether linkers, *e.g.*, PEG, *etc.* (*see, e.g.*, Hermanson (1996) *Bioconjugate Techniques*).

D. Additional Polypeptides

As described *supra*, the fusion protein may be linked to one or more additional polypeptides. For example, the fusion polypeptide may be linked to one or more copies of one of the two polypeptides of the fusion protein. Alternatively, the fusion protein may be linked to an additional heterologous polypeptide, such as, *e.g.*, Ra12 or LeIF, or to a peptide motif such as, *e.g.*, TcP0, as described *supra*. The fusion polypeptide may also be fused to an affinity tag for ease of purification upon expression. For example, multiple histidine residues encoded by the tag allow the use of metal chelate affinity chromatography methods for the purification of fusion polypeptides. Other examples of affinity tag molecules include, *e.g.*, Strep-tag, PinPoint, maltose binding protein, glutathione S-transferase, *etc.* (*see, e.g.*, Glick & Pasternak, *Molecular Biotechnology Principles and Applications of Recombinant DNA* (2nd ed. 1999)).

In one embodiment, the fusion polypeptide is optionally linked to a lipid moiety, such as mycolic acid, lipoaribdomanin ("LAMs"), or trehalose derivatives.

As described above, in one embodiment, such a fusion protein is produced by recombinant expression of a nucleic acid encoding the fusion protein. Such a fusion product can be made by ligating the appropriate nucleic acid sequences encoding the desired amino acid sequences to each other by methods known in the art, in the proper coding frame, and expressing the product by methods known in the art. Alternatively, such a product may be made by protein synthetic techniques, *e.g.*, by use of a peptide synthesizer. Coding sequences for other molecules such as a cytokine or an adjuvant can be added to the fusion polynucleotide as well.

E. Sequence Modifications

Variants of the fusion proteins of the invention that retain the ability to stimulate an immune response may generally be identified by modifying the sequence in one or more of the aspects described above and assaying the resulting fusion protein for the ability to stimulate an immune response, *e.g.*, a T cell response or an antibody

response. For example, such assays may generally be performed by contacting T cells with the modified fusion protein and assaying the response. Naturally occurring variants of the individual polypeptide components of the fusion protein may also be isolated by, for example, screening an appropriate cDNA or genomic library with a DNA sequence
5 encoding each individual polypeptide or a variant thereof.

The above-described sequence modifications may be introduced using standard recombinant techniques or by automated synthesis of the modified fusion protein. For example, mutations can be introduced at particular loci by synthesizing oligonucleotides containing a mutant sequence, flanked by restriction sites enabling
10 ligation to fragments of the native sequence. Following ligation, the resulting reconstructed sequence encodes an analogue having the desired amino acid insertion, substitution, or deletion.

Alternatively, oligonucleotide-directed site-specific mutagenesis procedures can be used to provide a gene in which particular codons are altered according
15 to the substitution, deletion, or insertion required. Exemplary methods of making the alterations set forth above are described by Walder *et al.* (1986) *Gene* 42:133; Bauer *et al.* (1985) *Gene* 37:73; Craik (1985) *BioTechniques* January:12-19; Smith *et al.* (1981) *Genetic Engineering: Principles and Methods*, Plenum Press; and U.S. Patent Nos. 4,518,584 and 4,737,462.

20 Mutations in nucleotide sequences constructed for expression of such Her-2/neu fusion proteins must, of course, preserve the reading frame of the coding sequences and preferably will not create complementary regions that could hybridize to produce secondary mRNA structures, such as loops or hairpins, which would adversely affect the translation of the mRNA. Although a mutation site may be predetermined, it is not
25 necessary that the nature of the mutation *per se* be predetermined. For example, in order to select for optimum characteristics of mutants at a given site, random mutagenesis may be conducted at the target codon and the expressed Her-2/neu fusion protein mutants screened for the desired activity. Not all mutations in a nucleotide sequence which encodes a Her-2/neu fusion protein will be expressed in the final product. For example,
30 nucleotide substitutions may be made to enhance expression, primarily to avoid secondary structure loops in the transcribed mRNA (*see, e.g.*, European Patent Application 75,444A), or to provide codons that are more readily translated by the selected host, such as the well-known *E. coli* preference codons for *E. coli* expression.

F. Expression Vectors

The Her-2/neu fusion proteins, and variants thereof, of the present invention, are preferably produced by recombinant DNA methods. Such methods include inserting a DNA sequence encoding a Her-2/neu fusion protein into a recombinant expression vector and expressing the DNA sequence in a recombinant microbial, mammalian, fungal or insect cell expression system under conditions promoting expression and, preferably, secretion of the fusion protein. DNA sequences encoding the Her-2/neu fusion proteins provided by this invention can be assembled from cDNA fragments and short oligonucleotide linkers, or from a series of oligonucleotides, to provide a synthetic gene which is capable of being inserted in a recombinant expression vector and expressed in a recombinant transcriptional unit.

Recombinant expression vectors contain a DNA sequence encoding a Her-2/neu fusion protein operably linked to suitable transcriptional or translational regulatory elements derived from mammalian, fungal, microbial, viral or insect genes. Such regulatory elements include a transcriptional promoter, an optional operator sequence to control transcription, a sequence encoding suitable mRNA ribosomal binding sites, and sequences which control the termination of transcription and translation. An origin of replication and a selectable marker to facilitate recognition of transformants may additionally be incorporated.

DNA regions are "operably linked" when they are functionally related to each other. For example, DNA for a signal peptide (secretory leader) is operably linked to DNA for a polypeptide if it is expressed as a precursor which participates in the secretion of the polypeptide; a promoter is "operably linked" to a coding sequence if it controls the transcription of the sequence; or a ribosome binding site is "operably linked" to a coding sequence if it is positioned so as to permit translation. Generally, "operably linked" means contiguous and, in the case of secretory leaders, in reading frame. DNA sequences encoding Her-2/neu fusion proteins which are to be expressed in a microorganism will preferably contain no introns that could prematurely terminate transcription of DNA into mRNA.

Expression vectors for bacterial use may comprise a selectable marker and a bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, e.g., pKK223-3 (Pharmacia Fine Chemicals, Uppsala, Sweden), pGEM1 (Promega Biotec, Madison, WI), pET28b (Novagen) and pPDM (a

modified pET28b, Corixa). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. *E. coli* is typically transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species (Bolivar *et al.* (1977) *Gene* 2:95). pBR322 contains genes for ampicillin and tetracycline resistance and thus provides simple means for identifying transformed cells. Promoters commonly used in recombinant microbial expression vectors include the β -lactamase (penicillinase) and lactose promoter system (Chang *et al.* (1978) *Nature* 275:615; and Goeddel *et al.* (1979) *Nature* 281:544), the tryptophan (*trp*) promoter system (Goeddel *et al.* (1980) *Nucl. Acids Res.* 8:4057; and European Patent Application 36,776) and the *tac* promoter (Maniatis, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, p.412 (1982)). A particularly useful bacterial expression system uses the phage λ P_L promoter and cI857ts thermolabile repressor. Plasmid vectors available from the American Type Culture Collection which incorporate derivatives of the λ P_L promoter include plasmid pHUB2, resident in *E. coli* strain JMB9 (ATCC 37092) and pPLc28, resident in *E. coli* RR1 (ATCC 53082).

Suitable promoter sequences in yeast vectors include the promoters for alcohol oxidase, metallothionein, 3-phosphoglycerate kinase (Hitzeman *et al.* (1980) *J. Biol. Chem.* 255:2073) or other glycolytic enzymes (Hess *et al.* (1968) *J. Adv. Enzyme Reg.* 7:149; and Holland *et al.* (1978) *Biochem.* 17:4900), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. Suitable vectors and promoters for use in yeast expression are further described in, *e.g.*, European Patent Application No. 73,657.

Preferred yeast vectors can be assembled using DNA sequences from pBR322 for selection and replication in *E. coli* (Amp^r gene and origin of replication) and yeast DNA sequences including a glucose-repressible ADH2 promoter and α -factor secretion leader. The ADH2 promoter has been described by Russell *et al.* (1982) *J. Biol. Chem.* 258:2674 and Beier *et al.* (1982) *Nature* 300:724. The yeast α -factor leader, which directs secretion of heterologous proteins, can be inserted between the promoter and the structural gene to be expressed (*see, e.g.*, Kurjan *et al.* (1982) *Cell* 30:933; and Bitter *et al.* (1984) *Proc. Natl. Acad. Sci. USA* 81:5330. The leader sequence may be

modified to contain, near its 3' end, one or more useful restriction sites to facilitate fusion of the leader sequence to foreign genes.

The transcriptional and translational control sequences in expression vectors to be used in transforming vertebrate cells may be provided by viral sources. For example, commonly used promoters and enhancers are derived, *e.g.*, from polyoma, adenovirus 2, simian virus (SV40), and human cytomegalovirus. DNA sequences derived from the SV40 viral genome, for example, the SV40 origin, early and late promoter, enhancer, splice, and polyadenylation sites may be used to provide the other genetic elements required for expression of a heterologous DNA sequence. The early and late promoters are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication (Fiers *et al.* (1978) *Nature* 273:113). Smaller or larger SV40 fragments may also be used, provided the approximately 250 bp sequence extending from the *Hind* III site toward the *Bgl* II site located in the viral origin of replication is included. Further, viral genomic promoter, control and/or signal sequences may be used, provided such control sequences are compatible with the host cell chosen. Exemplary vectors can be constructed as disclosed by Okayama *et al.* (1983) *Mol. Cell. Biol.* 3:280.

A useful system for stable high level expression of mammalian receptor cDNAs in C127 murine mammary epithelial cells can be constructed substantially as described by Cosman *et al.* (1986) *Mol. Immunol.* 23:935. A suitable eukaryotic vector for expression of the fusion proteins of the invention is pDC406 (McMahan *et al.* (1991) *EMBO J.* 10:2821 which includes regulatory sequences derived from SV40, human immunodeficiency virus (HIV), and Epstein-Barr virus (EBV). Other vectors include pDC409 and pDC410, which are derived from pDC406. pDC410 was derived from pDC406 by substituting the EBV origin of replication with sequences encoding the SV40 large T antigen. pDC409 differs from pDC406 in that a *Bgl* II restriction site outside of the multiple cloning site has been deleted, making the *Bgl* II site within the multiple cloning site unique. Additional suitable mammalian expression vectors include, for example, the pBIB and CR1012 vectors. Any other vector allowing expression of proteins under the direction of the CMV promoter, SV40 early promoter, SV40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, Rous sarcoma virus promoter, polyhedrin promoter, or other promoters shown effective for expression in mammalian cells are also suitable.

In addition to the transcriptional and translational control sequences, some expression systems have markers that provide gene amplification, such as neomycin, thymidine kinase, hygromycin B phosphotransferase, and dihydrofolate reductase.

5 A useful cell line that allows for episomal replication of expression vectors, such as pDC406 and pDC409, which contain the EBV origin of replication, is CV-1/EBNA (ATCC CRL 10478). The CV-1/EBNA cell line was derived by transfection of the CV-1 cell line with a gene encoding Epstein-Barr virus nuclear antigen-I (EBNA-1) and constitutively expresses EBNA-1 driven from human CMV immediate-early enhancer/promoter.

10 Preferred vectors for expression in mammalian cultured cells include pFLAGCMV-1 (Kodak), pcDNA3.1/hyg (Invitrogen), pEE14-GS (CellTech), pBIB and VR1012 (Hartikka *et al.* (1996) *Hum. Gene Ther.* 7(10):1205-1217).

G. Host Cells

15 Transformed host cells are cells which have been transformed or transfected with expression vectors constructed using recombinant DNA techniques and which contain sequences encoding Her-2/neu fusion proteins of the present invention. Transformed host cells may express the desired Her-2/neu fusion proteins, but host cells transformed for purposes of cloning or amplifying Her-2/neu DNA do not need to express the Her-2/neu fusion proteins. Expressed Her-2/neu fusion proteins will preferably be
20 secreted into the culture medium or supernatant, depending on the DNA selected. One skilled in the art will appreciate that if Her-2/neu fusion proteins are secreted into the culture supernatant, then they are also soluble in the culture supernatant.

Any of the well known procedures for introducing foreign nucleotide sequences into host cells may be used to introduce the expression vector. These include
25 the use of reagents such as Superfect (Qiagen), liposomes, calcium phosphate transfection, polybrene, protoplast fusion, electroporation, microinjection, plasmid vectors, viral vectors, biolistic particle acceleration (the gene gun), or any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (*see, e.g., Sambrook et al., supra*).

30 Suitable host cells for expression of recombinant proteins include prokaryotes, yeast or higher eukaryotic cells under the control of appropriate promoters. Prokaryotes include gram negative or gram positive organisms, for example *E. coli* or *Bacilli*. Higher eukaryotic cells include established cell lines of insect or mammalian

origin as described below. Cell-free translation systems could also be used to produce Her-2/neu fusion proteins using RNAs derived from DNA constructs. Appropriate cloning and expression vectors for use with bacterial, fungal, yeast, and mammalian cellular hosts are described, for example, by Pouwels *et al.*, *Cloning Vectors: A*

5 *Laboratory Manual*, Elsevier, NY (1985).

Prokaryotic expression hosts may be used for expression of Her-2/neu fusion proteins that do not require extensive proteolytic and disulfide processing. Prokaryotic expression vectors generally comprise one or more phenotypic selectable markers, *e.g.*, a gene encoding a protein conferring antibiotic resistance or supplying an autotrophic requirement, and an origin of replication recognized by the host to ensure
10 amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli* (*e.g.*, BL21 (DE3) CodonPlus *E. coli*), *Bacillus subtilis*, *Salmonella typhimurium*, and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although other hosts may also be used.

15 Recombinant Her-2/neu fusion proteins may also be expressed in yeast hosts such as *P. pastoris*. Yeast of other genera, such as *Saccharomyces*, *Schizosaccharomyces* or *Kluyveromyces*, may also be used. Expression in *Pichia* is achieved by ligation of the gene to be expressed into a bacterial shuttle vector (*e.g.*, the pPICZ series from Invitrogen Co.), transformation of the yeast with this vector and
20 chromosomal integration into the alcohol oxidase (AOX) locus of the yeast genome. Selection for recombinant yeast is then performed using, *e.g.*, Zeocin (Invitrogen Co.) and protein expression is induced by the addition of methanol to the growth medium (Higgin *et al.*, "Pichia Protocols," *Methods in Molecular Biology*, Vol. 103, Humana Press (1998)). Suitable strains of *Pichia* for protein expression include, *e.g.*, the SMD1168
25 *Pichia* strain. Expression systems based on other methodologies, such as the ESP system (Stratagene) may also be used.

Suitable yeast transformation protocols are known to one of skill in the art. An exemplary technique described by Hind *et al.* (1978) *Proc. Natl. Acad. Sci. USA* 75:1929 involves selecting for Trp⁺ transformants in a selective medium consisting of
30 0.67% yeast nitrogen base, 0.5% casamino acids, 2% glucose, 10 mg/ml adenine and 20 mg/ml uracil. Host strains transformed by vectors comprising the ADH2 promoter may be grown for expression in a rich medium consisting of 1% yeast extract, 2% peptone, and 1% glucose supplemented with 80 mg/ml adenine and 80 mg/ml uracil. Derepression

of the ADH2 promoter occurs upon exhaustion of medium glucose. Crude yeast supernatants are harvested by filtration and held at 4°C prior to further purification.

Various mammalian or insect (*e.g.*, *Spodoptera* or *Trichoplusia*) cell culture systems can also be used to express recombinant polypeptides. Baculovirus systems for production of heterologous polypeptides in insect cells are reviewed, for example, by Luckow *et al.* (1988) *BioTechnology* 6:47. Examples of suitable mammalian host cell lines include the COS-7 lines of monkey kidney cells, described by Gluzman (1981) *Cell* 23:175, and other cell lines capable of expressing an appropriate vector including, *e.g.*, CV-1/EBNA (ATCC CRL 10478), L cells, C127, 3T3, Chinese hamster ovary (CHO), COS, NS-1, HeLa, Human embryonic Kidney Fibroblasts (HEK 293), BHK and HEK293 cell lines. Mammalian expression vectors may comprise nontranscribed elements (*e.g.*, an origin of replication, a suitable promoter and/or an enhancer linked to the gene to be expressed, and other 5' or 3' flanking nontranscribed sequences) and 5' or 3' nontranslated sequences (*e.g.*, necessary ribosome binding sites, a polyadenylation site, splice donor and acceptor sites, and transcriptional termination sequences). Preferred mammalian expression systems are the Chinese hamster ovary (CHO), the HEK293 and the BHK cell lines.

H. Purification of the Fusion Proteins of the Invention

Purified Her-2/neu fusion proteins may be prepared by culturing suitable host/vector systems to express the recombinant translation products of the DNAs of the present invention, which are then purified from culture media or cell extracts. For example, supernatants from systems which secrete recombinant polypeptides into culture media may be first concentrated using a commercially available protein concentration filter, such as, *e.g.*, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate may be applied to a suitable purification matrix. For example, a suitable affinity matrix may comprise a counter structure protein (*i.e.*, a protein to which a Her-2/neu fusion protein binds in a specific interaction based on structure) or lectin or antibody molecule bound to a suitable support.

Alternatively, an anion exchange resin can be used, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose, polystyrene, sepharose or other types commonly used in protein purification. Alternatively, a cation exchange step can be used. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or carboxy-

methyl groups, preferably sulfopropyl groups. Gel filtration chromatography also provides a means of purifying Her-2/neu fusion proteins. The fusion proteins of the invention are preferably purified by anion exchange chromatography using, *e.g.*, monoQ columns or Q sepharose High Performance chromatography.

5 Affinity chromatography is another preferred method of purifying Her-2/neu fusion proteins. For example, monoclonal antibodies against the Her-2/neu fusion proteins may be useful in affinity chromatography purification, by using methods that are well-known in the art.

Finally, one or more reverse-phase high performance liquid
10 chromatography (RP-HPLC) steps using hydrophobic RP-HPLC media (*e.g.*, silica gel having pendant methyl or other aliphatic groups) may be used to further purify Her-2/neu fusion protein compositions. Some or all of the foregoing purification steps, in various combinations, can also be used to provide a homogeneous recombinant protein or polypeptide.

15 Recombinant Her-2/neu fusion proteins produced in bacterial culture may be purified by initial extraction from cell pellets, followed by one or more concentration, salting-out, aqueous ion exchange or size exclusion chromatography steps. High performance liquid chromatography (HPLC) may be used for final purification steps. Microbial cells used in expression of recombinant Her-2/neu fusion proteins can be
20 disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Fermentation of yeast cells which express Her-2/neu fusion proteins as a secreted protein greatly simplifies purification. The secreted recombinant proteins resulting from a large-scale fermentation can be purified by methods analogous to those
25 disclosed by Urdal *et al.* (1984) *J. Chromatog.* 296:171. This reference describes two sequential, reverse-phase HPLC steps for purification of recombinant human GM-CSF on a preparative HPLC column.

Preparations of Her-2/neu fusion proteins synthesized in recombinant cultures may contain non- Her-2/neu cell components, including proteins, in amounts and
30 of a character which depend upon the purification steps taken to recover the Her-2/neu fusion proteins from the culture. These components are ordinarily of yeast, prokaryotic or non-human eukaryotic origin. Such preparations are typically free of other proteins which may be normally associated with the Her-2/neu protein as it is found in nature in its species of origin.

Automated synthesis provides an alternate method for preparing proteins and polypeptides of this invention. For example, any of the commercially available solid-phase techniques may be used, such as, *e.g.*, the Merrifield solid phase synthesis method, in which amino acids are sequentially added to a growing amino acid chain (*see*,
5 Merrifield (1963) *J. Am. Chem. Soc.* 85:2149-2146). Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Applied Biosystems, Inc. (Foster City, CA), and may generally be operated according to the manufacturer's instructions.

In general, polypeptides (including fusion proteins) and polynucleotides as
10 described herein are isolated. An "isolated" polypeptide or polynucleotide is one that is removed from its original environment. For example, a naturally-occurring protein is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. A polynucleotide is
15 considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

V. BINDING AGENTS

The present invention further provides agents, such as antibodies and antigen-binding fragments thereof, that specifically bind to a Her-2/neu fusion protein of
20 the invention. As used herein, an antibody, or antigen-binding fragment thereof, is said to "specifically bind" to a Her-2/neu fusion protein if it reacts at a detectable level, *i.e.*, at least two fold over background signal (within, for example, an ELISA) with a Her-2/neu fusion protein, and does not react detectably with unrelated proteins under similar conditions. As used herein, "binding" refers to a noncovalent association between two
25 separate molecules such that a complex is formed. The ability to bind may be evaluated by, for example, determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind," in the context of the present invention, when the binding constant for complex
30 formation exceeds about 10^3 l/mol. The binding constant may be determined using methods well known in the art.

Binding agents may be further capable of differentiating between patients with and without a cancer, such as breast, ovarian, colon, lung or prostate cancer, using

the representative assays provided herein. In other words, antibodies or other binding agents that bind to a Her-2/neu fusion protein will generate a signal indicating the presence of a cancer in at least about 20% of patients with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without the cancer. To determine whether a binding agent satisfies this requirement, biological samples (*e.g.*, blood, sera, plasma, urine and/or tumor biopsies) from patients with and without a cancer (as determined using standard clinical tests) may be assayed as described herein for the presence of polypeptides that bind to the binding agent. It will be apparent that a statistically significant number of samples with and without the disease should be assayed. Each binding agent should satisfy the above criteria; however, those of ordinary skill in the art will recognize that binding agents may be used in combination to improve sensitivity.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome, with or without a peptide component, an RNA molecule or a polypeptide. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art (*see, e.g.*, Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory (1988)). In general, antibodies can be produced by cell culture techniques, including the generation of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising, *e.g.*, a fusion polypeptide or the sequence corresponding to the junction between the individual polypeptides of a fusion protein of interest (referred to as "junction region"), is initially injected into any of a wide variety of mammals (*e.g.*, mice, rats, rabbits, sheep or goats). In this step, the fusion protein of interest or the junction region of the fusion protein of the invention may serve as the immunogen without modification. Alternatively, particularly for relatively short sequences, a superior immune response may be elicited if the sequence is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the fusion polypeptide may then be purified from such antisera by, for example, affinity chromatography using the fusion polypeptide coupled to a suitable solid support.

Polyclonal antibodies raised to a fusion protein of the invention can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with the fusion protein of interest and not with the individual polypeptide components of the fusion proteins. This selection may be achieved by subtracting out antibodies that
5 cross-react with the individual polypeptide components of the fusion protein of interest.

Alternatively, antibodies that recognize each or all of the individual polypeptide components of a fusion protein may be useful in the context of the present invention.

Monoclonal antibodies specific for an immunogenic fusion polypeptide of
10 interest may be prepared, for example, using the technique of Kohler and Milstein (1976) *Eur. J. Immunol.* 6:511-519, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.*, reactivity with the fusion polypeptide of interest). Such cell lines may be produced, *e.g.*, from spleen cells obtained from an animal immunized as described above.
15 The spleen cells are then immortalized by, *e.g.*, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A
20 preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and their culture supernatants tested for binding activity against the fusion polypeptide. Hybridomas having high reactivity and specificity are preferred.

25 Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the ascites fluid or the blood. Contaminants may be removed from the antibodies by
30 conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The fusion polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Within certain embodiments, the use of antigen-binding fragments of antibodies may be preferred. Such fragments include Fab fragments, which may be

prepared using standard techniques. Briefly, immunoglobulins may be purified from rabbit serum by affinity chromatography on Protein A bead columns (Harlow and Lane, *supra*) and digested by papain to yield Fab and Fc fragments. The Fab and Fc fragments may be separated by affinity chromatography on protein A bead columns.

5 Monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers
10 include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, *Diphtheria* toxin, cholera toxin, gelonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct
15 reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a halide) on the other.

20 Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity
25 may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group.
30 Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, including, *e.g.*, U.S. Patent No. 4,671,958.

Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a

linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a disulfide bond (*e.g.*, U.S. Patent No. 4,489,710), by irradiation of a photolabile bond
5 (*e.g.*, U.S. Patent No. 4,625,014), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958), and by acid-catalyzed hydrolysis (*e.g.*, U.S. Patent No. 4,569,789).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In
10 another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers that provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

15 A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as, *e.g.*, albumins (*e.g.*, U.S. Patent No. 4,507,234), peptides and polysaccharides such as, *e.g.*, aminodextran (*e.g.*, U.S. Patent No. 4,699,784). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S.
20 Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example, U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or
25 metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562 discloses representative chelating compounds and their synthesis.

A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the
30 precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

Examples of available suitable antibodies to the fusion proteins of the invention include, but are not limited to, the 8029K rabbit polyclonal antibody, the mouse monoclonal c-neu-3 antibody (Calbiochem), and the mouse monoclonal Herceptin

antibody (U.S. Patent 5,677,171). The monoclonal c-neu-3 antibody recognizes a sequential epitope in the PD domain which is deleted (1242-1255 aa) in the ECD-ΔPD construct. The Herceptin antibody binds to a conformational epitope in the ECD domain.

VI. T CELLS

5 Immunotherapeutic compositions may also, or alternatively, comprise T cells specific for a fusion protein of the present invention. Such cells may generally be prepared *in vitro* or *ex vivo*, using standard procedures. For example, T cells may be isolated from bone marrow, peripheral blood or a fraction of bone marrow or peripheral blood of a patient, using a commercially available cell separation system (*see also*, U.S. 10 Patent Nos. 5,240,856 and 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from related or unrelated humans, non-human mammals, cell lines or cultures.

T cells may be stimulated with a Her-2/neu fusion polypeptide, a polynucleotide encoding a Her-2/neu fusion protein and/or an antigen presenting cell 15 (APC) that expresses such a fusion polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T cells that are specific for the fusion polypeptide. Preferably, a Her-2/neu fusion polypeptide or polynucleotide is present within a delivery vehicle, such as a microsphere, to facilitate the generation of specific T cells.

20 T cells are considered to be specific for a Her-2/neu fusion polypeptide if the T cells kill target cells coated with the fusion polypeptide or expressing a polynucleotide encoding the fusion polypeptide. T cell specificity may be evaluated using any of a variety of standard techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis 25 and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as described in Chen *et al.* (1994) *Cancer Res.* 54:1065-1070. Alternatively, detection of the proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (*e.g.*, by pulse-labeling 30 cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Contact with a Her-2/neu fusion polypeptide (100 ng/ml-100 μg/ml, preferably 200 ng/ml-25 μg/ml) for 3-7 days should result in at least a two fold increase in proliferation of the T cells. Contact as described above for 2-3 hours

should result in the activation of the T cells, as measured using standard cytokine assays in which a two fold increase in the level of cytokine release (*e.g.*, TNF or IFN- γ) is indicative of T cell activation (*see*, Coligan *et al.*, *Current Protocols in Immunology*, vol. 1, Wiley Interscience, Greene (1998)). T cells that have been activated in response to a Her-2/neu fusion polypeptide, polynucleotide or fusion polypeptide-expressing APC may be CD4⁺ and/or CD8⁺. Her-2/neu fusion protein-specific T cells may be expanded using standard techniques. Within preferred embodiments, the T cells are derived from a patient, or from a related or unrelated donor, and are administered to the patient following stimulation and expansion.

For therapeutic purposes, CD4⁺ or CD8⁺ T cells that proliferate in response to a Her-2/neu fusion polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to a Her-2/neu fusion polypeptide with or without the addition of T cell growth factors, such as interleukin-2, and/or stimulator cells that synthesize a Her-2/neu fusion polypeptide. Alternatively, one or more T cells that proliferate in the presence of a Her-2/neu fusion protein can be expanded in number by cloning. Methods for cloning cells are well known in the art, and include limiting dilution. Following expansion, the cells may be administered back to the patient as described, for example, by Chang *et al.* (1996) *Crit. Rev. Oncol. Hematol.* 22:213.

VII. PHARMACEUTICAL COMPOSITIONS AND VACCINES COMPRISING FUSION PROTEINS OF THE INVENTION

In another preferred embodiment, the present invention is directed to compositions comprising Her-2/neu fusion proteins, or variants thereof, and Her-2/neu ICD proteins, or variants thereof. The fusion proteins are preferably ECD-ICD fusion proteins, ECD-PD fusion proteins, or variants thereof, of the present invention, as described in detail herein. The ICD protein is preferably the human ICD protein, which spans the region from Lys 676 to Val 1255, inclusively, as shown in Fig. 7 (SEQ ID NO:1), or the rat ICD protein, which is the amino acid sequence inclusively spanning Lys 677 to Val 1256 as shown in Fig. 8 (SEQ ID NO:2). Alternatively, the Her-2/neu ICD protein can be any variant or portion of the ICD protein that is immunogenic or that provides enhanced immunogenicity to the composition. For example, the portion of the ICD protein may be the Her-2/neu PD protein, as described herein, the Her-2/neu Δ PD

protein, as described herein, the Her-2/neu KD protein, as described herein, or a Her-2/neu ICD protein where anywhere from 1 to 100 amino acids are sequentially removed from the N-terminus or C-terminus of the ICD protein. In addition, amino acid substitutions can generally be made in a variety of ways to provide other embodiments of variants within the present invention. In one embodiment, conservative amino acid substitutions are made, as described *supra*.

Within certain aspects, polypeptides, polynucleotides, T cells and/or binding agents described herein may be incorporated into pharmaceutical compositions or immunogenic compositions (*i.e.*, vaccines). Pharmaceutical compositions comprise one or more such compounds and a physiologically acceptable carrier. Vaccines may comprise one or more such compounds and a non-specific immune response enhancer. A non-specific immune response enhancer may be any substance that enhances an immune response to an exogenous antigen. Examples of non-specific immune response enhancers include adjuvants, biodegradable microspheres (*e.g.*, polylactic galactide) and liposomes (into which the compound is incorporated; *see, e.g.*, U.S. Patent No. 4,235,877). Vaccine preparation is generally described in, for example, Powell and Newman, eds., *Vaccine Design* (the subunit and adjuvant approach), Plenum Press, NY(1995). Vaccines may be designed to generate antibody immunity and/or cellular immunity such as that arising from CTL or CD4⁺ T cells.

Pharmaceutical compositions and vaccines within the scope of the present invention may also contain other compounds, which may be biologically active or inactive. For example, one or more immunogenic portions of other tumor antigens may be present, either incorporated into a fusion polypeptide or as a separate compound, within the composition or vaccine. Polypeptides may, but need not, be conjugated to other macromolecules as described, for example, within U.S. Patent Nos. 4,372,945 and 4,474,757. Pharmaceutical compositions and vaccines may generally be used for prophylactic and therapeutic purposes.

A pharmaceutical composition or vaccine may contain a polynucleotide encoding one or more of the Her-2/neu fusion proteins, *e.g.*, Her-2/neu ECD-ICD and/or Her-2/neu ECD-PD, as described above, such that the fusion protein is generated *in situ*. Such a polynucleotide may comprise DNA, RNA, a modified nucleic acid or a DNA/RNA hybrid. As noted above, a polynucleotide may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Numerous gene delivery

techniques are well known in the art, such as those described by Rolland (1998) *Crit. Rev. Therap. Drug Carrier Systems* 15:143-198, and references cited therein. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus Calmette-Guerin*) that expresses an immunogenic portion of the fusion protein on its cell surface or secretes such an epitope.

In a preferred embodiment, the DNA may be introduced using a viral expression system (e.g., vaccinia, pox virus, retrovirus, adenovirus or alphavirus such as SFV), which may involve the use of a part of a virus, of a non-pathogenic (defective), replication competent virus or of a whole virus.

In one illustrative embodiment, retroviruses provide a convenient and effective platform for gene delivery systems. A selected nucleotide sequence encoding a fusion protein of the present invention can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to a subject. A number of illustrative retroviral systems have been described (e.g., U.S. Patent No. 5,219,740; Miller and Rosman (1989) *BioTechniques* 7:980-990; Miller (1990) *Human Gene Therapy* 1:5-14; Scarpa *et al.* (1991) *Virology* 180:849-852; Burns *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:8033-8037; and Boris-Lawric and Temin (1993) *Curr. Opin. Genet. Develop.* 3:102-109.

In addition, a number of illustrative adenovirus-based systems have also been described. Unlike retroviruses which integrate into the host genome, adenoviruses persist extrachromosomally thus minimizing the risks associated with insertional mutagenesis (Haj-Ahmad and Graham (1986) *J. Virol.* 57:267-274; Bett *et al.* (1993) *J. Virol.* 67:5911-5921; Mittereder *et al.* (1994) *Human Gene Therapy* 5:717-729; Seth *et al.* (1994) *J. Virol.* 68:933-940; Barr *et al.* (1994) *Gene Therapy* 1:51-58; Berkner (1988) *BioTechniques* 6:616-629; and Rich *et al.* (1993) *Human Gene Therapy* 4:461-476).

Various adeno-associated virus (AAV) vector systems have also been developed for polynucleotide delivery. AAV vectors can be readily constructed using techniques well known in the art (see, e.g., U.S. Patent Nos. 5,173,414 and 5,139,941; PCT Publication Nos. WO 92/01070 and WO 93/03769; Lebkowski *et al.* (1988) *Mol. Cell. Biol.* 8:3988-3996; Vincent *et al.* (1990) *Vaccines* 90 (Cold Spring Harbor Laboratory Press); Carter (1992) *Current Opinion in Biotechnology* 3:533-539; Muzyczka (1992) *Current Topics in Microbiol. and Immunol.* 158:97-129; Kotin (1994)

Human Gene Therapy 5:793-801; Shelling and Smith (1994) *Gene Therapy* 1:165-169; and Zhou *et al.* (1994) *J. Exp. Med.* 179:1867-1875).

Additional viral vectors useful for delivering the nucleic acid molecules encoding fusion proteins of the present invention by gene transfer include those derived from the pox family of viruses, such as vaccinia virus and avian poxvirus. By way of example, vaccinia virus recombinants expressing the novel molecules can be constructed as follows. The DNA encoding a fusion protein is first inserted into an appropriate vector so that it is adjacent to a vaccinia promoter and flanking vaccinia DNA sequences, such as the sequence encoding thymidine kinase (TK). This vector is then used to transfect cells which are simultaneously infected with vaccinia. Homologous recombination serves to insert the vaccinia promoter plus the gene encoding the fusion protein of interest into the viral genome. The resulting TK^{sup}(-) recombinant can be selected by culturing the cells in the presence of 5-bromodeoxyuridine and picking viral plaques resistant thereto.

A vaccinia-based infection/transfection system can be conveniently used to provide for inducible, transient expression or coexpression of one or more fusion proteins described herein in host cells of an organism. In this particular system, cells are first infected *in vitro* with a vaccinia virus recombinant that encodes the bacteriophage T7 RNA polymerase. This polymerase displays exquisite specificity in that it only transcribes templates bearing T7 promoters. Following infection, cells are transfected with the polynucleotide or polynucleotides of interest, driven by a T7 promoter. The polymerase expressed in the cytoplasm from the vaccinia virus recombinant transcribes the transfected DNA into RNA which is then translated into polypeptide by the host translational machinery. The method provides for high level, transient, cytoplasmic production of large quantities of RNA and its translation products (*see, e.g.,* Elroy-Stein and Moss (1990) *Proc. Natl. Acad. Sci. USA* 87:6743-6747; Fuerst *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:8122-8126).

Alternatively, avipoxviruses, such as the fowlpox and canarypox viruses, can also be used to deliver the coding sequences of interest. Recombinant avipox viruses, expressing immunogens from mammalian pathogens, are known to confer protective immunity when administered to non-avian species. The use of an Avipox vector is particularly desirable in human and other mammalian species since members of the Avipox genus can only productively replicate in susceptible avian species and therefore are not infective in mammalian cells. Methods for producing recombinant Avipoxviruses are known in the art and employ genetic recombination, as described above with respect

to the production of vaccinia viruses (*see, e.g.*, WO 91/12882; WO 89/03429; and WO 92/03545).

Any of a number of alphavirus vectors (*e.g.*, Semliki Forest Virus) can also be used for delivery of polynucleotide compositions of the present invention, such as those vectors described in U.S. Patent Nos. 5,843,723; 6,015,686; 6,008,035 and 6,015,694. Certain vectors based on Venezuelan Equine Encephalitis (VEE) can also be used, illustrative examples of which can be found in U.S. Patent Nos. 5,505,947 and 5,643,576.

Moreover, molecular conjugate vectors, such as the adenovirus chimeric vectors described in Michael *et al.* (1993) *J. Biol. Chem.* 268:6866-6869 and Wagner *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:6099-6103, can also be used for gene delivery under the invention.

Additional illustrative information on these and other known viral-based delivery systems can be found, for example, in Fisher-Hoch *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:317-321; Flexner *et al.* (1989) *Ann. N.Y. Acad. Sci.* 569:86-103; Flexner *et al.* (1990) *Vaccine* 8:17-21; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner (1988) *Biotechniques* 6:616-627; Rosenfeld *et al.* (1991) *Science* 252:431-434; Kolls *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:215-219; Kass-Eisler *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:11498-11502; Guzman *et al.* (1993) *Circulation* 88:2838-2848; and Guzman *et al.* (1993) *Cir. Res.* 73:1202-1207.

Techniques for incorporating DNA into such expression systems are well known to those of ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer *et al.* (1993) *Science* 259:1745-1749 and reviewed by Cohen (1993) *Science* 259:1691-1692. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells. It will be apparent that a vaccine may comprise both a polynucleotide and a polypeptide component. Such vaccines may provide for an enhanced immune response.

It will be apparent that a vaccine may contain pharmaceutically acceptable salts of the polynucleotides and fusion polypeptides provided herein. Such salts may be prepared from pharmaceutically acceptable non-toxic bases, including organic bases (*e.g.*, salts of primary, secondary and tertiary amines and basic amino acids) and inorganic bases (*e.g.*, sodium, potassium, lithium, ammonium, calcium and magnesium salts).

While any suitable carrier known to those of ordinary skill in the art may be employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration, including for example, 5 topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous or intramuscular administration. For parenteral administration, such as subcutaneous injection, the carrier preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, 10 sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactate polyglycolate) may also be employed as carriers for the pharmaceutical compositions of this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268; 5,075,109; 5,928,647; 5,811,128; 5,820,883. The Her-2/neu fusion protein may be encapsulated within the biodegradable microsphere 15 or associated with the surface of the microsphere. In one embodiment, an ECD-ICD fusion protein described herein is encapsulated within a biodegradable microsphere. Alternatively or in addition, an ECD-PD fusion protein described herein is encapsulated within a biodegradable microsphere. The microsphere can comprise, *e.g.*, both an ECD-ICD fusion protein and an ECD-PD fusion protein. Preferably the microsphere be less 20 than about 25 μm , preferably about 1 μm to about 10 μm . Encapsulation in liposomes is described, *e.g.*, in U.S. Patent No. 4,235,877.

Such compositions may also comprise buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, 25 bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide), solutes that render the formulation isotonic, hypotonic or weakly hypertonic with the blood of a recipient, suspending agents, thickening agents and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate. Compounds may also be encapsulated within liposomes using well known technology.

30 Any of a variety of immune response enhancers or immunostimulatory substances may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of

immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Suitable adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI); Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ); AS-2 and variants thereof
5 (SmithKline Beecham, Philadelphia, PA); aluminum salts such as aluminum hydroxide gel (alum) or aluminum phosphate; salts of calcium, iron or zinc; an insoluble suspension of acylated tyrosine; acylated sugars; cationically or anionically derivatized polysaccharides; polyphosphazenes; biodegradable microspheres; monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF or interleukin-2, -7, or -12, may also be used
10 as adjuvants.

Within the vaccines provided herein, the adjuvant composition can be designed to induce, *e.g.*, an immune response predominantly of the Th1 type. High levels of Th1-type cytokines (*e.g.*, IFN- γ , TNF α , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of
15 Th2-type cytokines (*e.g.*, IL-4, IL-5, IL-6 and IL-10) tend to favor the induction of humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type
20 cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review of the families of cytokines, *see*, Mosmann and Coffman (1989) *Ann. Rev. Immunol.* 7:145-173.

Preferred adjuvants for use in eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL), together with an aluminum salt. MPL
25 adjuvants are available from Corixa Corporation (Seattle, WA; *see*, U.S. Patent Nos. 4,436,727; 4,877,611; 4,866,034; and 4,912,094). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555,
30 WO 99/33488 and U.S. Patent Nos. 6,008,200 and 5,856,462. Immunostimulatory DNA sequences are also described, for example, by Sato *et al.* (1996) *Science* 273:352. Another preferred adjuvant is a saponin, preferably QS21 (Aquila Biopharmaceuticals Inc., Framingham, MA), which may be used alone or in combination with other

adjuvants. For example, an enhanced system involves the combination of a monophosphoryl lipid A and saponin derivative, such as the combination of QS21 and 3D-MPL as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred
5 formulations comprise an oil-in-water emulsion and tocopherol. A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil-in-water emulsion is described in WO 95/17210.

Other preferred adjuvants include Montanide ISA 720 (Seppic, France), SAF (Chiron, California, United States), ISCOMS (CSL), MF-59 (Chiron), the SBAS
10 series of adjuvants (*e.g.*, SBAS-2, SBAS-4 or SBAS-6 or variants thereof, available from SmithKline Beecham, Rixensart, Belgium), Detox (Corixa, Hamilton, MT), RC-529 (Corixa, Hamilton, MT) and other aminoalkyl glucosaminide 4-phosphates (AGPs), such as those described in pending U.S. Patent Application Serial Nos. 08/853,826 and 09/074,720.

15 In a preferred embodiment, the adjuvant is SBAS-2 or SBAS-4 (*see, e.g.*, EP 735898B1).

Any vaccine provided herein may be prepared using well known methods that result in a combination of antigen, immune response enhancer and a suitable carrier or excipient. The compositions described herein may be administered as part of a
20 sustained release formulation (*i.e.*, a formulation such as a capsule or sponge that effects a slow release of compound following administration). Such formulations may generally be prepared using well known technology (*see, e.g.*, Coombes *et al.* (1996) *Vaccine* 14:1429-1438) and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations
25 may contain a polypeptide, polynucleotide or antibody dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane.

Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. Such carriers include microparticles of poly(lactide-co-glycolide), as
30 well as polyacrylate, latex, starch, cellulose and dextran. Other delayed-release carriers include supramolecular biovectors, which comprise a non-liquid hydrophilic core (*e.g.*, a cross-linked polysaccharide or oligosaccharide) and, optionally, an external layer comprising an amphiphilic compound, such as a phospholipid (*see, e.g.*, U.S. Patent No. 5,151,254; and PCT applications WO 94/20078; WO/94/23701; and WO 96/06638). The

amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

Any of a variety of delivery vehicles may be employed within
5 pharmaceutical compositions and vaccines to facilitate production of an antigen-specific immune response that targets tumor cells. Delivery vehicles include antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or
10 maintenance of the T cell response, to have anti-tumor effects *per se* and/or to be immunologically compatible with the receiver (*i.e.*, matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, including tumor and peritumoral tissues, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

15 Certain preferred embodiments of the present invention use dendritic cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau *et al.* (1998) *Nature* 392:245-251) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic antitumor immunity (*see*, Timmerman *et al.* (1999) *Ann. Rev. Med.* 50:507-529). In general, dendritic cells may be
20 identified based on their typical shape (stellate *in situ*, with marked cytoplasmic processes (dendrites) visible *in vitro*), their ability to take up, process and present antigens with high efficiency, and their ability to activate naïve T cell responses. Dendritic cells may, of course, be engineered to express specific cell-surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells
25 are contemplated by the present invention. As an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within a vaccine (*see*, Zitvogel *et al.* (1998) *Nature Med.* 4:594-600).

Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, tumor-infiltrating cells, peritumoral tissues-infiltrating cells, lymph nodes,
30 spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF α to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells harvested from peripheral blood, umbilical cord

blood or bone marrow may be differentiated into dendritic cells by adding to the culture medium combinations of GM-CSF, IL-3, TNF α , CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce maturation and proliferation of dendritic cells.

Dendritic cells are conveniently categorized as "immature" and "mature" cells, which allows a simple way to discriminate between two well characterized phenotypes. However, this nomenclature should not be construed to exclude all possible intermediate stages of differentiation. Immature dendritic cells are characterized as APCs with a high capacity for antigen uptake and processing, which correlates with the high expression of Fc γ receptor and mannose receptor. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (*e.g.*, CD54 and CD11) and costimulatory molecules (*e.g.*, CD40, CD80, CD86 and 4-1BB).

APCs may generally be transfected with a polynucleotide encoding a fusion protein of the invention (or variant thereof) such that the fusion protein, or a variant thereof, is expressed on the cell surface. Such transfection may take place *ex vivo*, and a composition or vaccine comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi *et al.* (1997) *Immunology and cell Biology* 75:456-460. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the fusion protein of interest, DNA (naked or within a plasmid vector) or RNA; or with fusion protein-expressing recombinant bacteria or viruses (*e.g.*, vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the fusion protein of interest may be covalently conjugated to an immunological partner that provides T cell help (*e.g.*, a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the fusion protein.

Vaccines and pharmaceutical compositions may be presented in unit-dose or multi-dose containers, such as sealed ampoules or vials. Such containers are preferably hermetically sealed to preserve sterility of the formulation until use. In

general, formulations may be stored as suspensions, solutions or emulsions in oily or aqueous vehicles. Alternatively, a vaccine or pharmaceutical composition may be stored in a freeze-dried condition requiring only the addition of a sterile liquid carrier immediately prior to use.

- 5 It will be evident to one skilled in the art that a Her-2/neu fusion protein or nucleic acid for a vaccine may be prepared synthetically or may be naturally derived.

VIII. IMMUNE RESPONSE TO FUSION PROTEINS OF THE INVENTION

A. Detection of an Immune Response to Fusion Proteins of the Invention

- In one aspect of the invention, Her-2/neu fusion proteins (or
10 polynucleotides that encode Her-2/neu fusion proteins) are used to generate an immune response to the Her-2/neu protein, including that expressed in a malignancy in which a Her-2/neu oncogene is associated. Representative examples of such malignancies include breast, ovarian, colon, lung and prostate cancers. An immune response to the Her-2/neu protein, once generated by Her-2/neu fusion proteins, can be long-lived and can be
15 detected long after immunization, regardless of whether the protein is present or absent in the body at the time of testing. An immune response to the Her-2/neu protein generated by reaction to a Her-2/neu fusion protein can be detected by examining for the presence or absence, or enhancement, of specific activation of CD4⁺ or CD8⁺ T cells or by antibodies. For instance, T cells isolated from an immunized individual by routine
20 techniques (*e.g.*, by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes) are incubated with a Her-2/neu fusion protein. For example, T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with a Her-2/neu fusion protein (typically, 5 µg/ml of whole protein or graded numbers of cells synthesizing Her-2/neu protein). It may be desirable to incubate another aliquot of a T cell sample in the absence
25 of Her-2/neu fusion protein to serve as a control.

- Specific activation of CD4⁺ or CD8⁺ T cells may be detected in a variety of ways. Methods for detecting specific T cell activation include, but are not limited to, detecting the proliferation of T cells, the production of cytokines (*e.g.*, lymphokines), or the generation of cytolytic activity (*i.e.*, generation of cytotoxic T cells specific for a Her-
30 2/neu fusion protein). For CD4⁺ T cells, a preferred method for detecting specific T cell activation is the detection of the proliferation of T cells. For CD8⁺ T cells, a preferred method for detecting specific T cell activation is the detection of the generation of cytolytic activity.

Detection of the proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring the rate of DNA synthesis. T cells which have been stimulated to proliferate exhibit an increased rate of DNA synthesis. A typical way to measure the rate of DNA synthesis is, for example, by pulse-labeling cultures of T cells with tritiated thymidine, a nucleoside precursor which is incorporated into newly synthesized DNA. The amount of tritiated thymidine incorporated can be determined using a liquid scintillation spectrophotometer. Other ways to detect T cell proliferation include measuring increases in interleukin-2 (IL-2) production, Ca^{2+} flux, or dye uptake, such as 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium. Alternatively, synthesis of lymphokines (*e.g.*, interferon-gamma) can be measured or the relative number of T cells that can respond to intact p185^{Her-2/neu} protein may be quantified.

B. Detection of Antibody Production in Response to Fusion Proteins of the Invention

The present invention is also directed to Her-2/neu fusion proteins that, in addition to being immunogenic to T cells, appear to stimulate B-cells to produce antibodies capable of recognizing Her-2/neu fusion proteins. Detection of such antibodies provides another way to diagnose a malignancy in which a Her-2/neu oncogene is associated with the malignancy. Antibodies specific (*i.e.*, which exhibit a binding affinity of about 10^7 l/mole or better) for Her-2/neu fusion proteins may be found in a variety of body fluids including sera and ascites. Briefly, a body fluid sample is isolated from a warm-blooded animal, such as a human, for whom it is desired to determine whether antibodies specific for the fusion proteins are present. The body fluid is incubated with Her-2/neu fusion proteins under conditions and for a time sufficient to permit immunocomplexes to form between the Her-2/neu fusion proteins and antibodies specific for the fusion proteins. For example, a body fluid and Her-2/neu fusion proteins may be incubated at 46°C for 24-48 hours. Following the incubation, the reaction mixture is tested for the presence of immunocomplexes. Detection of one or more immunocomplexes formed between Her-2/neu fusion protein and antibodies specific for Her-2/neu fusion protein may be accomplished by a variety of known techniques, such as radioimmunoassays (RIA) and enzyme linked immunosorbent assays (ELISA).

Suitable immunoassays include the double monoclonal antibody sandwich immunoassay technique of David *et al.* (U.S. Patent No. 4,376,110); monoclonal-polyclonal antibody sandwich assays (Wide *et al.*, in Kirkham and Hunter, eds.,

Radioimmunoassay Methods, E. and S. Livingstone, Edinburgh (1970)); the "western blot" method of Gordon *et al.* (U.S. Patent No. 4,452,901); immunoprecipitation of labeled ligand (Brown *et al.* (1980) *J. Biol. Chem.* 255:4980-4983); enzyme-linked immunosorbent assays (ELISA) as described, for example, by Raines *et al.* (1982) *J. Biol.*
5 *Chem.* 257:5154-5160; immunocytochemical techniques, including the use of fluorochromes (Brooks *et al.* (1980) *Clin. Exp. Immunol.* 39:477); and neutralization of activity (Bowen-Pope *et al.* (1984) *Proc. Natl. Acad. Sci. USA* 81:2396-2400). In addition to the immunoassays described above, a number of other immunoassays are available, including those described in U.S. Patent Nos. 3,817,827; 3,850,752; 3,901,654;
10 3,935,074; 3,984,533; 3,996,345; 4,034,074; and 4,098,876.

For detection purposes, Her-2/neu fusion proteins (*i.e.*, antigens) may either be labeled or unlabeled. When unlabeled, fusion proteins find use in agglutination assays. In addition, unlabeled fusion proteins can be used in combination with labeled molecules that are reactive with immunocomplexes, or in combination with labeled
15 antibodies (second antibodies) that are reactive with the antibody directed against Her-2/neu fusion protein, such as antibodies specific for immunoglobulin. Alternatively, the fusion protein can be directly labeled. Where it is labeled, the reporter group can include, *e.g.*, radioisotopes, fluorophores, enzymes, luminescers, dye particles and the like. These and other labels are well known in the art and are described, for example, in U.S. Patent
20 Nos. 3,766,162; 3,791,932; 3,817,837; 3,996,345; and 4,233,402.

Typically in an ELISA, the fusion protein of interest is adsorbed to the surface of a microtiter well. Residual protein-binding sites on the surface are then blocked with an appropriate agent, such as bovine serum albumin (BSA), heat-inactivated normal goat serum (NGS), or BLOTTO (buffered solution of nonfat dry milk which also
25 contains a preservative, salts, and an antifoaming agent). The well is then incubated with a sample suspected of containing specific antibody. The sample can be applied neat, or, more often, it can be diluted; usually in a buffered solution which contains a small amount (0.1%-5.0% by weight) of protein, such as BSA, NGS, or BLOTTO. After incubating for a sufficient length of time to allow specific binding to occur, the well is
30 washed to remove unbound protein and then incubated with an anti-species specific immunoglobulin antibody labeled with a reporter group. The reporter group can be chosen from a variety of enzymes, including, *e.g.*, horseradish peroxidase, beta-galactosidase, alkaline phosphatase, and glucose oxidase. Sufficient time is allowed for specific binding to occur, then the well is again washed to remove unbound conjugate,

and the substrate for the enzyme is added. Color is allowed to develop and the optical density of the contents of the well is determined visually or instrumentally.

In one preferred embodiment of this aspect of the present invention, a reporter group is bound to the Her-2/neu fusion protein of interest. The step of detecting immunocomplexes involves removing substantially any unbound Her-2/neu fusion protein and then detecting the presence or absence of the reporter group.

In another preferred embodiment, a reporter group is bound to a second antibody capable of binding to the antibodies specific for Her-2/neu fusion proteins. The step of detecting immunocomplexes involves (a) removing substantially any unbound antibody, (b) adding the second antibody, (c) removing substantially any unbound second antibody and then (d) detecting the presence or absence of the reporter group. Where the antibody specific for the Her-2/neu fusion protein of interest is derived from a human, the second antibody is an anti-human antibody.

In a third preferred embodiment for detecting immunocomplexes, a reporter group is bound to a molecule capable of binding to the immunocomplexes. The step of detecting involves (a) adding the molecule, (b) removing substantially any unbound molecule, and then (c) detecting the presence or absence of the reporter group. An example of a molecule capable of binding to the immunocomplexes is protein A.

It will be evident to one skilled in the art that a variety of methods for detecting the immunocomplexes may be used within the present invention. Reporter groups suitable for use in any of the methods include, *e.g.*, radioisotopes, fluorophores, enzymes, luminescers, and dye particles.

In a related aspect of the present invention, detection of immunocomplexes formed between Her-2/neu fusion proteins and antibodies in body fluid which are specific for Her-2/neu fusion proteins may be used to monitor the effectiveness of cancer therapy, which involves a Her-2/neu fusion protein, for a malignancy in which the Her-2/neu oncogene is associated. Samples of body fluid taken from an individual prior to and subsequent to initiation of therapy may be analyzed for the immunocomplexes by the methodologies described above. Briefly, the number of immunocomplexes detected in both samples are compared. A substantial change in the number of immunocomplexes in the second sample (post-therapy initiation) relative to the first sample (pre-therapy) reflects successful therapy.

IX. CANCER THERAPY

In further aspects of the present invention, the compositions described herein may be used for immunotherapy of cancer, such as breast, ovarian, colon, lung and prostate cancer. Within such methods, pharmaceutical compositions and vaccines are typically administered to a patient. As used herein, a "patient" refers to any warm-blooded animal, preferably a human. A patient may or may not be afflicted with cancer. Accordingly, the above pharmaceutical compositions and vaccines may be used to prevent the development of a cancer or to treat a patient afflicted with a cancer. A cancer may be diagnosed using criteria generally accepted in the art, including the presence of a malignant tumor. Pharmaceutical compositions and vaccines may be administered either prior to or following surgical removal of primary tumors and/or treatment such as administration of radiotherapy or conventional chemotherapeutic drugs. Administration may be by any suitable method, including administration by intravenous, intraperitoneal, intramuscular, subcutaneous, intranasal, intradermal, anal, vaginal, topical, sublingual and oral routes.

Within certain embodiments, immunotherapy may be active immunotherapy, in which treatment relies on the *in vivo* stimulation of the endogenous host immune system to react against tumors with the administration of immune response-modifying agents (such as fusion polypeptides and polynucleotides as provided herein).

Within other embodiments, immunotherapy may be passive immunotherapy, in which treatment involves the delivery of agents with established tumor-immune reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T cells as discussed *supra*, T lymphocytes (such as CD8⁺ cytotoxic T lymphocytes and CD4⁺ T-helper tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells and lymphokine-activated killer cells), B cells and antigen-presenting cells (such as dendritic cells and macrophages) expressing a fusion protein provided herein. T cell receptors and antibody receptors specific for the fusion polypeptides recited herein may be cloned, expressed and transferred into other vectors or effector cells for adoptive immunotherapy. The fusion polypeptides provided herein may also be used to generate antibodies or anti-idiotypic antibodies (as described above and in U.S. Patent No. 4,918,164) for passive immunotherapy.

Effector cells may generally be obtained in sufficient quantities for adoptive immunotherapy by growth *in vitro*, as described herein. Culture conditions for expanding single antigen-specific effector cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. Such *in vitro* culture conditions typically use intermittent stimulation with antigen, often in the presence of cytokines (such as IL-2) and non-dividing feeder cells. As noted above, immunoreactive fusion polypeptides as provided herein may be used to rapidly expand antigen-specific T cell cultures in order to generate a sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage or B cells, may be pulsed with immunoreactive fusion polypeptides or transfected with one or more polynucleotides using standard techniques well known in the art. For example, antigen-presenting cells can be transfected with a polynucleotide having a promoter appropriate for increasing expression in a recombinant virus or other expression system. Cultured effector cells for use in therapy must be able to grow and distribute widely, and to survive long term *in vivo*. Studies have shown that cultured effector cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (*see, for example, Cheever et al. (1997) Immunological Reviews 157:177*).

Alternatively, a vector expressing a fusion polypeptide recited herein may be introduced into antigen presenting cells taken from a patient and clonally propagated *ex vivo* for transplant back into the same patient. Transfected cells may be reintroduced into the patient using any means known in the art, preferably in sterile form by intravenous, intracavitary, intraperitoneal or intratumor administration.

Routes and frequency of administration of the therapeutic compositions described herein, as well as dosage, will vary from individual to individual, and may be readily established using standard techniques. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g.*, intracutaneous, intramuscular, intravenous or subcutaneous), intranasally (*e.g.*, by aspiration) or orally. Preferably, between 1 and 10 doses may be administered over a 52 week period. Preferably, 6 doses are administered, at intervals of 1 month, and booster vaccinations may be given periodically thereafter. Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of a compound that, when administered as described above, is capable of promoting an anti-tumor immune response, and is at least 10-50% above the basal (*i.e.*, untreated) level. Such response can be monitored by

measuring the anti-tumor antibodies in a patient or by vaccine-dependent generation of cytolytic effector cells capable of killing the patient's tumor cells *in vitro*. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome (e.g., more frequent remissions, complete or partial or longer disease-free survival) in vaccinated patients as compared to non-vaccinated patients. In general, for pharmaceutical compositions and vaccines comprising one or more fusion polypeptides, the amount of each fusion protein present in a dose ranges from about 1 µg to 5 mg, preferably 100 µg to 5 mg, and most preferably 5 µg to 250 µg per kg of host. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 ml to about 5 ml.

Preferably, an initial or primary immunization will be made with a Her-2/neu fusion protein having, e.g., at least one of an ECD and/or a ICD or PD, and a subsequent or booster immunization will be made with a Her-2/neu fusion protein having, e.g., at least one of a ECD and/or a ICD or PD. Preferred ECD-ICD and/or ECD-PD fusion proteins for immunization include those described herein. It will be appreciated by one skilled in the art that the present invention contemplates the use of an intact Her-2/neu fusion protein as well as division of the Her-2/neu fusion protein into a plurality of peptides. In addition, variants of the entire Her-2/neu protein, such as, e.g., the Delta Her-2/Neu protein (e.g., the mouse Delta Her-2/neu protein) described herein can also be used for immunization in the context of the present invention. Neither intact human p185^{Her-2/neu} protein nor a peptide having the amino acid sequence of the entire Her-2/neu ECD domain (or a portion of the Her-2/neu ECD domain) are used alone for immunization.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome (e.g., more frequent remissions, complete or partial, or longer disease-free survival) in treated patients as compared to non-treated patients. Increases in preexisting immune responses to a Her-2/neu protein or fusion protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

X. DETECTING CANCER

A. Methods of Detecting Cancer

In general, a cancer may be detected in a patient based on the presence of Her-2/neu proteins and/or polynucleotides encoding such proteins in a biological sample (such as blood, sera, plasma, urine and/or tumor biopsies) obtained from the patient. In other words, such proteins may be used as markers to indicate the presence or absence of a cancer such as, *e.g.*, breast, ovarian, colon, lung, prostate cancer, *etc.* The binding agents provided herein generally permit detection of the level of Her-2/neu protein that binds to the agent in the biological sample. Polynucleotide primers and probes may be used to detect the level of mRNA encoding a Her-2/neu tumor protein, which is also indicative of the presence or absence of a cancer. In general, a Her-2/neu tumor sequence should be present at a level that is at least three fold higher in tumor tissue than in normal tissue.

There are a variety of assay formats known to those of ordinary skill in the art for using a binding agent to detect polypeptide markers in a sample (*see, e.g.*, Harlow and Lane, *supra*). In general, the presence or absence of a cancer in a patient may be determined by (a) contacting a biological sample obtained from a patient with a binding agent; (b) detecting in the sample a level of polypeptide that binds to the binding agent; and (c) comparing the level of polypeptide with a predetermined cut-off value.

In a preferred embodiment, the assay involves the use of a binding agent immobilized on a solid support to bind to and remove the polypeptide from the remainder of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding agent after incubation of the binding agent with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include full length Her-2/neu tumor proteins and portions thereof to which the binding agent binds, and Her-2/neu fusion proteins and portions thereof to which the

binding agent binds, as described above.

The solid support may be any material known to those of ordinary skill in the art to which the tumor protein may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane.

5 Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the
10 patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases,
15 adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to
20 about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of a binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an
25 appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.,* Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a
30 solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a detection reagent (preferably a second antibody capable of binding to a different

site on the polypeptide) containing a reporter group is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as
5 described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered
10 saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is a period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with breast, ovarian, colon, lung or prostate cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of
15 ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody,
20 which contains a reporter group, may then be added to the solid support. Preferred reporter groups include those groups recited above.

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of
25 binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and
30 fluorescent groups. Biotin may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

To determine the presence or absence of a cancer, such as breast, ovarian, colon, lung or prostate cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value for the detection of a cancer is the average mean signal obtained when the immobilized antibody is incubated with samples from patients without the cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for the cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett *et al.*, *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-107. Briefly, in this embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (*i.e.*, sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (*i.e.*, the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for a cancer.

In a related embodiment, the assay is performed in a flow-through or strip test format, wherein the binding agent is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized binding agent as the sample passes through the membrane. A second, labeled binding agent then binds to the binding agent-polypeptide complex as a solution containing the second binding agent flows through the membrane. The detection of bound second binding agent may then be performed as described above. In the strip test format, one end of the membrane to which binding agent is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second binding agent and to the area of immobilized binding agent. Concentration of second binding agent at the area of immobilized antibody indicates the presence of a cancer. Typically, the concentration of second binding agent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of binding agent immobilized

on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferred binding agents for use in such assays are antibodies and antigen-binding fragments thereof. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the tumor proteins, or binding agents of the present invention. The above descriptions are intended to be exemplary only. For example, it will be apparent to those of ordinary skill in the art that the above protocols may be readily modified to use Her-2/neu polypeptides to detect antibodies that bind to such polypeptides in a biological sample. The detection of such Her-2/neu protein specific antibodies may correlate with the presence of a cancer.

A cancer may also, or alternatively, be detected based on the presence of T cells that specifically react with a Her-2/neu fusion protein in a biological sample. Within certain methods, a biological sample comprising CD4⁺ and/or CD8⁺ T cells isolated from a patient is incubated with a Her-2/neu fusion polypeptide, a polynucleotide encoding such a fusion polypeptide and/or an APC that expresses at least such fusion polypeptide, and the presence or absence of specific activation of the T cells is detected. Suitable biological samples include, but are not limited to, isolated T cells. For example, T cells may be isolated from a patient by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes). T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with a Her-2/neu fusion polypeptide (e.g., 5-25 μ g/ml). It may be desirable to incubate another aliquot of a T cell sample in the absence of Her-2/neu fusion polypeptide to serve as a control. For CD4⁺ T cells, activation is preferably detected by evaluating proliferation of the T cells. For CD8⁺ T cells, activation is preferably detected by evaluating cytolytic activity. A level of proliferation that is at least two fold greater and/or a level of cytolytic activity that is at least 20% greater than in disease-free patients indicates the presence of a cancer in the patient.

As noted above, a cancer may also, or alternatively, be detected based on the level of mRNA encoding a Her-2/neu protein in a biological sample. For example, at

least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify a portion of a Her-2/neu cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is specific for (*i.e.*, hybridizes to) a polynucleotide encoding the Her-2/neu protein. The amplified cDNA is then separated
5 and detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes that specifically hybridize to a polynucleotide encoding a Her-2/neu protein or fusion protein may be used in a hybridization assay to detect the presence of a polynucleotide encoding the Her-2/neu protein in a biological sample.

To permit hybridization under assay conditions, oligonucleotide primers
10 and probes should comprise an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to a portion of a polynucleotide encoding a Her-2/neu protein, or fusion protein, that is at least 10 nucleotides, and preferably at least 20 nucleotides, in length. Preferably, oligonucleotide primers and/or probes hybridize to a polynucleotide encoding a Her-2/neu protein or
15 fusion protein described herein under moderately stringent conditions, as defined above. Oligonucleotide primers and/or probes which may be usefully employed in the diagnostic methods described herein preferably are at least 10-40 nucleotides in length. In a preferred embodiment, the oligonucleotide primers comprise at least 10 contiguous nucleotides, more preferably at least 15 contiguous nucleotides, of a DNA molecule
20 having a sequence recited SEQ ID NO:9, 10 or 11. Techniques for both PCR based assays and hybridization assays are well known in the art (*see*, for example, Mullis *et al.* (1987) *Cold Spring Harbor Symp. Quant. Biol.* 51:263; Erlich ed., *PCR Technology*, Stockton Press, NY (1989)).

One preferred assay employs RT-PCR, in which PCR is applied in
25 conjunction with reverse transcription. Typically, RNA is extracted from a biological sample such as a biopsy tissue and is reverse transcribed to produce cDNA molecules. PCR amplification using at least one specific primer generates a cDNA molecule, which may be separated and visualized using, for example, gel electrophoresis. Amplification may be performed on biological samples taken from a test patient and from an individual
30 who is not afflicted with a cancer. The amplification reaction may be performed on several dilutions of cDNA spanning two orders of magnitude. A two-fold or greater increase in expression in several dilutions of the test patient sample as compared to the same dilutions of the non-cancerous sample is typically considered positive.

In another embodiment, Her-2/neu proteins or fusion proteins and

polynucleotides encoding such proteins or fusion proteins may be used as markers for monitoring the progression of cancer. In this embodiment, assays as described above for the diagnosis of a cancer may be performed over time, and the change in the level of reactive polypeptide(s) evaluated. For example, the assays may be performed every 24-
5 72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, a cancer is progressing in those patients in whom the level of polypeptide detected by the binding agent increases over time. In contrast, the cancer is not progressing when the level of reactive polypeptide either remains constant or decreases with time.

10 Certain *in vivo* diagnostic assays may be performed directly on a tumor. One such assay involves contacting tumor cells with a binding agent. The bound binding agent may then be detected directly or indirectly via a reporter group. Such binding agents may also be used in histological applications. Alternatively, polynucleotide probes may be used within such applications.

15 As noted above, to improve sensitivity, multiple Her-2/neu fusion protein markers may be assayed within a given sample. It will be apparent that binding agents specific for different proteins provided herein may be combined within a single assay. Further, multiple primers or probes may be used concurrently. The selection of tumor protein markers may be based on routine experiments to determine which combinations
20 result in optimal sensitivity. In addition, or alternatively, assays for tumor proteins provided herein may be combined with assays for other known tumor antigens.

B. Diagnostic Kits

The present invention further provides kits for use within any of the above diagnostic methods. Such kits typically comprise two or more components necessary for
25 performing a diagnostic assay. Components may be compounds, reagents, containers and/or equipment. For example, one container within a kit may contain a monoclonal antibody or fragment thereof that specifically binds to a Her-2/neu fusion protein. Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements, such as reagents or
30 buffers, to be used in the assay. Such kits may also, or alternatively, contain a detection reagent as described above that contains a reporter group suitable for direct or indirect detection of antibody binding.

Alternatively, a kit may be designed to detect the level of mRNA encoding

a Her-2/neu protein in a biological sample. Such kits generally comprise at least one oligonucleotide probe or primer, as described above, that hybridizes to a polynucleotide encoding a Her-2/neu protein or fusion protein. Such an oligonucleotide may be used, for example, within a PCR or hybridization assay. Additional components that may be
5 present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding a Her-2/neu protein.

All publications and patent applications cited in this specification are herein incorporated by reference as if each individual publication or patent application
10 were specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit
15 or scope of the appended claims.

XI. EXAMPLES

In the examples, general molecular biology reagents, such as oligonucleotide primers, lipofectamine and restriction endonucleases, were primarily obtained from Gibco/BRL (Grand Island, NY). Restriction endonucleases, Aat II and
20 PflM-I were obtained from New England Biolabs (Beverly, MA). The Her-2/neu ELISA assay kit and Her-2/neu specific monoclonal antibody Ab-3 were purchased from Oncogene Science (Manhasset, NY). The pFLAGCMV-1 expression vector and FLAG-Tag M2 antibody were purchased from Kodak (Rochester, NY). Pfu polymerase was obtained from Strategene (La Jolla, CA). The pcDNA3.1/hyg expression vector was
25 purchased from Invitrogen (Carlsbad, CA).

Example 1

CLONING OF THE ICD, KD AND PD FRAGMENTS OF HER-2/NEU INTO THE pFLAGCMV-1 EXPRESSION VECTOR

The DNA fragments of Her-2/neu that encode the intracellular domain
30 (ICD), kinase domain (KD) and phosphorylation domain (PD) were obtained separately by polymerase chain reaction. Restriction digestion sites, Hind III and Xho I, were introduced at their 5' and 3' end respectively. This design allowed the cloning of the

DNA fragments into the pFLAGCMV-1 expression vector (Kodak) in frame with a preprotrypsin leader sequence and a FLAG-Tag sequence at their N-terminus. The PCR products were gel purified and cloned into the Hind III and Sal I sites of pFLAGCMV-1. The resulting expression plasmids were designated as pFLAGCMV-1/ICD (Fig. 1),
5 pFLAGCMV-1/KD (Fig. 2) and pFLAGCMV-1/PD (Fig. 3).

Example 2

CLONING OF THE ECD-PD FUSION PROTEIN INTO THE pcDNA3.1/hyg EXPRESSION VECTOR

The DNA fragment encoding the Her-2/neu PD was amplified by
10 polymerase chain reaction. After gel purification, it was cloned into the Aat II and Xho I sites of the pT7-Her-2/neu plasmid. This procedure generated a new cloning vector, pT7/ECD-PD which linked the ECD and PD together (including a Ser from the transmembrane domain). The pT7/ECD-PD plasmid was digested with Hind-III and Xho I at 37°C for 1 hour. The 2.7 kb DNA fragment encoding the ECD-PD fusion protein was
15 gel purified and subcloned into the Hind III and Xho I sites of pcDNA3.1/hyg (Invitrogen). The resulting expression vector was designated as pcDNA3.1/hyg/ECD-PD (Fig. 5).

Example 3

EXPRESSION OF THE ICD, KD AND PD FRAGMENTS OF HER-2/NEU IN HEK-293 20 CELLS

The pFLAGCMV-1 expression plasmid (Kodak) was used to determine which region of the Her-2/neu intracellular domain could be secreted in culture media. The proteins were expressed as fusions with a preprotrypsin secretion signal and a FLAG-Tag at their N-terminus, as described in Example 1.

25 Transfection and growth of ICD, KD and PD expressing cell lines was conducted as follows: human embryonic kidney fibroblasts (HEK-293 cells) were grown in Dulbecco's modified Eagle's medium containing 10% fetal calf serum. One day prior to transfection, 1.5×10^5 cells were seeded into each well of a six-well dish. Transfection was carried out using 1 µg of plasmid-DNA by lipofectamine (Gibco/BRL) in serum free
30 media. Culture media and cells were harvested 72 hours later. For selection of stable transformants, the transfected cells were grown in media containing 200 µg/ml hygromycin.

Cells and culture media were assayed for FLAG-Tag fusion proteins by Western blot analysis as follows: culture media and cell lysates from transfected HEK-293 cells were separated on 7.5% SDS polyacrylamide gel. The proteins were transferred electrophoretically to polyvinylidene difluoride (PVDF) filters. The PVDF filters were first incubated with 5% bovine serum albumin in TBST (20 mM Tris, pH 7.5, 150 mM NaCl, 0.01% Tween 20), then incubated for 1 hour with the primary antibody, and finally incubated for another hour with peroxidase-conjugated goat anti-mouse antibody. Immunoblots were developed using ECL system (Amersham Corp.). The mouse monoclonal antibody, c-neu Ab-3 ("Ab-3") (Oncogene Science), was used for detecting the Her-2/neu protein. Ab-3 recognizes the carboxyl terminal of the human Her-2/neu protein. For detecting FLAG-Tag fusion proteins, the M2 monoclonal antibody (Kodak) was used as primary antibody in the analysis.

The results presented in Fig. 4 show that neither full length ICD nor KD were secreted, but that PD was detected in the culture media. The results indicate that the structure of KD did not permit passage of the protein through the cell membrane.

Example 4

EXPRESSION OF THE ECD-PD FUSION PROTEIN IN HEK-293 AND CHO CELLS USING THE pcDNA3.1/hyg EXPRESSION VECTOR

The ECD-PD fusion protein with a preprotrypsin secretion signal and FLAG-Tag at its N-terminus was constructed, as described in Example 2.

Transfection and growth of ECD-PD expressing cell lines was conducted as follows: HEK-293 cells and Chinese hamster ovary cells (CHO) were grown in Dulbecco's modified Eagle's medium containing 10% fetal calf serum. One day prior to transfection, 1.5×10^5 cells were seeded into each well of a six-well dish. Transfection was carried out using 1 μ g of plasmid DNA by lipofectamine (Gibco/BRL) in serum free media. Culture media and cells were harvested 72 hour later. For selection of stable transformants, the transfected cells were grown in media containing 200 μ g/ml hygromycin.

The secretion of the soluble ECD-PD fusion protein was determined by an ELISA assay with Her-2/neu ECD specific antibodies as follows: the microplate was pre-coated with the Her-2/neu specific mouse antibody (Oncogene Science) to capture Her-2/neu protein in the samples. Test samples were incubated in the microplate overnight at room temperature, incubated for 1 hour with detector antibody, and incubated for another

hour with a horseradish peroxidase-conjugated goat anti-rabbit antibody. After adding the peroxidase substrate o-phenylenediamine, a colored product was formed. The colored product was quantitated by spectrophotometry. Absorbance at 490 nm reflected the amount of Her-2/neu protein in the samples.

5 Thereafter, the secretion of the soluble ECD-PD fusion protein was determined by Western blot analysis with Her-2/neu PD specific antibodies as follows: culture media and cell lysates from transfected HEK-293 cells were separated on 7.5% SDS polyacrylamide gel. For western blot analysis, the proteins were transferred electrophoretically to PVDF filters. The PVDF filters were incubated with 5% bovine
10 serum albumin in TBST, incubated for 1 hour with the primary antibody, and incubated for another hour with peroxidase-conjugated goat anti-mouse antibody. Immunoblots were developed using the ECL system (Amersham Corp.). The mouse Ab-3 monoclonal antibody (Oncogene Science), was used for detecting the Her-2/neu protein in the present experiments. Ab-3 recognizes the carboxyl terminal of human Her-2/neu protein. For
15 detecting FLAG-Tag fusion proteins, M2 monoclonal antibody (Kodak) was used as primary antibody in the analysis. The results are presented in Fig. 6.

Example 5

CLONING OF THE HUMAN ECD-APD FUSION PROTEIN INTO THE pcDNA3.1/hyg EXPRESSION VECTOR

20 The human ECD-APD fusion protein, shown in Fig. 13 (SEQ ID NO:7), was prepared by polymerase chain reaction (PCR) with the following primers:

PDM-251 5'-cctgaatcgcgaacccaagtgtgcaccggcac-3' (SEQ ID NO:15) Tm 69°C

PDM-279 5'-ctggactcgagtcattagcgggtgcctgtgtgtg-3' (SEQ ID NO:16) Tm 69°C.

 The polymerase chain reaction conditions were: 10 µl 10x Pfu Buffer
25 (Stratagene), 1 µl 10 mM dNTPs, 2 µl 10 µM each primer, 83 µl sterile water, 1.5 µl Pfu DNA polymerase, and 50 ng template at 96°C for 2 minutes x 1 cycle; (96°C for 20 seconds, 69°C for 15 seconds, 72°C for 5 minutes) x 40 cycles; 72°C for 5 minutes x 1 cycle. The PCR product was digested with Nru I and Xho I, and cloned into a pPDM His vector (a modified pET28 vector which has a His-tag in-frame with a blunt Restriction
30 enzyme cutter Eco 72I), which was cut with Eco 72I and Xho I. The sequence was confirmed and the recombinant plasmid was then transformed into BL21 pLys S for *E. coli* expression. The plasmid construct was then digested with BamHI and XhoI and cloned into pcDNA3.1/hyg/ECD-PD which was cut with the same restriction enzymes.

Example 6

EXPRESSION IN *E. coli* of HUMAN ECD-PD-C_T-HIS TAG FUSION PROTEIN

The human ECD-PD fusion protein was cloned into the pcDNA3.1/hyg vector as described in example 2 and was used as template for constructing the hECD-PD in frame with a C-terminal 6 x histidine tag. The hECD-PD was amplified by PCR using the following primers:

AWO28 hECD-PD sense primer, with Nco I site:

5'-GGGccatgggAGCACCCAAGTGTGCACCGGC-3' (SEQ ID NO:17)

AW029 hECD-PD antisense primer, with Xho I site without stop:

10 5'-GGGctcgagCACTGGCACGTCCAGACCCAGG-3' (SEQ ID NO:18)

The PCR product was then cut with the Nco I and Xho I restriction enzymes, purified, and ligated into the pET28b expression vector linearized with the same two restriction enzymes. The ligation product was transformed into NovaBlue cells and several colonies were selected for screening. Of those, the hECD-PD.C_This clones were confirmed by DNA sequence and used for subsequent protein expression.

For protein expression, a hECD-PD.C_This clone was transformed into BL21 (DE3) CodonPlus-RIU *E. coli* competent cells. A standard mini expression screen was carried out with clones from the transformation to determine the induction yield. The best results were obtained when the cells were grown in TB media at 37°C for 2 hours.

20 The *E. coli* produced hECD-PD.C_This was then purified on a monoQ column, and refolded in 20 mM Tris-HCl (pH 8.0) buffer. The refolded protein was tested and found to be positive for Herceptin binding by Western blot and ELISA (Fig. 17). The Herceptin binding activity was, however, lost later on, probably due to denaturation of the protein.

Example 7

EXPRESSION IN *E. coli* of HUMAN N_T-HIS TAG-ECD-PD FUSION PROTEIN

The human ECD-PD fusion protein was cloned into the pPDM expression vector with a 5'Nde I and a 3' Xho I restriction sites. The ECD-PD insert was fused in frame with a N-terminal 6 x histidine tag.

30 For protein expression, the pPDM vector containing the N_This-hECD-PD fusion protein was transformed into BL21 (DE3) CodonPlus-RIU *E. coli* competent cells. A standard mini expression screen was carried out with clones from the transformation to

determine the induction yield. The best results were obtained when the cells were grown in TB media at 30°C for 2 hours.

The *E. coli* derived unpurified N_This-hECD-PD fusion protein was recognized by the mouse c-neu-3 antibody and by a rabbit anti-ECD antibody. Following purification, the *E. coli* derived N_This-hECD-PD was recognized by Herceptin both in Western blots and in ELISA assays (Fig. 17).

Example 8

EXPRESSION IN *E. coli* of MOUSE ECD-PD-C_THIS TAG FUSION PROTEIN

The mouse ECD-PD fusion protein was cloned into the pcDNA3.1 vector following a protocol similar to that described in example 2 for cloning the human ECD-PD fusion protein. This construct was then used as template for constructing the mECD-PD in frame with a C-terminal 6 x histidine tag followed by a stop codon. The internal Nco I site in the mECD-PD/pcDNA3.1 construct (base pair 1932 of the ORF) was silently mutated by site-directed mutagenesis using the following primers:

AW038 primer:

5'-GGCCCCTCCAGCCCGATGGACAGCACCTTCTACCG-3' (SEQ ID NO:19)

AW039 primer:

5'-CGGTAGAAGGTGCTGTCCATCGGGCTGGAGGGGCC-3' (SEQ ID NO:20)

After the sequence was confirmed, the mECD-PD fusion construct was amplified by PCR using the following primers:

AW036 sense primer, with Nco I restriction site:

5'-GGGGcatggGTACCCAAGTGTGTACCGG-3' (SEQ ID NO:21)

AW037 antisense primer, with Xho I restriction site:

5'-GGGGctcgagTCAATGGTGATGGTGATGGTGTCATGG
CACATCCAGGCCTAGGTACTCAGGG-3' (SEQ ID NO:22)

The PCR product was then cut with the Nco I and Xho I restriction enzymes, purified, and ligated into the pET28b expression vector linearized with the same two restriction enzymes.

The ligation product was transformed into NovaBlue and yielded multiple colonies. Four colonies were selected for sequence analysis. Of those, a mECD-PD.C_This clone which had the correct sequence was transformed into BL21 (DE3)

CodonPlus-RIU *E. coli* competent cells. A standard mini expression screen was carried out with clones from the transformation to determine the induction yield. The best results were obtained when the cells were grown in 2xYT media at 30°C for 3 hours.

Example 9

5 EXPRESSION IN *E. coli* of the Ra12-mECD-PD-C_T HIS TAG FUSION PROTEIN

The mouse ECD-PD fusion protein was cloned into the pET28b expression vector as described in Example 8. The Ra12 sequence was amplified using the following PCR fragments that added Nco I sites on the 5' and 3' ends:

10 Ra12.JC05: 5'-CCGccatggGCACGGCCGCGTCCGATAACTTCC-3'
(SEQ ID NO:23)

Ra12.JC06: 5'-GCGccatggCGGCCGGGGGTCCCTCGGCC-3' (SEQ
ID NO:24)

To obtain the Ra12 adjuvant fusion with mECD-PD, the Ra12 PCR product was then digested with the Nco I restriction enzyme, and ligated into the Nco I
15 digested and CIAP treated pET28b-mECD-PD vector. The ligation product was transformed into NovaBlue cells and yielded multiple colonies. Due to the non-directional specific ligation reaction, twice as many clones were picked for plasmid miniprep. These clones were screened by digestion with the Afi III restriction enzyme for correct orientation of the insert. The sequence of a few correctly oriented clones was
20 analyzed. One clone with a correct sequence was transformed into BL21 (DE3) CodonPlus-RIU *E. coli* competent cells for expression. A standard mini expression screen was carried out with clones from the transformation to determine the induction yield. The best results were obtained when the cells were grown in LB media at 37°C for 3 hours.

25 Example 10

EXPRESSION IN *E. coli* of LeIF.mECD-PD-C_T HIS TAG FUSION PROTEIN

The mouse ECD-PD fusion protein was cloned into the pET28b expression vector as described in Example 8. The LeIF sequence was amplified using the following PCR fragments that added Nco I sites on the 5' and 3' ends:

30 LeIF.JC03: 5'-CGCccatggCGCAGAATGATAAGATCGCCC-3'
(SEQ ID NO:25)

LeIF JC04: 5'-GCC~~ccatgg~~CGTCGCGCATGAACTTCTTCGTC-3'
(SEQ ID NO:26)

To obtain the LeIF adjuvant fusion with mECD-PD, the LeIF PCR product was then digested by the Nco I restriction enzyme, and ligated into the Nco I digested and
5 CIAP treated pET28b-mECD-PD vector. The ligation product was transformed into NovaBlue cells and yielded multiple colonies. Due to the non-directional specific ligation reaction, twice as many clones were picked for plasmid miniprep. These clones were screened by digestion with the Kpn I restriction enzyme for correct orientation of the insert. The sequence of a few correctly oriented clones was analyzed. One clone with
10 a correct sequence was transformed into BL21 (DE3) CodonPlus-RIU *E. coli* competent cells for expression. A standard mini expression screen was carried out with clones from the transformation to determine the induction yield. The best results were obtained when the cells were grown in 2xYT media at 30°C for 3 hours.

Example 11

15 EXPRESSION IN PICHIA OF AN ECD-PD FUSION PROTEIN

The ECD-PD recombinant protein used for expression in *Pichia* had the same design as for CHO expression with two modifications: (i) the native secretion signal sequence of the Her-2/neu gene had been replaced by the *Saccharomyces cerevisiae* alpha pre-pro signal sequence; and (ii) the C-terminal part of the recombinant protein was
20 elongated by one glycine and six histidines.

The ECD-PD fusion protein expression cassette was integrated into the SMD1168 *Pichia* strain using the Spheroplast method. Six multicopy integrant clones were selected among 250 clones by quantitative Dot Blot analysis. The selected clones were induced during 72 hours in Buffered Methanol-complex medium (BMMY-1%
25 methanol) in shake-flasks conditions. The six candidate clones showed the same expression profile in the cell-free supernatants and in total cellular extracts.

In the cell-free supernatants, secretion of full-length ECD-PD recombinant protein was very weak and only detected on Western Blots using the c-neu-3 mouse antibody (Calbiochem). Secretion and accumulation (maximum after 72 hours) of a \pm 70
30 kDa protein was visible on Silver stained SDS-PAGE and detected on Western blot under non-reducing conditions with Herceptin mouse antibody. This protein was not detected using the mouse c-neu-3 antibody or a mouse anti-histidine antibody (QIAGEN).

In total cellular extracts, no specific band was detected on SDS-PAGE using the Silver staining DAIICHI coloration kit. Two bands were detected on Western blots using the mouse c-neu-3 or the mouse anti-histidine antibody. One band has the same molecular size as that observed for the secreted ECD-PD product following expression in CHO cells. The other band appeared as a "smear" ranging from 100-120 kDa. These two signals could correspond to ECD-PD recombinant proteins retained in the E.R. and presenting various forms of glycosylations.

Example 12

EXPRESSION IN CHOK1 CELLS OF ECD-PD AND ECD-APD FUSION PROTEINS

The pcDNA3.1/hyg/ECD-PD and pcDNA3.1/hyg/ECD-APD plasmids were digested by the Xba I restriction enzyme and the DNA fragments encompassing the ECD-PD and ECD-APD fusion proteins were gel purified as Xba I fragments of 2783 and 2166 bp, respectively. Each fragment was transferred into the pEE14-GS vector (CellTech) linearized with Xba I (cloning site downstream of the CMV immediate early promoter). After ligation, transformation was performed into DH5 α competent *E. coli* cells. Out of sixteen colonies analyzed by restriction enzyme digestion, 2 positive colonies were found for ECD-PD, and 1 positive colony was found for ECD-APD. The obtained plasmids were prepared at large scale and purified by double CsCl-EtBr gradient centrifugation. The plasmids were analyzed by restriction enzyme digestion and sequencing of the 5' and 3' junctions between insert and vector and no abnormalities were found.

Transfection of CHO-K1 cells derived from Master Cell Bank (MCB) CHO-K1 028W 1996/2 SHF P31, growing under suspension serum free conditions, was carried out with both the pEE14-ECD-PD and the pEE14-ECD-APD plasmids using the classical DNA Ca Phosphate co-precipitation technique. Cells were counted 48 hours after transfection and transferred into 96-well plates at a density of 5000 cells/well. Transfected cells were selected according to the procedure of the glutamine synthetase (GS) expression system described by Crockett *et al.* ((1990) *Biotech.* 8:662) and amplified in the presence of 30 μ M methionine sulfoximine (MSX) in GMEM medium containing no glutamine and supplemented with additives (glutamate/asparagine/nucleosides) and 5% dialysed Fetal Bovine Serum (FBS). Cells were washed three times during the first week and twice during the second week following transfection. During the third wash, 20% conditioned medium was added.

During the fifth wash, the concentration of MSX was raised to 50 μ M to increase the level of selection.

The MSX transfectant clones were transferred 3-5 weeks after transfection into 24-well plates and the culture supernatants were harvested. Expression of the ECD-PD or ECD- Δ PD fusion proteins was tested by Western blot analysis using Herceptin antibody under non reducing conditions. Expression of the ECD-PD fusion protein was detected in 18 out of 52 clones tested, while 13 out of 47 clones tested were positive for ECD- Δ PD expression. The selected clones expressing the fusion proteins were then readapted to suspension serum-free conditions. Based on the level of expression, growth and viability, 5 clones carrying the ECD-PD construct and 3 clones carrying the ECD- Δ PD construct were further evaluated and characterized. For the ECD-PD construct, clone 560 F3 showed the highest expression level.

Expression was evaluated at 33°C in the presence or absence of sodium butyrate (2 mM) and of DMSO (2 %). Some of the clones were inducible by NaB or DMSO. Expression in CHO-K1 cells of ECD-PD and ECD- Δ PD was analyzed by Western blots and SDS-PAGE followed by either Silver or Coomassie staining. The Herceptin and the c-neu-3 mouse monoclonal antibodies, as well as the 8029K rabbit polyclonal antibody were used for Western blot analysis. Analysis of the culture supernatants from ECD-PD and ECD- Δ PD clones showed a band in Coomassie/Silver stained gels at 150 kDa and at 98 kDa, respectively. The same bands were revealed by Herceptin and by the 8029K polyclonal antiserum, as well as by the c-neu-3 antibody for ECD-PD only (Fig. 18). The CHO-expressed Her-2/neu fusion proteins are recognized by the Herceptin antibody (Fig. 18).

The expression level of the fusion proteins was also followed in terms of stability during the different cell passages. Five ECD-PD clones and two ECD- Δ PD clones were followed during passages and the stability of expression was evaluated by Western blot analysis. Out of the seven analyzed clones, four were stable after more than 32 passages, although one of them showed a high mortality and the other presented big cells.

Small scale production runs were carried out with the two best ECD-PD and ECD- Δ PD clones. Cells were cultured in suspension under serum-free conditions for 120 hours at 33°C in the presence of 2 mM Sodium butyrate. The expression of both fusion proteins was evaluated by Western blot using the Herceptin antibody and by SDS-

PAGE followed by silver staining using the Daiichi kit. Both fusion proteins were found to be expressed at +/- 100 µg/ml.

Example 13

PURIFICATION OF ECD-PD AND ECD-APD FUSION PROTEINS FOLLOWING EXPRESSION IN CHO CELLS

5

Following expression in CHO cells and secretion, the ECD-PD and ECD-APD fusion proteins were purified by anion exchange chromatography on Q sepharose High Performance columns. Before loading the supernatant onto the column, the pH was adjusted to 6.5 by adding 1N HCl. For the chromatography, 1 ml of Q Sepharose High Performance resin (Pharmacia) was used for a C10/10 column (Pharmacia).

10

The column was first equilibrated with 10 column volumes of H₂O at 4 ml/min, followed by 1 column volume of 0.5 M NaOH at 4 ml/min, and 10 column volumes of Buffer A (20 mM Bis-Tris propane pH 6.5 – 50 mM NaCl) at 4 ml/min. The sample was then loaded onto the column and allowed to pass through at a flow rate of 1 ml/min. The column was then washed with Buffer A at 1 ml/min until the O.D. at 280 nm reached 0.1, and an additional wash step of 20 column volumes was then performed. Before elution, the flow stream was reversed and an additional wash step of 3 column volumes was performed.

15

The elution was carried out at 1 ml/min, first with Buffer B (20 mM Bis-Tris propane pH 6.5-250 mM NaCl), and then with Buffer C (20 mM Bis-Tris propane pH 6.5-1 mM NaCl). The fusion proteins of interest were eluted with buffer B.

20

The fusion proteins were further purified by hydrophobic chromatography on Phenyl Sepharose 6 Fast Flow low substitution. The eluate containing the ECD-PD and ECD-APD fusion proteins (Buffer B eluate) was adjusted to obtain a concentration of 1M ammonium sulfate (AMS) by addition of solid AMS (140 g/liter of solution). The pH of the solution was checked to be 7.0.

25

For the chromatography, 0.5 ml of Phenyl Sepharose 6 Fast Flow low substitution (Pharmacia) was used with a C10/10 (Pharmacia). The column was then equilibrated with 10 column volumes of H₂O, at 4 ml/min, 1 column volume of 0.5 M NaOH, at 4 ml/min, and 10 column volumes of buffer D (1 mM PO₄ pH 7.0 – 1M AMS), at 4 ml/min. Following equilibration, the sample was loaded and allowed to pass through the column at a flow rate of 0.5 ml/min. The column was then washed in Buffer D at 0.5 ml/min until the O.D. at 280 nm reached the baseline, and then at 1 ml/min for 10 column

30

volumes. Before elution, the flow stream was reversed and an additional wash step of 3 column volumes was performed. Elution was carried out at 1 ml/min with buffer E (1 mM PO₄ pH 7.0).

The purified fusion proteins were analyzed by SDS-PAGE followed by silver staining using the Daiichi kit, and by Western blot, using the 8029K rabbit polyclonal antibody or the mouse Herceptin antibody. The analysis showed that the level of purity following the two purification steps was estimated at +/- 90% by densitometry (Biorad GS-700 Imaging Densitometer). The Western blot analysis showed that the monomers remained the major band all along the purification, that the level of oxydation was not increased, and that the detection of the epitope of interest was not modified by the conditions of purification, as shown by using the Herceptin antibody. The total amount of each fusion protein recovered was measured using a colorimetric protein assay (DOC TCA BCA). This assays estimated that 2 and 4 mg of ECD-PD and ECD-ΔPD fusion protein, respectively, were purified from 75 ml of culture, with a level of purity of +/- 90%.

Example 14

EXPRESSION IN MAMMALIAN CELLS OF THE MOUSE ECD-PD AND ECD-PDTcP0 FUSION PROTEINS

Transiently transfected mammalian HEK293 cells were generated that expressed full-length mouse Neu, Delta mouse Neu, mouse ECD-PD and mouse ECD-PD-TcP0 recombinant proteins. Stable HEK clones that express mouse ECDPD and mouse ECD-PD-TcP0 recombinant proteins were also generated.

For recombinant expression in mammalian cells, various forms of mouse Neu (mNeu) cDNAs were subcloned into mammalian expression vectors. Full-length mNeu (Fig. 19) was subcloned into pBIB and VR1012 vectors, and delta mNeu, which contained an engineered A-T basepair change (and subsequent lysine to methionine amino acid change) at position 2261 of the mNeu cDNA, was subcloned into the pBIB vector. A cDNA encoding the extracellular domain (ECD) and phosphorylation domain (PD), and a cDNA encoding the ECD, PD and 6 amino acid TcP0 domains were generated by PCR amplification using full-length mouse Neu cDNA as template. These ECD-PD and ECD-PD-TcP0 products were then subcloned into the mammalian expression vector pcDNA3.1 (Invitrogen) that was modified to contain the DHFR genomic fragment. All constructs were transfected into HEK293 cells (ATCC) using

Fugene 6 reagent (Roche). Briefly, the HEK cells were plated at a density of 100,000 cells/ml in DMEM (Gibco) containing 10% FBS (Hyclone) and grown overnight. The following day, 2 μ l of Fugene6 was added to 100 μ l of DMEM containing no FBS and incubated for 15 minutes at RT. The Fugene6/DMEM mixture was then added to 1 μ g of mNeu/expression vector DNA and incubated for 15 minutes at RT. The Fugene/DNA mix was then added to the HEK293 cells and incubated for 48-72 hours at 37°C with 7% CO₂. Cells were rinsed with PBS and then collected and pelleted by centrifugation. For Western blot analysis, whole cell lysates were generated by incubating the cells in Triton-X 100 containing lysis buffer for 30 minutes on ice. Lysates were then cleared by centrifugation at 10,000 rpm for 5 minutes at 4°C. Samples were diluted with SDS-PAGE loading buffer containing beta-mercaptoethanol, then boiled for 10 minutes prior to loading the SDS-PAGE gel. The proteins were transferred to nitrocellulose and probed using the mouse monoclonal Ab-3 (c-neu 3; CalBiochem) at a concentration of 1 μ g/ml. The blot was revealed with a goat anti-mouse Ig coupled to HRP followed by incubation in ECL substrate (Fig. 21).

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for the purpose of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

WHAT IS CLAIMED IS:

- 1 1. An isolated protein comprising a Her-2/neu extracellular domain
2 fused to a Her-2/neu phosphorylation domain, wherein the protein is capable of producing
3 an immune response in a warm-blooded animal.
- 1 2. The protein of claim 1, wherein the protein has a sequence at least
2 80% identical to the sequence of SEQ ID NO:6, or wherein the protein comprises a
3 sequence at least 80% identical to the sequence of SEQ ID NO:3 fused to a sequence at
4 least 80% identical to the sequence of SEQ ID NO:4.
- 1 3. The protein of claim 1, wherein the protein comprises a sequence at
2 least 80 % identical to the sequence of SEQ ID NO:3 directly fused to an amino acid
3 sequence at least 80% identical to the sequence inclusive of Gln 991 to Val 1256 of SEQ
4 ID NO:2, or wherein the protein comprises a sequence at least 80 % identical to the
5 sequence of SEQ ID NO:3 fused to the amino acid sequence at least 80% identical to the
6 sequence inclusive of Gln 991 to Val 1256 of SEQ ID NO:2.
- 1 4. The protein of claim 1, wherein the protein comprises a sequence at
2 least 80% identical to the sequence of SEQ ID NO:8 directly fused to a sequence at least
3 80% identical to the sequence of SEQ ID NO:4, or wherein the protein comprises a
4 sequence at least 80% identical to the sequence of SEQ ID NO:8 fused to a sequence at
5 least 80% identical to the sequence of SEQ ID NO:4.
- 1 5. The protein of claim 1, wherein the protein comprises a sequence at
2 least 80% identical to the sequence of SEQ ID NO:8 directly fused to the amino acid
3 sequence inclusive of Gln 991 to Val 1256 of SEQ ID NO:2, or wherein the protein
4 comprises a sequence at least 80% identical to the sequence of SEQ ID NO:8 fused to a
5 sequence at least 80% identical to the amino acid sequence inclusive of Gln 991 to Val
6 1256 of SEQ ID NO:2.
- 1 6. The protein of claim 1, wherein the protein has a sequence at least
2 80% identical to the sequence of SEQ ID NO:27.
- 1 7. The protein of claim 1, wherein the Her-2/neu extracellular domain
2 is fused to the Her-2/neu phosphorylation domain via a chemical linker.

- 1 8. The protein of claim 7, wherein the chemical linker is an amino
2 acid linker.
- 1 9. The protein of claim 1, wherein the protein further comprises a
2 peptide having the sequence set forth in SEQ ID NO:31.
- 1 10. The protein of claim 9, wherein said peptide is fused to the C-
2 terminus of the Her-2/neu phosphorylation domain.
- 1 11. The protein of claim 10, wherein the protein comprises a sequence
2 at least 80% identical to the sequence set forth in SEQ ID NO:29.
- 1 12. The protein of claim 1, wherein the protein further comprises an
2 NS1 antigen or an immunogenic fragment thereof.
- 1 13. The protein of claim 12, wherein the NS1 antigen or the
2 immunogenic fragment thereof is fused to the N-terminus of the protein.
- 1 14. A nucleic acid molecule encoding the protein of claim 1.
- 1 15. A viral vector comprising a polynucleotide sequence encoding the
2 protein of claim 1.
- 1 16. The viral vector of claim 15, wherein the viral vector is an
2 adenovirus.
- 1 17. The viral vector of claim 15, wherein the viral vector is an
2 alphavirus.
- 1 18. A pharmaceutical composition comprising the protein of claim 1,
2 and a pharmaceutically acceptable carrier or diluent.
- 1 19. The pharmaceutical composition of claim 18, wherein the
2 pharmaceutical composition is a vaccine.
- 1 20. The pharmaceutical composition of claim 18, further comprising an
2 immunostimulatory substance.

1 21. The pharmaceutical composition of claim 20, wherein the protein is
2 presented in an oil-in-water emulsion.

1 22. The pharmaceutical composition of claim 20, wherein the
2 immunostimulatory substance is SBAS-2, SBAS-4, 3D-MPL, QS21, or a combination of
3 3D-MPL and QS21.

1 23. A pharmaceutical composition comprising the nucleic acid
2 molecule of claim 14, and a pharmaceutically acceptable carrier or diluent.

1 24. The pharmaceutical composition of claim 23, wherein the
2 pharmaceutical composition is a vaccine.

1 25. The pharmaceutical composition of claim 23, further comprising an
2 immunostimulatory substance.

1 26. The pharmaceutical composition of claim 23, wherein the nucleic
2 acid molecule is a DNA molecule.

1 27. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the protein of claim 1 in an amount effective to elicit or enhance the immune response.

1 28. The method of claim 27, wherein the protein is administered in the
2 form of a vaccine.

1 29. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the nucleic acid molecule of claim 14 in an amount effective to elicit or enhance the
4 immune response.

1 30. The method of claim 29, wherein the nucleic acid molecule is in
2 the form of a vaccine.

1 31. The method of claim 29, wherein the step of administering
2 comprises transfecting cells of the warm-blooded animal *ex vivo* with the nucleic acid
3 molecule and subsequently delivering the transfected cells to the warm-blooded animal.

1 32. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the viral vector of claim 15 in an amount effective to elicit or enhance the immune
4 response.

1 33. The method of claim 32, wherein the viral vector is an adenovirus.

1 34. The method of claim 32, wherein the viral vector is an alphavirus.

1 35. The method of claim 32, wherein the step of administering
2 comprises infecting cells of the warm-blooded animal *ex vivo* with the viral vector and
3 subsequently delivering the infected cells to the warm-blooded animal.

1 36. An isolated protein comprising a Her-2/neu extracellular domain
2 fused to a fragment of the Her-2/neu phosphorylation domain, wherein the protein is
3 capable of producing an immune response in a warm-blooded animal.

1 37. The protein of claim 36, wherein the protein has a sequence at least
2 80% identical to the sequence of SEQ ID NO:7, or wherein the protein comprises a
3 sequence at least 80% identical to the sequence of SEQ ID NO:3 fused to a sequence at
4 least 80% identical to the sequence of SEQ ID NO:5.

1 38. The protein of claim 36, wherein the protein comprises a sequence
2 at least 80% identical to the sequence of SEQ ID NO:3 directly fused to a sequence at
3 least 80% identical to the amino acid sequence inclusive of Gln 991 to Arg 1049 of SEQ
4 ID NO:2, or wherein the protein comprises a sequence at least 80% identical to the
5 sequence of SEQ ID NO:3 fused to a sequence at least 80% identical to the amino acid
6 sequence inclusive of Gln 991 to Arg 1049 of SEQ ID NO:2.

1 39. The protein of claim 36, wherein the protein comprises a sequence
2 at least 80% identical to the sequence of SEQ ID NO:8 directly fused to a sequence at
3 least 80% identical to the sequence of SEQ ID NO:5, or wherein the protein comprises a
4 sequence at least 80% identical to the sequence of SEQ ID NO:8 fused to a sequence at
5 least 80% identical to the sequence of SEQ ID NO:5.

1 40. The protein of claim 36, wherein the protein comprises a sequence
2 at least 80% identical to the sequence of SEQ ID NO:8 directly fused to a sequence at
3 least 80% identical to the amino acid sequence inclusive of Gln 991 to Arg 1049 of SEQ
4 ID NO:2, or wherein the protein comprises a sequence at least 80% identical to the
5 sequence of SEQ ID NO:8 fused to a sequence at least 80% identical to the amino acid
6 sequence inclusive of Gln 991 to Arg 1049 of SEQ ID NO:2.

1 41. The protein of claim 36, wherein the Her-2/neu extracellular
2 domain is fused to the fragment of the Her-2/neu phosphorylation domain via a chemical
3 linker.

1 42. The protein of claim 41, wherein the chemical linker is an amino
2 acid linker.

1 43. The protein of claim 36, wherein the protein further comprises a
2 peptide having the sequence set forth in SEQ ID NO:31.

1 44. The protein of claim 43, wherein said peptide is fused to the C-
2 terminus of the fragment of the Her-2/neu phosphorylation domain.

1 45. A nucleic acid molecule encoding the protein of claim 36.

1 46. A viral vector comprising a polynucleotide sequence encoding the
2 protein of claim 36.

1 47. The viral vector of claim 46, wherein the viral vector is an
2 adenovirus.

1 48. The viral vector of claim 46, wherein the viral vector is an
2 alphavirus.

1 49. A pharmaceutical composition comprising the protein of claim 36,
2 and a pharmaceutically acceptable carrier or diluent.

1 50. The pharmaceutical composition of claim 49, wherein the
2 pharmaceutical composition is a vaccine.

1 51. The pharmaceutical composition of claim 49, further comprising an
2 immunostimulatory substance.

1 52. The pharmaceutical composition of claim 51, wherein the protein is
2 presented in an oil-in-water emulsion.

1 53. The pharmaceutical composition of claim 51, wherein the
2 immunostimulatory substance is SBAS-2, SBAS-4, 3D-MPL, QS21, or a combination of
3 3D-MPL and QS21.

1 54. A pharmaceutical composition comprising the nucleic acid
2 molecule of claim 45, and a pharmaceutically acceptable carrier or diluent.

1 55. The pharmaceutical composition of claim 54, wherein the
2 pharmaceutical composition is a vaccine.

1 56. The pharmaceutical composition of claim 54, further comprising an
2 immunostimulatory substance.

1 57. The pharmaceutical composition of claim 54, wherein the nucleic
2 acid molecule is a DNA molecule.

1 58. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the protein of claim 36 in an amount effective to elicit or enhance the immune response.

1 59. The method of claim 58, wherein the protein is administered in the
2 form of a vaccine.

1 60. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the nucleic acid molecule of claim 45 in an amount effective to elicit or enhance the
4 immune response.

1 61. The method of claim 60, wherein the nucleic acid molecule is in
2 the form of a vaccine.

1 62. The method of claim 60, wherein the step of administering
2 comprises transfecting cells of the warm-blooded animal *ex vivo* with the nucleic acid
3 molecule and subsequently delivering the transfected cells to the warm-blooded animal.

1 63. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the viral vector of claim 46 in an amount effective to elicit or enhance the immune
4 response.

1 64. The method of claim 63, wherein the viral vector is an adenovirus.

1 65. The method of claim 63, wherein the viral vector is an alphavirus.

1 66. The method of claim 63, wherein the step of administering
2 comprises infecting cells of the warm-blooded animal *ex vivo* with the viral vector and
3 subsequently delivering the infected cells to the warm-blooded animal.

1 67. An isolated protein comprising a Her-2/neu extracellular domain
2 fused to a Her-2/neu intracellular domain, wherein the protein is capable of producing an
3 immune response in a warm-blooded animal.

1 68. The protein of claim 67, wherein the protein comprises a sequence
2 at least 80% identical to the sequence of SEQ ID NO:3 fused directly to a sequence at
3 least 80% identical to the amino acid sequence inclusive of Lys 676 to Val 1255 in SEQ
4 ID NO:1, or wherein the protein comprises a sequence at least 80% identical to the
5 sequence of SEQ ID NO:3 fused to a sequence at least 80% identical to the amino acid
6 sequence inclusive of Lys 676 to Val 1255 of SEQ ID NO:1 via at least one of a chemical
7 or amino acid linking group.

1 69. The protein of claim 67, wherein the protein comprises a sequence
2 at least 80% identical to the sequence of SEQ ID NO:3 directly fused to a sequence at
3 least 80% identical to the amino acid sequence inclusive of Lys 677 to Val 1256 of SEQ
4 ID NO:2, or wherein the protein comprises a sequence at least 80% identical to the
5 sequence of SEQ ID NO:3 fused to a sequence at least 80% identical to the amino acid
6 sequence inclusive of Lys 677 to Val 1256 of SEQ ID NO:2 via at least one of a chemical
7 or amino acid linking group.

1 70. The protein of claim 67, wherein the protein comprises a sequence
2 at least 80% identical to the sequence of SEQ ID NO:8 directly fused to a sequence at
3 least 80% identical to the amino acid sequence inclusive of Lys 676 to Val 1255 of SEQ
4 ID NO:1, or wherein the protein comprises a sequence at least 80% identical to the
5 sequence of SEQ ID NO:8 fused to a sequence at least 80% identical to the amino acid
6 sequence inclusive of Lys 676 to Val 1255 of SEQ ID NO:1 via at least one of a chemical
7 or amino acid linking group.

1 71. The protein of claim 67, wherein the protein comprises a sequence
2 at least 80% identical to the sequence of SEQ ID NO:8 directly fused to a sequence at
3 least 80% identical to the amino acid sequence inclusive of Lys 677 to Val 1256 of SEQ
4 ID NO:2, or wherein the protein comprises a sequence at least 80% identical to the
5 sequence of SEQ ID NO:8 fused to a sequence at least 80% identical to the amino acid
6 sequence inclusive of Lys 677 to Val 1256 of SEQ ID NO:2 via at least one of a chemical
7 or amino acid linking group.

1 72. The protein of claim 67, wherein the Her-2/neu extracellular
2 domain is fused to the Her-2/neu intracellular domain via a chemical linker.

1 73. The protein of claim 72, wherein the chemical linker is an amino
2 acid linker.

1 74. The protein of claim 67, wherein the protein further comprises a
2 peptide having the sequence set forth in SEQ ID NO:31.

1 75. The protein of claim 74, wherein said peptide is fused to the C-
2 terminus of the Her-2/neu intracellular domain.

1 76. A nucleic acid molecule encoding the protein of claim 67.

1 77. A viral vector comprising a polynucleotide sequence encoding the
2 protein of claim 67.

1 78. The viral vector of claim 77, wherein the viral vector is an
2 adenovirus.

1 79. The viral vector of claim 77, wherein the viral vector is an
2 alphavirus.

1 80. A pharmaceutical composition comprising the protein of claim 67,
2 and a pharmaceutically acceptable carrier or diluent.

1 81. The pharmaceutical composition of claim 80, wherein the
2 pharmaceutical composition is a vaccine.

1 82. The pharmaceutical composition of claim 80, further comprising an
2 immunostimulatory substance.

1 83. The pharmaceutical composition of claim 82, wherein the protein is
2 presented in an oil-in-water emulsion.

1 84. The pharmaceutical composition of claim 82, wherein the
2 immunostimulatory substance is SBAS-2, SBAS-4, 3D-MPL, QS21, or a combination of
3 3D-MPL and QS21.

1 85. A pharmaceutical composition comprising the nucleic acid
2 molecule of claim 76, and a pharmaceutically acceptable carrier or diluent.

1 86. The pharmaceutical composition of claim 85, wherein the
2 pharmaceutical composition is a vaccine.

1 87. The pharmaceutical composition of claim 85, further comprising an
2 immunostimulatory substance.

1 88. The pharmaceutical composition of claim 85, wherein the nucleic
2 acid molecule is a DNA molecule.

1 89. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the protein of claim 67 in an amount effective to elicit or enhance the immune response.

1 90. The method of claim 89, wherein the protein is administered in the
2 form of a vaccine.

1 91. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the nucleic acid molecule of claim 76 in an amount effective to elicit or enhance the
4 immune response.

1 92. The method of claim 91, wherein the nucleic acid molecule is in
2 the form of a vaccine.

1 93. The method of claim 91, wherein the step of administering
2 comprises transfecting cells of the warm-blooded animal *ex vivo* with the nucleic acid
3 molecule and subsequently delivering the transfected cells to the warm-blooded animal.

1 94. A method for eliciting or enhancing an immune response to Her-
2 2/neu protein, the method comprising the step of administering to a warm-blooded animal
3 the viral vector of claim 77 in an amount effective to elicit or enhance the immune
4 response.

1 95. The method of claim 94, wherein the viral vector is an adenovirus.

1 96. The method of claim 94, wherein the viral vector is an alphavirus.

1 97. The method of claim 94, wherein the step of administering
2 comprises infecting cells of the warm-blooded animal *ex vivo* with the viral vector and
3 subsequently delivering the infected cells to the warm-blooded animal.

1 98. A method for inhibiting the development of a cancer in a patient,
2 the method comprising the step of administering to a patient an effective amount of a
3 fusion polypeptide according to claim 1, 36, or 67 and thereby inhibiting the development
4 of a cancer in the patient.

1 99. A method for inhibiting the development of a cancer in a patient,
2 the method comprising the step of administering to a patient an effective amount of a
3 polynucleotide according to claim 14, 45, or 76 and thereby inhibiting the development of
4 a cancer in the patient.

1 100. A method for inhibiting the development of a cancer in a patient,
2 the method comprising the step of administering to a patient an effective amount of an

3 antigen-presenting cell that expresses a fusion polypeptide according to claim 1, 36, or
4 67, and thereby inhibiting the development of a cancer in the patient.

1 101. A method according to claim 100, wherein the antigen-presenting
2 cell is a dendritic cell.

1 102. A method according to any one of claims 98-101, wherein the
2 cancer is breast, ovarian, colon, lung or prostate cancer.

1 103. A method for removing tumor cells from a biological sample, the
2 method comprising the step of contacting a biological sample with T cells that
3 specifically react with a Her-2/neu fusion protein, wherein the fusion protein comprises
4 an amino acid sequence that is encoded by a polynucleotide sequence selected from the
5 group consisting of:

6 (i) polynucleotides of claims 14, 45, or 76; and

7 (ii) complements of the foregoing polynucleotides;

8 wherein the step of contacting is performed under conditions and for a
9 time sufficient to permit the removal of cells expressing the antigen from the sample.

1 104. A method according to claim 103, wherein the biological sample is
2 blood or a fraction thereof.

1 105. A method for inhibiting the development of a cancer in a patient,
2 comprising the step of administering to a patient a biological sample treated according to
3 the method of claim 103.

1 106. A method for stimulating and/or expanding T cells specific for a
2 Her-2/neu fusion protein, the method comprising the step of contacting T cells with one
3 or more of:

4 (i) a fusion protein according to claims 1, 36, or 67;

5 (ii) a polynucleotide encoding such a fusion protein; or

6 (iii) an antigen presenting cell that expresses such a fusion protein;

7 under conditions and for a time sufficient to permit the stimulation and/or
8 expansion of T cells.

1 107. An isolated T cell population, comprising T cells prepared
2 according to the method of claim 106.

1 108. A method for inhibiting the development of a cancer in a patient,
2 the method comprising the step of administering to a patient an effective amount of a T
3 cell population according to claim 107.

1 109. A method for inhibiting the development of a cancer in a patient,
2 the method comprising the steps of:

3 (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with
4 at least one component selected from the group consisting of:

- 5 (i) a fusion protein according to claims 1, 36, or 67;
6 (ii) a polynucleotide encoding such a fusion protein; and
7 (iii) an antigen-presenting cell that expresses such a fusion
8 protein;

9 such that T cells proliferate; and

10 (b) administering to the patient an effective amount of the proliferated
11 T cells, thereby inhibiting the development of a cancer in the patient.

1 110. A method for inhibiting the development of a cancer in a patient,
2 the method comprising the steps of:

3 (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with
4 at least one component selected from the group consisting of:

- 5 (i) a fusion protein according to claims 1, 36, or 67;
6 (ii) a polynucleotide encoding such a fusion protein; and
7 (iii) an antigen-presenting cell that expresses such a fusion
8 protein;

9 such that T cells proliferate;

10 (b) cloning at least one proliferated cell; and

11 (c) administering to the patient an effective amount of the cloned T
12 cells, thereby inhibiting the development of a cancer in the patient.

1 111. A method of making a fusion protein according to claims 1, 36, or
2 67, the method comprising the steps of:

- 3 (a) introducing into a cell an expression vector comprising a
4 polynucleotide according to claims 14, 45, or 76;
5 (b) culturing the transfected cell; and
6 (c) purifying the expressed protein.
- 1 112. The method of claim 111, wherein the cell is a CHO cell.
- 1 113. The method of claim 111, wherein the cell is a BHK cell.
- 1 114. The method of claim 111, wherein the cell is cultured in
2 suspension, under serum-free conditions.
- 1 115. The method of claim 111, wherein the expressed protein is purified
2 by a two-step procedure, the procedure comprising:
3 (a) anion exchange chromatography on Q sepharose High Performance
4 Columns; and
5 (b) hydrophobic chromatography on Phenyl Sepharose 6 Fast Flow
6 low substitution.


**Fig. 1**

Fig. 2




Fig. 3





Fig. 4






Fig. 5

pcDNA3.1hyg/ECD-PD expression




Fig. 6A





Fig. 6B



																			10																				20
Met	Glu	Leu	Ala	Ala	Leu	Cys	Arg	Trp	Gly	Leu	Leu	Leu	Ala	Leu	Leu	Pro	Pro	Gly	Ala	20																			
Ala	Ser	Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys	Leu	Arg	Leu	Pro	Ala	Ser	Pro	Glu	40																			
Thr	His	Leu	Asp	Met	Leu	Arg	His	Leu	Tyr	Gln	Gly	Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	60																			
Glu	Leu	Thr	Tyr	Leu	Pro	Thr	Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val	80																			
Gln	Gly	Tyr	Val	Leu	Ile	Ala	His	Asn	Gln	Val	Arg	Gln	Val	Pro	Leu	Gln	Arg	Leu	Arg	100																			
																			110																				120
Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Asn	Tyr	Ala	Leu	Ala	Val	Leu	Asp	Asn	Gly	120																			
Asp	Pro	Leu	Asn	Asn	Thr	Thr	Pro	Val	Thr	Gly	Ala	Ser	Pro	Gly	Gly	Leu	Arg	Glu	Leu	140																			
Gln	Leu	Arg	Ser	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Gln	Arg	Asn	Pro	Gln	160																			
Leu	Cys	Tyr	Gln	Asp	Thr	Ile	Leu	Trp	Lys	Asp	Ile	Phe	His	Lys	Asn	Asn	Gln	Leu	Ala	180																			
Leu	Thr	Leu	Ile	Asp	Thr	Asn	Arg	Ser	Arg	Ala	Cys	His	Pro	Cys	Ser	Pro	Met	Cys	Lys	200																			
																			210																				220
Gly	Ser	Arg	Cys	Trp	Gly	Glu	Ser	Ser	Glu	Asp	Cys	Gln	Ser	Leu	Thr	Arg	Thr	Val	Cys	220																			
Ala	Gly	Gly	Cys	Ala	Arg	Cys	Lys	Gly	Pro	Leu	Pro	Thr	Asp	Cys	Cys	His	Glu	Gln	Cys	240																			
Ala	Ala	Gly	Cys	Thr	Gly	Pro	Lys	His	Ser	Asp	Cys	Leu	Ala	Cys	Leu	His	Phe	Asn	His	260																			
Ser	Gly	Ile	Cys	Glu	Leu	His	Cys	Pro	Ala	Leu	Val	Thr	Tyr	Asn	Thr	Asp	Thr	Phe	Glu	280																			
Ser	Met	Pro	Asn	Pro	Glu	Gly	Arg	Tyr	Thr	Phe	Gly	Ala	Ser	Cys	Val	Thr	Ala	Cys	Pro	300																			
																			310																				320
Tyr	Asn	Tyr	Leu	Ser	Thr	Asp	Val	Gly	Ser	Cys	Thr	Leu	Val	Cys	Pro	Leu	His	Asn	Gln	320																			
Glu	Val	Thr	Ala	Glu	Asp	Gly	Thr	Gln	Arg	Cys	Glu	Lys	Cys	Ser	Lys	Pro	Cys	Ala	Arg	340																			
Val	Cys	Tyr	Gly	Leu	Gly	Met	Glu	His	Leu	Arg	Glu	Val	Arg	Ala	Val	Thr	Ser	Ala	Asn	360																			
Ile	Gln	Glu	Phe	Ala	Gly	Cys	Lys	Lys	Ile	Phe	Gly	Ser	Leu	Ala	Phe	Leu	Pro	Glu	Ser	380																			
Phe	Asp	Gly	Asp	Pro	Ala	Ser	Asn	Thr	Ala	Pro	Leu	Gln	Pro	Glu	Gln	Leu	Gln	Val	Phe	400																			
																			410																				420
Glu	Thr	Leu	Glu	Glu	Ile	Thr	Gly	Tyr	Leu	Tyr	Ile	Ser	Ala	Trp	Pro	Asp	Ser	Leu	Pro	420																			
Asp	Leu	Ser	Val	Phe	Gln	Asn	Leu	Gln	Val	Ile	Arg	Gly	Arg	Ile	Leu	His	Asn	Gly	Ala	440																			
Tyr	Ser	Leu	Thr	Leu	Gln	Gly	Leu	Gly	Ile	Ser	Trp	Leu	Gly	Leu	Arg	Ser	Leu	Arg	Glu	460																			
Leu	Gly	Ser	Gly	Leu	Ala	Glu	Ile	His	His	Asn	Thr	His	Leu	Cys	Phe	Val	His	Thr	Val	480																			
Pro	Trp	Asp	Gln	Leu	Phe	Arg	Asn	Pro	His	Gln	Ala	Leu	Leu	His	Thr	Ala	Asn	Arg	Pro	500																			
																			510																				520
Glu	Asp	Glu	Cys	Val	Gly	Glu	Gly	Leu	Ala	Cys	His	Gln	Leu	Cys	Ala	Arg	Gly	His	Cys	520																			
Trp	Gly	Pro	Gly	Pro	Thr	Gln	Cys	Val	Asn	Cys	Ser	Gln	Phe	Leu	Arg	Gly	Gln	Glu	Cys	540																			
Val	Glu	Glu	Cys	Arg	Val	Leu	Gln	Gly	Leu	Pro	Arg	Glu	Tyr	Val	Asn	Ala	Arg	His	Cys	560																			
Leu	Pro	Cys	His	Pro	Glu	Cys	Gln	Pro	Gln	Asn	Gly	Ser	Val	Thr	Cys	Phe	Gly	Pro	Glu	580																			
Ala	Asp	Gln	Cys	Val	Ala	Cys	Ala	His	Tyr	Lys	Asp	Pro	Pro	Phe	Cys	Val	Ala	Arg	Cys	600																			

Fig. 7 (SEQ ID NO: 1)

610																			620			
Pro	Ser	Gly	Val	Lys	Pro	Asp	Leu	Ser	Tyr	Met	Pro	Ile	Trp	Lys	Phe	Pro	Asp	Glu	Glu	620		
Gly	Ala	Cys	Gln	Pro	Cys	Pro	Ile	Asn	Cys	Thr	His	Ser	Cys	Val	Asp	Leu	Asp	Lys		640		
Gly	Cys	Pro	Ala	Glu	Gln	Arg	Ala	Ser	Pro	Leu	Thr	Ser	Ile	Ile	Ser	Ala	Val	Val	Gly	660		
Ile	Leu	Leu	Val	Val	Val	Leu	Gly	Val	Val	Phe	Gly	Ile	Leu	Ile	Lys	Arg	Arg	Gln	Gln	680		
Lys	Ile	Arg	Lys	Tyr	Thr	Met	Arg	Arg	Leu	Leu	Gln	Glu	Thr	Glu	Leu	Val	Glu	Pro	Leu	700		
710																			720			
Thr	Pro	Ser	Gly	Ala	Met	Pro	Asn	Gln	Ala	Gln	Met	Arg	Ile	Leu	Lys	Glu	Thr	Glu	Leu	720		
Arg	Lys	Val	Lys	Val	Leu	Gly	Ser	Gly	Ala	Phe	Gly	Thr	Val	Tyr	Lys	Gly	Ile	Trp	Ile	740		
Pro	Asp	Gly	Glu	Asn	Val	Lys	Ile	Pro	Val	Ala	Ile	Lys	Val	Leu	Arg	Glu	Asn	Thr	Ser	760		
Pro	Lys	Ala	Asn	Lys	Glu	Ile	Leu	Asp	Glu	Ala	Tyr	Val	Met	Ala	Gly	Val	Gly	Ser	Pro	780		
Tyr	Val	Ser	Arg	Leu	Leu	Gly	Ile	Cys	Leu	Thr	Ser	Thr	Val	Gln	Leu	Val	Thr	Gln	Leu	800		
810																			820			
Met	Pro	Tyr	Gly	Cys	Leu	Leu	Asp	His	Val	Arg	Glu	Asn	Arg	Gly	Arg	Leu	Gly	Ser	Gln	820		
Asp	Leu	Leu	Asn	Trp	Cys	Met	Gln	Ile	Ala	Lys	Gly	Met	Ser	Tyr	Leu	Glu	Asp	Val	Arg	840		
Leu	Val	His	Arg	Asp	Leu	Ala	Ala	Arg	Asn	Val	Leu	Val	Lys	Ser	Pro	Asn	His	Val	Lys	860		
Ile	Thr	Asp	Phe	Gly	Leu	Ala	Arg	Leu	Leu	Asp	Ile	Asp	Glu	Thr	Glu	Tyr	His	Ala	Asp	880		
Gly	Gly	Lys	Val	Pro	Ile	Lys	Trp	Met	Ala	Leu	Glu	Ser	Ile	Leu	Arg	Arg	Arg	Phe	Thr	900		
910																			920			
His	Gln	Ser	Asp	Val	Trp	Ser	Tyr	Gly	Val	Thr	Val	Trp	Glu	Leu	Met	Thr	Phe	Gly	Ala	920		
Lys	Pro	Tyr	Asp	Gly	Ile	Pro	Ala	Arg	Glu	Ile	Pro	Asp	Leu	Leu	Glu	Lys	Gly	Glu	Arg	940		
Leu	Pro	Gln	Pro	Pro	Ile	Cys	Thr	Ile	Asp	Val	Tyr	Met	Ile	Met	Val	Lys	Cys	Trp	Met	960		
Ile	Asp	Ser	Glu	Cys	Arg	Pro	Arg	Phe	Arg	Glu	Leu	Val	Ser	Glu	Phe	Ser	Arg	Met	Ala	980		
Arg	Asp	Pro	Gln	Arg	Phe	Val	Val	Ile	Gln	Asn	Glu	Asp	Leu	Gly	Pro	Ala	Ser	Pro	Leu	1000		
1010																			1020			
Asp	Ser	Thr	Phe	Tyr	Arg	Ser	Leu	Leu	Glu	Asp	Asp	Asp	Met	Gly	Asp	Leu	Val	Asp	Ala	1020		
Glu	Glu	Tyr	Leu	Val	Pro	Gln	Gln	Gly	Phe	Phe	Cys	Pro	Asp	Pro	Ala	Pro	Gly	Ala	Gly	1040		
Gly	Met	Val	His	His	Arg	His	Arg	Ser	Ser	Ser	Thr	Arg	Ser	Gly	Gly	Gly	Asp	Leu	Thr	1060		
Leu	Gly	Leu	Glu	Pro	Ser	Glu	Glu	Glu	Ala	Pro	Arg	Ser	Pro	Leu	Ala	Pro	Ser	Glu	Gly	1080		
Ala	Gly	Ser	Asp	Val	Phe	Asp	Gly	Asp	Leu	Gly	Met	Gly	Ala	Ala	Lys	Gly	Leu	Gln	Ser	1100		
1110																			1120			
Leu	Pro	Thr	His	Asp	Pro	Ser	Pro	Leu	Gln	Arg	Tyr	Ser	Glu	Asp	Pro	Thr	Val	Pro	Leu	1120		
Pro	Ser	Glu	Thr	Asp	Gly	Tyr	Val	Ala	Pro	Leu	Thr	Cys	Ser	Pro	Gln	Pro	Glu	Tyr	Val	1140		
Asn	Gln	Pro	Asp	Val	Arg	Pro	Gln	Pro	Pro	Ser	Pro	Arg	Glu	Gly	Pro	Leu	Pro	Ala	Ala	1160		
Arg	Pro	Ala	Gly	Ala	Thr	Leu	Glu	Arg	Pro	Lys	Thr	Leu	Ser	Pro	Gly	Lys	Asn	Gly	Val	1180		
Val	Lys	Asp	Val	Phe	Ala	Phe	Gly	Gly	Ala	Val	Glu	Asn	Pro	Glu	Tyr	Leu	Thr	Pro	Gln	1200		
1210																			1220			
Gly	Gly	Ala	Ala	Pro	Gln	Pro	His	Pro	Pro	Pro	Ala	Phe	Ser	Pro	Ala	Phe	Asp	Asn	Leu	1220		
Tyr	Tyr	Trp	Asp	Gln	Asp	Pro	Pro	Glu	Arg	Gly	Ala	Pro	Pro	Ser	Thr	Phe	Lys	Gly	Thr	1240		
Pro	Thr	Ala	Glu	Asn	Pro	Glu	Tyr	Leu	Gly	Leu	Asp	Val	Pro	Val						1257		

610																			620				
Cys	Pro	Ser	Gly	Val	Lys	Pro	Asp	Leu	Ser	Tyr	Met	Pro	Ile	Trp	Lys	Tyr	Pro	Asp	Glu	620			
Glu	Gly	Ile	Cys	Gln	Pro	Cys	Pro	Ile	Asn	Cys	Thr	His	Ser	Cys	Val	Asp	Leu	Asp	Glu	640			
Arg	Gly	Cys	Pro	Ala	Glu	Gln	Arg	Ala	Ser	Pro	Val	Thr	Phe	Ile	Ile	Ala	Thr	Val	Val	660			
Gly	Val	Leu	Leu	Phe	Leu	Ile	Leu	Val	Val	Val	Val	Gly	Ile	Leu	Ile	Lys	Arg	Arg	Arg	680			
Gln	Lys	Ile	Arg	Lys	Tyr	Thr	Met	Arg	Arg	Leu	Leu	Gln	Glu	Thr	Glu	Leu	Val	Glu	Pro	700			
710																			720				
Leu	Thr	Pro	Ser	Gly	Ala	Met	Pro	Asn	Gln	Ala	Gln	Met	Arg	Ile	Leu	Lys	Glu	Thr	Glu	720			
Leu	Arg	Lys	Val	Lys	Val	Leu	Gly	Ser	Gly	Ala	Phe	Gly	Thr	Val	Tyr	Lys	Gly	Ile	Trp	740			
Ile	Pro	Asp	Gly	Glu	Asn	Val	Lys	Ile	Pro	Val	Ala	Ile	Lys	Val	Leu	Arg	Glu	Asn	Thr	760			
Ser	Pro	Lys	Ala	Asn	Lys	Glu	Ile	Leu	Asp	Glu	Ala	Tyr	Val	Met	Ala	Gly	Val	Gly	Ser	780			
Pro	Tyr	Val	Ser	Arg	Leu	Leu	Gly	Ile	Cys	Leu	Thr	Ser	Thr	Val	Gln	Leu	Val	Thr	Gln	800			
810																			820				
Leu	Met	Pro	Tyr	Gly	Cys	Leu	Leu	Asp	His	Val	Arg	Glu	His	Arg	Gly	Arg	Leu	Gly	Ser	820			
Gln	Asp	Leu	Leu	Asn	Trp	Cys	Val	Gln	Ile	Ala	Lys	Gly	Met	Ser	Tyr	Leu	Glu	Asp	Val	840			
Arg	Leu	Val	His	Arg	Asp	Leu	Ala	Ala	Arg	Asn	Val	Leu	Val	Lys	Ser	Pro	Asn	His	Val	860			
Lys	Ile	Thr	Asp	Phe	Gly	Leu	Ala	Arg	Leu	Leu	Asp	Ile	Asp	Glu	Thr	Glu	Tyr	His	Ala	880			
Asp	Gly	Gly	Lys	Val	Pro	Ile	Lys	Trp	Met	Ala	Leu	Glu	Ser	Ile	Leu	Arg	Arg	Arg	Phe	900			
910																			920				
Thr	His	Gln	Ser	Asp	Val	Trp	Ser	Tyr	Gly	Val	Thr	Val	Trp	Glu	Leu	Met	Thr	Phe	Gly	920			
Ala	Lys	Pro	Tyr	Asp	Gly	Ile	Pro	Ala	Arg	Glu	Ile	Pro	Asp	Leu	Leu	Glu	Lys	Gly	Glu	940			
Arg	Leu	Pro	Gln	Pro	Pro	Ile	Cys	Thr	Ile	Asp	Val	Tyr	Met	Ile	Met	Val	Lys	Cys	Trp	960			
Met	Ile	Asp	Ser	Gly	Cys	Arg	Pro	Arg	Phe	Arg	Glu	Leu	Val	Ser	Glu	Phe	Ser	Arg	Met	980			
Ala	Arg	Asp	Pro	Gln	Arg	Phe	Val	Val	Ile	Gln	Asn	Glu	Asp	Leu	Gly	Pro	Ser	Ser	Pro	1000			
1010																			1020				
Met	Asp	Ser	Thr	Phe	Tyr	Arg	Ser	Leu	Leu	Glu	Asp	Asp	Asp	Met	Gly	Asp	Leu	Val	Asp	1020			
Ala	Glu	Glu	Tyr	Leu	Val	Pro	Gln	Gln	Gly	Phe	Phe	Ser	Pro	Asp	Pro	Thr	Pro	Gly	Thr	1040			
Gly	Ser	Thr	Ala	His	Arg	Arg	His	Arg	Ser	Ser	Ser	Ser	Thr	Arg	Ser	Gly	Gly	Gly	Leu	1060			
Thr	Leu	Gly	Leu	Glu	Pro	Ser	Glu	Glu	Gly	Pro	Pro	Arg	Ser	Pro	Leu	Ala	Pro	Ser	Glu	1080			
Gly	Ala	Gly	Ser	Asp	Val	Phe	Asp	Gly	Asp	Leu	Ala	Met	Gly	Val	Thr	Lys	Gly	Leu	Gln	1100			
1110																			1120				
Ser	Leu	Ser	Pro	His	Asp	Leu	Ser	Pro	Leu	Gln	Arg	Tyr	Ser	Glu	Asp	Pro	Thr	Leu	Pro	1120			
Leu	Pro	Pro	Glu	Thr	Asp	Gly	Tyr	Val	Ala	Pro	Leu	Ala	Cys	Ser	Pro	Gln	Pro	Glu	Tyr	1140			
Val	Asn	Gln	Ser	Glu	Val	Gln	Pro	Gln	Pro	Pro	Leu	Thr	Pro	Glu	Gly	Pro	Leu	Pro	Pro	1160			
Val	Arg	Pro	Ala	Gly	Ala	Thr	Leu	Glu	Arg	Pro	Lys	Thr	Leu	Ser	Pro	Gly	Lys	Asn	Gly	1180			
Val	Val	Lys	Asp	Val	Phe	Ala	Phe	Gly	Gly	Ala	Val	Glu	Asn	Pro	Glu	Tyr	Leu	Val	Pro	1200			
1210																			1220				
Arg	Glu	Gly	Thr	Ala	Ser	Pro	Pro	His	Pro	Ser	Pro	Ala	Phe	Ser	Pro	Ala	Phe	Asp	Asn	1220			
Leu	Tyr	Tyr	Trp	Asp	Gln	Asn	Ser	Ser	Glu	Gln	Gly	Pro	Pro	Pro	Ser	Asn	Phe	Glu	Gly	1240			
Thr	Pro	Thr	Ala	Glu	Asn	Pro	Glu	Tyr	Leu	Gly	Leu	Asp	Val	Pro	Val	.	.	.	1258				

																			10																				20
Met	Glu	Leu	Ala	Ala	Trp	Cys	Arg	Trp	Gly	Phe	Leu	Leu	Ala	Leu	Leu	Pro	Pro	Gly	Ile	20																			
Ala	Gly	Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys	Leu	Arg	Leu	Pro	Ala	Ser	Pro	Glu	40																			
Thr	His	Leu	Asp	Met	Leu	Arg	His	Leu	Ser	Gln	Gly	Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	60																			
Glu	Leu	Thr	Tyr	Val	Pro	Ala	Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val	80																			
Gln	Gly	Tyr	Met	Leu	Ile	Ala	His	Asn	Gln	Val	Lys	Arg	Val	Pro	Leu	Gln	Arg	Leu	Arg	100																			
																			110																				120
Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Lys	Tyr	Ala	Leu	Ala	Val	Leu	Asp	Asn	Arg	120																			
Asp	Pro	Gln	Asp	Asn	Val	Ala	Ala	Ser	Thr	Pro	Gly	Arg	Thr	Pro	Glu	Gly	Leu	Arg	Glu	140																			
Leu	Gln	Leu	Arg	Ser	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Arg	Gly	Asn	Pro	160																			
Gln	Leu	Cys	Tyr	Gln	Asp	Met	Val	Leu	Trp	Lys	Asp	Val	Phe	Arg	Lys	Asn	Asn	Gln	Leu	180																			
Ala	Pro	Val	Asp	Ile	Asp	Thr	Asn	Arg	Ser	Arg	Ala	Cys	Pro	Pro	Cys	Ala	Pro	Ala	Cys	200																			
																			210																				220
Lys	Asp	Asn	His	Cys	Trp	Gly	Glu	Ser	Pro	Glu	Asp	Cys	Gln	Ile	Leu	Thr	Gly	Thr	Ile	220																			
Cys	Thr	Ser	Gly	Cys	Ala	Arg	Cys	Lys	Gly	Arg	Leu	Pro	Thr	Asp	Cys	Cys	His	Glu	Gln	240																			
Cys	Ala	Ala	Gly	Cys	Thr	Gly	Pro	Lys	His	Ser	Asp	Cys	Leu	Ala	Cys	Leu	His	Phe	Asn	260																			
His	Ser	Gly	Ile	Cys	Glu	Leu	His	Cys	Pro	Ala	Leu	Val	Thr	Tyr	Asn	Thr	Asp	Thr	Phe	280																			
Glu	Ser	Met	His	Asn	Pro	Glu	Gly	Arg	Tyr	Thr	Phe	Gly	Ala	Ser	Cys	Val	Thr	Thr	Cys	300																			
																			310																				320
Pro	Tyr	Asn	Tyr	Leu	Ser	Thr	Glu	Val	Gly	Ser	Cys	Thr	Leu	Val	Cys	Pro	Pro	Asn	Asn	320																			
Gln	Glu	Val	Thr	Ala	Glu	Asp	Gly	Thr	Gln	Arg	Cys	Glu	Lys	Cys	Ser	Lys	Pro	Cys	Ala	340																			
Arg	Val	Cys	Tyr	Gly	Leu	Gly	Met	Glu	His	Leu	Arg	Gly	Ala	Arg	Ala	Ile	Thr	Ser	Asp	360																			
Asn	Val	Gln	Glu	Phe	Asp	Gly	Cys	Lys	Lys	Ile	Phe	Gly	Ser	Leu	Ala	Phe	Leu	Pro	Glu	380																			
Ser	Phe	Asp	Gly	Asp	Pro	Ser	Ser	Gly	Ile	Ala	Pro	Leu	Arg	Pro	Glu	Gln	Leu	Gln	Val	400																			
																			410																				420
Phe	Glu	Thr	Leu	Glu	Glu	Ile	Thr	Gly	Tyr	Leu	Tyr	Ile	Ser	Ala	Trp	Pro	Asp	Ser	Leu	420																			
Arg	Asp	Leu	Ser	Val	Phe	Gln	Asn	Leu	Arg	Ile	Ile	Arg	Gly	Arg	Ile	Leu	His	Asp	Gly	440																			
Ala	Tyr	Ser	Ser	Thr	Leu	Gln	Gly	Leu	Gly	Ile	His	Ser	Leu	Gly	Leu	Arg	Ser	Leu	Arg	460																			
Glu	Leu	Gly	Ser	Gly	Leu	Ala	Leu	Ile	His	Arg	Asn	Ala	His	Leu	Cys	Phe	Val	His	Thr	480																			
Val	Pro	Trp	Asp	Gln	Leu	Phe	Arg	Asn	Pro	His	Gln	Ala	Leu	Leu	His	Ser	Gly	Asn	Arg	500																			
																			510																				520
Pro	Glu	Glu	Asp	Cys	Gly	Leu	Glu	Gly	Leu	Val	Cys	Asn	Ser	Leu	Cys	Ala	His	Gly	His	520																			
Cys	Trp	Gly	Pro	Gly	Pro	Thr	Gln	Cys	Val	Asn	Cys	Ser	His	Phe	Leu	Arg	Gly	Gln	Glu	540																			
Cys	Val	Glu	Glu	Cys	Arg	Val	Trp	Lys	Gly	Leu	Pro	Arg	Glu	Tyr	Val	Ser	Asp	Lys	Arg	560																			
Cys	Leu	Pro	Cys	His	Pro	Glu	Cys	Gln	Pro	Gln	Asn	Ser	Ser	Glu	Thr	Cys	Phe	Gly	Ser	580																			
Glu	Ala	Asp	Gln	Cys	Ala	Ala	Cys	Ala	His	Tyr	Lys	Asp	Ser	Ser	Ser	Cys	Val	Ala	Arg	600																			

10																			20		
Met	Glu	Leu	Ala	Ala	Leu	Cys	Arg	Trp	Gly	Leu	Leu	Leu	Ala	Leu	Leu	Pro	Pro	Gly	Ala	20	
Ala	Ser	Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys	Leu	Arg	Leu	Pro	Ala	Ser	Pro	Glu	40	
Thr	His	Leu	Asp	Met	Leu	Arg	His	Leu	Tyr	Gln	Gly	Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	60	
Glu	Leu	Thr	Tyr	Leu	Pro	Thr	Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val	80	
Gln	Gly	Tyr	Val	Leu	Ile	Ala	His	Asn	Gln	Val	Arg	Gln	Val	Pro	Leu	Gln	Arg	Leu	Arg	100	
110																			120		
Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Asn	Tyr	Ala	Leu	Ala	Val	Leu	Asp	Asn	Gly	120	
Asp	Pro	Leu	Asn	Asn	Thr	Thr	Pro	Val	Thr	Gly	Ala	Ser	Pro	Gly	Gly	Leu	Arg	Glu	Leu	140	
Gln	Leu	Arg	Ser	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Gln	Arg	Asn	Pro	Gln	160	
Leu	Cys	Tyr	Gln	Asp	Thr	Ile	Leu	Trp	Lys	Asp	Ile	Phe	His	Lys	Asn	Asn	Gln	Leu	Ala	180	
Leu	Thr	Leu	Ile	Asp	Thr	Asn	Arg	Ser	Arg	Ala	Cys	His	Pro	Cys	Ser	Pro	Met	Cys	Lys	200	
210																			220		
Gly	Ser	Arg	Cys	Trp	Gly	Glu	Ser	Ser	Glu	Asp	Cys	Gln	Ser	Leu	Thr	Arg	Thr	Val	Cys	220	
Ala	Gly	Gly	Cys	Ala	Arg	Cys	Lys	Gly	Pro	Leu	Pro	Thr	Asp	Cys	Cys	His	Glu	Gln	Cys	240	
Ala	Ala	Gly	Cys	Thr	Gly	Pro	Lys	His	Ser	Asp	Cys	Leu	Ala	Cys	Leu	His	Phe	Asn	His	260	
Ser	Gly	Ile	Cys	Glu	Leu	His	Cys	Pro	Ala	Leu	Val	Thr	Tyr	Asn	Thr	Asp	Thr	Phe	Glu	280	
Ser	Met	Pro	Asn	Pro	Glu	Gly	Arg	Tyr	Thr	Phe	Gly	Ala	Ser	Cys	Val	Thr	Ala	Cys	Pro	300	
310																			320		
Tyr	Asn	Tyr	Leu	Ser	Thr	Asp	Val	Gly	Ser	Cys	Thr	Leu	Val	Cys	Pro	Leu	His	Asn	Gln	320	
Glu	Val	Thr	Ala	Glu	Asp	Gly	Thr	Gln	Arg	Cys	Glu	Lys	Cys	Ser	Lys	Pro	Cys	Ala	Arg	340	
Val	Cys	Tyr	Gly	Leu	Gly	Met	Gln	His	Leu	Arg	Glu	Val	Arg	Ala	Val	Thr	Ser	Ala	Asn	360	
Ile	Gln	Glu	Phe	Ala	Gly	Cys	Lys	Lys	Ile	Phe	Gly	Ser	Leu	Ala	Phe	Leu	Pro	Glu	Ser	380	
Phe	Asp	Gly	Asp	Pro	Ala	Ser	Asn	Thr	Ala	Pro	Leu	Gln	Pro	Glu	Gln	Leu	Gln	Val	Phe	400	
410																			420		
Glu	Thr	Leu	Glu	Glu	Ile	Thr	Gly	Tyr	Leu	Tyr	Ile	Ser	Ala	Trp	Pro	Asp	Ser	Leu	Pro	420	
Asp	Leu	Ser	Val	Phe	Gln	Asn	Leu	Gln	Val	Ile	Arg	Gly	Arg	Ile	Leu	His	Asn	Gly	Ala	440	
Tyr	Ser	Leu	Thr	Leu	Gln	Gly	Leu	Gly	Ile	Ser	Trp	Leu	Gly	Leu	Arg	Ser	Leu	Arg	Glu	460	
Leu	Gly	Ser	Gly	Leu	Ala	Leu	Ile	His	His	Asn	Thr	His	Leu	Cys	Phe	Val	His	Thr	Val	480	
Pro	Trp	Asp	Gln	Leu	Phe	Arg	Asn	Pro	His	Gln	Ala	Leu	Leu	His	Thr	Ala	Asn	Arg	Pro	500	
510																			520		
Glu	Asp	Glu	Cys	Val	Gly	Glu	Gly	Leu	Ala	Cys	His	Gln	Leu	Cys	Ala	Arg	Gly	His	Cys	520	
Trp	Gly	Pro	Gly	Pro	Thr	Gln	Cys	Val	Asn	Cys	Ser	Gln	Phe	Leu	Arg	Gly	Gln	Glu	Cys	540	
Val	Glu	Glu	Cys	Arg	Val	Leu	Gln	Gly	Leu	Pro	Arg	Glu	Tyr	Val	Asn	Ala	Arg	His	Cys	560	
Leu	Pro	Cys	His	Pro	Glu	Cys	Gln	Pro	Gln	Asn	Gly	Ser	Val	Thr	Cys	Phe	Gly	Pro	Glu	580	
Ala	Asp	Gln	Cys	Val	Ala	Cys	Ala	His	Tyr	Lys	Asp	Pro	Pro	Phe	Cys	Val	Ala	Arg	Cys	600	
610																			620		
Pro	Ser	Gly	Val	Lys	Pro	Asp	Leu	Ser	Tyr	Met	Pro	Ile	Trp	Lys	Phe	Pro	Asp	Glu	Glu	620	
Gly	Ala	Cys	Gln	Pro	Cys	Pro	Ile	Asn	Cys	Thr	His	Ser	Cys	Val	Asp	Leu	Asp	Asp	Lys	640	
Gly	Cys	Pro	Ala	Glu	Gln	Arg	Ala	Ser	Pro	Leu	Thr	Ser	653								

[illegible][illegible]

Fig. 12 (SEQ ID NO: 6)

10										20										
Met	Glu	Leu	Ala	Ala	Leu	Cys	Arg	Trp	Gly	Leu	Leu	Leu	Ala	Leu	Pro	Pro	Gly	Ala	20	
Ala	Ser	Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys	Leu	Arg	Leu	Pro	Ala	Ser	Pro	Glu	40
Thr	His	Leu	Asp	Met	Leu	Arg	His	Leu	Tyr	Gln	Gly	Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	60
Glu	Leu	Thr	Tyr	Leu	Pro	Thr	Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val	80
Gln	Gly	Tyr	Val	Leu	Ile	Ala	His	Asn	Gln	Val	Arg	Gln	Val	Pro	Leu	Gln	Arg	Leu	Arg	100
110										120										
Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Asn	Tyr	Ala	Leu	Ala	Val	Leu	Asp	Asn	Gly	120
Asp	Pro	Leu	Asn	Asn	Thr	Thr	Pro	Val	Thr	Gly	Ala	Ser	Pro	Gly	Gly	Leu	Arg	Glu	Leu	140
Gln	Leu	Arg	Ser	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Gln	Arg	Asn	Pro	Gln	160
Leu	Cys	Tyr	Gln	Asp	Thr	Ile	Leu	Trp	Lys	Asp	Ile	Phe	His	Lys	Asn	Asn	Gln	Leu	Ala	180
Leu	Thr	Leu	Ile	Asp	Thr	Asn	Arg	Ser	Arg	Ala	Cys	His	Pro	Cys	Ser	Pro	Met	Cys	Lys	200
210										220										
Gly	Ser	Arg	Cys	Trp	Gly	Glu	Ser	Ser	Glu	Asp	Cys	Gln	Ser	Leu	Thr	Arg	Thr	Val	Cys	220
Ala	Gly	Gly	Cys	Ala	Arg	Cys	Lys	Gly	Pro	Leu	Pro	Thr	Asp	Cys	Cys	His	Glu	Gln	Cys	240
Ala	Ala	Gly	Cys	Thr	Gly	Pro	Lys	His	Ser	Asp	Cys	Leu	Ala	Cys	Leu	His	Phe	Asn	His	260
Ser	Gly	Ile	Cys	Glu	Leu	His	Cys	Pro	Ala	Leu	Val	Thr	Tyr	Asn	Thr	Asp	Thr	Phe	Glu	280
Ser	Met	Pro	Asn	Pro	Glu	Gly	Arg	Tyr	Thr	Phe	Gly	Ala	Ser	Cys	Val	Thr	Ala	Cys	Pro	300
310										320										
Tyr	Asn	Tyr	Leu	Ser	Thr	Asp	Val	Gly	Ser	Cys	Thr	Leu	Val	Cys	Pro	Leu	His	Asn	Gln	320
Glu	Val	Thr	Ala	Glu	Asp	Gly	Thr	Gln	Arg	Cys	Glu	Lys	Cys	Ser	Lys	Pro	Cys	Ala	Arg	340
Val	Cys	Tyr	Gly	Leu	Gly	Met	Glu	His	Leu	Arg	Glu	Val	Arg	Ala	Val	Thr	Ser	Ala	Asn	360
Ile	Gln	Glu	Phe	Ala	Gly	Cys	Lys	Lys	Ile	Phe	Gly	Ser	Leu	Ala	Phe	Leu	Pro	Glu	Ser	380
Phe	Asp	Gly	Asp	Pro	Ala	Ser	Asn	Thr	Ala	Pro	Leu	Gln	Pro	Glu	Gln	Leu	Gln	Val	Phe	400
410										420										
Glu	Thr	Leu	Glu	Glu	Ile	Thr	Gly	Tyr	Leu	Tyr	Ile	Ser	Ala	Trp	Pro	Asp	Ser	Leu	Pro	420
Asp	Leu	Ser	Val	Phe	Gln	Asn	Leu	Gln	Val	Ile	Arg	Gly	Arg	Ile	Leu	His	Asn	Gly	Ala	440
Tyr	Ser	Leu	Thr	Leu	Gln	Gly	Leu	Gly	Ile	Ser	Trp	Leu	Gly	Leu	Arg	Ser	Leu	Arg	Glu	460
Leu	Gly	Ser	Gly	Leu	Ala	Leu	Ile	His	His	Asn	Thr	His	Leu	Cys	Phe	Val	His	Thr	Val	480
Pro	Trp	Asp	Gln	Leu	Phe	Arg	Asn	Pro	His	Gln	Ala	Leu	Leu	His	Thr	Ala	Asn	Arg	Pro	500
510										520										
Glu	Asp	Glu	Cys	Val	Gly	Glu	Gly	Leu	Ala	Cys	His	Gln	Leu	Cys	Ala	Arg	Gly	His	Cys	520
Trp	Gly	Pro	Gly	Pro	Thr	Gln	Cys	Val	Asn	Cys	Ser	Gln	Phe	Leu	Arg	Gly	Gln	Glu	Cys	540
Val	Glu	Glu	Cys	Arg	Val	Leu	Gln	Gly	Leu	Pro	Arg	Glu	Tyr	Val	Asn	Ala	Arg	His	Cys	560
Leu	Pro	Cys	His	Pro	Glu	Cys	Gln	Pro	Gln	Asn	Gly	Ser	Val	Thr	Cys	Phe	Gly	Pro	Glu	580
Ala	Asp	Gln	Cys	Val	Ala	Cys	Ala	His	Tyr	Lys	Asp	Pro	Pro	Phe	Cys	Val	Ala	Arg	Cys	600

Fig. 12 (SEQ ID NO: 6)

610															620					
Pro	Ser	Gly	Val	Lys	Pro	Asp	Leu	Ser	Tyr	Met	Pro	Ile	Trp	Lys	Phe	Pro	Asp	Glu	Glu	620
Gly	Ala	Cys	Gln	Pro	Cys	Pro	Ile	Asn	Cys	Thr	His	Ser	Cys	Val	Asp	Leu	Asp	Asp	Lys	640
Gly	Cys	Pro	Ala	Glu	Gln	Arg	Ala	Ser	Pro	Leu	Thr	Ser	Gln	Asn	Glu	Asp	Leu	Gly	Pro	660
Ala	Ser	Pro	Leu	Asp	Ser	Thr	Phe	Tyr	Arg	Ser	Leu	Leu	Glu	Asp	Asp	Asp	Met	Gly	Asp	680
Leu	Val	Asp	Ala	Glu	Glu	Tyr	Leu	Val	Pro	Gln	Gln	Gly	Phe	Phe	Cys	Pro	Asp	Pro	Ala	700
710															720					
Pro	Gly	Ala	Gly	Gly	Met	Val	His	His	Arg	His	Arg	Ser	Ser	Ser	Thr	Arg	Ser	Gly	Gly	720
Gly	Asp	Leu	Thr	Leu	Gly	Leu	Glu	Pro	Ser	Glu	Glu	Glu	Ala	Pro	Arg	Ser	Pro	Leu	Ala	740
Pro	Ser	Glu	Gly	Ala	Gly	Ser	Asp	Val	Phe	Asp	Gly	Asp	Leu	Gly	Met	Gly	Ala	Ala	Lys	760
Gly	Leu	Gln	Ser	Leu	Pro	Thr	His	Asp	Pro	Ser	Pro	Leu	Gln	Arg	Tyr	Ser	Glu	Asp	Pro	780
Thr	Val	Pro	Leu	Pro	Ser	Glu	Thr	Asp	Gly	Tyr	Val	Ala	Pro	Leu	Thr	Cys	Ser	Pro	Gln	800
810															820					
Pro	Glu	Tyr	Val	Asn	Gln	Pro	Asp	Val	Arg	Pro	Gln	Pro	Pro	Ser	Pro	Arg	Glu	Gly	Pro	820
Leu	Pro	Ala	Ala	Arg	Pro	Ala	Gly	Ala	Thr	Leu	Glu	Arg	Pro	Lys	Thr	Leu	Ser	Pro	Gly	840
Lys	Asn	Gly	Val	Val	Lys	Asp	Val	Phe	Ala	Phe	Gly	Gly	Ala	Val	Glu	Asn	Pro	Glu	Tyr	860
Leu	Thr	Pro	Gln	Gly	Gly	Ala	Ala	Pro	Gln	Pro	His	Pro	Pro	Pro	Ala	Phe	Ser	Pro	Ala	880
Phe	Asp	Asn	Leu	Tyr	Tyr	Trp	Asp	Gln	Asp	Pro	Pro	Glu	Arg	Gly	Ala	Pro	Pro	Ser	Thr	900
910															920					
Phe	Lys	Gly	Thr	Pro	Thr	Ala	Glu	Asn	Pro	Glu	Tyr	Leu	Gly	Leu	Asp	Val	Pro	Val	•	920

	10		20	
Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu Pro Pro Gly Ala			20	
Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys Leu Arg Leu Pro Ala Ser Pro Glu			40	
Thr His Leu Asp Met Leu Arg His Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu			60	
Glu Leu Thr Tyr Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val			80	
Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu Gln Arg Leu Arg			100	
	110		120	
Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr Ala Leu Ala Val Leu Asp Asn Gly			120	
Asp Pro Leu Asn Asn Thr Thr Pro Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu			140	
Gln Leu Arg Ser Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln			160	
Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn Asn Gln Leu Ala			180	
Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys His Pro Cys Ser Pro Met Cys Lys			200	
	210		220	
Gly Ser Arg Cys Trp Gly Glu Ser Ser Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys			220	
Ala Gly Gly Cys Ala Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys			240	
Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu His Phe Asn His			260	
Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val Thr Tyr Asn Thr Asp Thr Phe Glu			280	
Ser Met Pro Asn Pro Glu Gly Arg Tyr Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro			300	
	310		320	
Tyr Asn Tyr Leu Ser Thr Asp Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln			320	
Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys Pro Cys Ala Arg			340	
Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu Val Arg Ala Val Thr Ser Ala Asn			360	
Ile Gln Glu Phe Ala Gly Cys Lys Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser			380	
Phe Asp Gly Asp Pro Ala Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Gln Val Phe			400	
	410		420	
Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pro Asp Ser Leu Pro			420	
Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile Arg Gly Arg Ile Leu His Asn Gly Ala			440	
Tyr Ser Leu Thr Leu Gln Gly Leu Gly Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu			460	
Leu Gly Ser Gly Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val			480	
Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr Ala Asn Arg Pro			500	
	510		520	
Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His Gln Leu Cys Ala Arg Gly His Cys			520	
Trp Gly Pro Gly Pro Thr Gln Cys Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys			540	
Val Glu Glu Cys Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys			560	
Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys Phe Gly Pro Glu			580	
Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp Pro Pro Phe Cys Val Ala Arg Cys			600	
	610		620	
Pro Ser Gly Val Lys Pro Asp Leu Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu			620	
Gly Ala Cys Gln Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys			640	
Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Gln Asn Glu Asp Leu Gly Pro			660	
Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu Leu Glu Asp Asp Asp Met Gly Asp			680	
Leu Val Asp Ala Glu Glu Tyr Leu Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala			700	
	710		720	
Pro Gly Ala Gly Gly Met Val His His Arg His Arg . . .				714

Fig. 14 (SEQ ID NO: 8)

																			10																				20
Met	Glu	Leu	Ala	Ala	Trp	Cys	Arg	Trp	Gly	Phe	Leu	Leu	Ala	Leu	Leu	Pro	Pro	Gly	Ile	20																			
Ala	Gly	Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys	Leu	Arg	Leu	Pro	Ala	Ser	Pro	Glu	40																			
Thr	His	Leu	Asp	Met	Leu	Arg	His	Leu	Tyr	Gln	Gly	Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	60																			
Glu	Leu	Thr	Tyr	Val	Pro	Ala	Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val	80																			
Gln	Gly	Tyr	Met	Leu	Ile	Ala	His	Asn	Gln	Val	Lys	Arg	Val	Pro	Leu	Gln	Arg	Leu	Arg	100																			
																			110																				120
Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Lys	Tyr	Ala	Leu	Ala	Val	Leu	Asp	Asn	Arg	120																			
Asp	Pro	Gln	Asp	Asn	Val	Ala	Ala	Ser	Thr	Pro	Gly	Arg	Thr	Pro	Glu	Gly	Leu	Arg	Glu	140																			
Leu	Gln	Leu	Arg	Ser	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Arg	Gly	Asn	Pro	160																			
Gln	Leu	Cys	Tyr	Gln	Asp	Met	Val	Leu	Trp	Lys	Asp	Val	Phe	Arg	Lys	Asn	Asn	Gln	Leu	180																			
Ala	Pro	Val	Asp	Ile	Asp	Thr	Asn	Arg	Ser	Arg	Ala	Cys	Pro	Pro	Cys	Ala	Pro	Ala	Cys	200																			
																			210																				220
Lys	Asp	Asn	His	Cys	Trp	Gly	Glu	Ser	Pro	Glu	Asp	Cys	Gln	Ile	Leu	Thr	Gly	Thr	Ile	220																			
Cys	Thr	Ser	Gly	Cys	Ala	Arg	Cys	Lys	Gly	Arg	Leu	Pro	Thr	Asp	Cys	Cys	His	Glu	Gln	240																			
Cys	Ala	Ala	Gly	Cys	Thr	Gly	Pro	Lys	His	Ser	Asp	Cys	Leu	Ala	Cys	Leu	His	Phe	Asn	260																			
His	Ser	Gly	Ile	Cys	Glu	Leu	His	Cys	Pro	Ala	Leu	Val	Thr	Tyr	Asn	Thr	Asp	Thr	Phe	280																			
Glu	Ser	Met	His	Asn	Pro	Glu	Gly	Arg	Tyr	Thr	Phe	Gly	Ala	Ser	Cys	Val	Thr	Thr	Cys	300																			
																			310																				320
Pro	Tyr	Asn	Tyr	Leu	Ser	Thr	Glu	Val	Gly	Ser	Cys	Thr	Leu	Val	Cys	Pro	Pro	Asn	Asn	320																			
Gln	Glu	Val	Thr	Ala	Glu	Asp	Gly	Thr	Gln	Arg	Cys	Glu	Lys	Cys	Ser	Lys	Pro	Cys	Ala	340																			
Arg	Val	Cys	Tyr	Gly	Leu	Gly	Met	Glu	His	Leu	Arg	Gly	Ala	Arg	Ala	Ile	Thr	Ser	Asp	360																			
Asn	Val	Gln	Glu	Phe	Asp	Gly	Cys	Lys	Lys	Ile	Phe	Gly	Ser	Leu	Ala	Phe	Leu	Pro	Glu	380																			
Ser	Phe	Asp	Gly	Asp	Pro	Ser	Ser	Gly	Ile	Ala	Pro	Leu	Arg	Pro	Glu	Gln	Leu	Gln	Val	400																			
																			410																				420
Phe	Glu	Thr	Leu	Glu	Glu	Ile	Thr	Gly	Tyr	Leu	Tyr	Ile	Ser	Ala	Trp	Pro	Asp	Ser	Leu	420																			
Arg	Asp	Leu	Ser	Val	Phe	Gln	Asn	Leu	Arg	Ile	Ile	Arg	Gly	Arg	Ile	Leu	His	Asp	Gly	440																			
Ala	Tyr	Ser	Leu	Thr	Leu	Gln	Gly	Leu	Gly	Ile	His	Ser	Leu	Gly	Leu	Arg	Ser	Leu	Arg	460																			
Glu	Leu	Gly	Ser	Gly	Leu	Ala	Leu	Ile	His	Arg	Asn	Ala	His	Leu	Cys	Phe	Val	His	Thr	480																			
Val	Pro	Trp	Asp	Gln	Leu	Phe	Arg	Asn	Pro	His	Gln	Ala	Leu	Leu	His	Ser	Gly	Asn	Arg	500																			
																			510																				520
Pro	Glu	Glu	Asp	Cys	Gly	Leu	Glu	Gly	Leu	Val	Cys	Asn	Ser	Leu	Cys	Ala	His	Gly	His	520																			
Cys	Trp	Gly	Pro	Gly	Pro	Thr	Gln	Cys	Val	Asn	Cys	Ser	His	Phe	Leu	Arg	Gly	Gln	Glu	540																			
Cys	Val	Glu	Glu	Cys	Arg	Val	Trp	Lys	Gly	Leu	Pro	Arg	Glu	Tyr	Val	Ser	Asp	Lys	Arg	560																			
Cys	Leu	Pro	Cys	His	Pro	Glu	Cys	Gln	Pro	Gln	Asn	Ser	Ser	Glu	Thr	Cys	Phe	Gly	Ser	580																			
Glu	Ala	Asp	Gln	Cys	Ala	Ala	Cys	Ala	His	Tyr	Lys	Asp	Ser	Ser	Ser	Cys	Val	Ala	Arg	600																			
																			610																				620
Cys	Pro	Ser	Gly	Val	Lys	Pro	Asp	Leu	Ser	Tyr	Met	Pro	Ile	Trp	Lys	Tyr	Pro	Asp	Glu	620																			
Glu	Gly	Ile	Cys	Gln	Pro	Cys	Pro	Ile	Asn	Cys	Thr	His	Ser	Cys	Val	Asp	Leu	Asp	Glu	640																			
Arg	Gly	Cys	Pro	Ala	Glu	Gln	Arg	Ala	Ser	Pro	Val	Thr	Phe	654																									

Fig. 15 (SEQ ID NO: 9)

ATG GAG CTG GCG GCC TTG TGC CGC TGG GGG CTC CTC CTC GCC CTC TTG	48
Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu	
1 5 10 15	
CCC CCC GGA GCC GCG AGC ACC CAA GTG TGC ACC GGC ACA GAC ATG AAG	96
Pro. Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys	
20 25 30	
CTG CGG CTC CCT GCC AGT CCC GAG ACC CAC CTG GAC ATG CTC CGC CAC	144
Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His	
35 40 45	
CTC TAC CAG GGC TGC CAG GTG GTG CAG GGA AAC CTG GAA CTC ACC TAC	192
Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr	
50 55 60	
CTG CCC ACC AAT GCC AGC CTG TCC TTC CTG CAG GAT ATC CAG GAG GTG	240
Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val	
65 70 75 80	
CAG GGC TAC GTG CTC ATC GCT CAC AAC CAA GTG AGG CAG GTC CCA CTG	288
Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu	
85 90 95	
CAG AGG CTG CGG ATT GTG CGA GGC ACC CAG CTC TTT GAG GAC AAC TAT	336
Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr	
100 105 110	
GCC CTG GCC GTG CTA GAC AAT GGA GAC CCG CTG AAC AAT ACC ACC CCT	384
Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro	
115 120 125	

17/42
Fig. 15 (SEQ ID NO: 9)

GTC ACA GGG GCC TCC CCA GGA GGC CTG CGG GAG CTG CAG CTT CGA AGC Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser 130 135 140	432
CTC ACA GAG ATC TTG AAA GGA GGG GTC TTG ATC CAG CGG AAC CCC CAG Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln 145 150 155 160	480
CTC TGC TAC CAG GAC ACG ATT TTG TGG AAG GAC ATC TTC CAC AAG AAC Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn 165 170 175	528
AAC CAG CTG GCT CTC ACA CTG ATA GAC ACC AAC CGC TCT CGG GCC TGC Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys 180 185 190	576
CAC CCC TGT TCT CCG ATG TGT AAG GGC TCC CGC TGC TGG GGA GAG AGT His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser 195 200 205	624
TCT GAG GAT TGT CAG AGC CTG ACG CGC ACT GTC TGT GCC GGT GGC TGT Ser Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys Ala Gly Gly Cys 210 215 220	672
GCC CGC TGC AAG GGG CCA CTG CCC ACT GAC TGC TGC CAT GAG CAG TGT Ala Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys 225 230 235 240	720
GCT GCC GGC TGC ACG GGC CCC AAG CAC TCT GAC TGC CTG GCC TGC CTC Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu 245 250 255	768
CAC TTC AAC CAC AGT GGC ATC TGT GAG CTG CAC TGC CCA GCC CTG GTC His Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val 260 265 270	816
ACC TAC AAC ACA GAC ACG TTT GAG TCC ATG CCC AAT CCC GAG GGC CGG Thr Tyr Asn Thr Asp Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg 275 280 285	864
TAT ACA TTC GGC GCC AGC TGT GTG ACT GCC TGT CCC TAC AAC TAC CTT Tyr Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu 290 295 300	912

Fig. 15 (SEQ ID NO: 9)

CCC TGG GAC CAG CTC TTT CGG AAC CCG CAC CAA GCT CTG CTC CAC ACT	1488
Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr	
485 490 495	
GCC AAC CGG CCA GAG GAC GAG TGT GTG GGC GAG GGC CTG GCC TGC CAC	1536
Ala Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His	
500 505 510	
CAG CTG TGC GCC CGA GGG CAC TGC TGG GGT CCA GGG CCC ACC CAG TGT	1584
Gln Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln Cys	
515 520 525	
GTC AAC TGC AGC CAG TTC CTT CGG GGC CAG GAG TGC GTG GAG GAA TGC	1632
Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys	
530 535 540	
CGA GTA CTG CAG GGG CTC CCC AGG GAG TAT GTG AAT GCC AGG CAC TGT	1680
Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys	
545 550 555 560	
TTG CCG TGC CAC CCT GAG TGT CAG CCC CAG AAT GGC TCA GTG ACC TGT	1728
Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys	
565 570 575	
TTT GGA CCG GAG GCT GAC CAG TGT GTG GCC TGT GCC CAC TAT AAG GAC	1776
Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp	
580 585 590	
CCT CCC TTC TGC GTG GCC CGC TGC CCC AGC GGT GTG AAA CCT GAC CTC	1824
Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu	
595 600 605	
TCC TAC ATG CCC ATC TGG AAG TTT CCA GAT GAG GAG GGC GCA TGC CAG	1872
Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln	
610 615 620	
CCT TGC CCC ATC AAC TGC ACC CAC TCC TGT GTG GAC CTG GAT GAC AAG	1920
Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys	
625 630 635 640	
GGC TGC CCC GCC GAG CAG AGA GCC AGC CCT CTG ACG TCC ATC ATC TCT	1968
Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Ile Ser	
645 650 655	
GCG GTG GTT GGC ATT CTG CTG GTC GTG GTC TTG GGG GTG GTC TTT GGG	2016
Ala Val Val Gly Ile Leu Leu Val Val Val Leu Gly Val Val Phe Gly	
660 665 670	

Fig. 15 (SEQ ID NO: 9)

ATC CTC ATC AAG CGA CGG CAG CAG AAG ATC CGG AAG TAC ACG ATG CGG Ile Leu Ile Lys Arg Arg Gln Gln Lys Ile Arg Lys Tyr Thr Met Arg 675 680 685	2064
AGA CTG CTG CAG GAA ACG GAG CTG GTG GAG CCG CTG ACA CCT AGC GGA Arg Leu Leu Gln Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly 690 695 700	2112
GCG ATG CCC AAC CAG GCG CAG ATG CGG ATC CTG AAA GAG ACG GAG CTG Ala Met Pro Asn Gln Ala Gln Met Arg Ile Leu Lys Glu Thr Glu Leu 705 710 715 720	2160
AGG AAG GTG AAG GTG CTT GGA TCT GGC GCT TTT GGC ACA GTC TAC AAG Arg Lys Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys 725 730 735	2208
GGC ATC TGG ATC CCT GAT GGG GAG AAT GTG AAA ATT CCA GTG GCC ATC Gly Ile Trp Ile Pro Asp Gly Glu Asn Val Lys Ile Pro Val Ala Ile 740 745 750	2256
AAA GTG TTG AGG GAA AAC ACA TCC CCC AAA GCC AAC AAA GAA ATC TTA Lys Val Leu Arg Glu Asn Thr Ser Pro Lys Ala Asn Lys Glu Ile Leu 755 760 765	2304
GAC GAA GCA TAC GTG ATG GCT GGT GTG GGC TCC CCA TAT GTC TCC CGC Asp Glu Ala Tyr Val Met Ala Gly Val Gly Ser Pro Tyr Val Ser Arg 770 775 780	2352
CTT CTG GGC ATC TGC CTG ACA TCC ACG GTG CAG CTG GTG ACA CAG CTT Leu Leu Gly Ile Cys Leu Thr Ser Thr Val Gln Leu Val Thr Gln Leu 785 790 795 800	2400
ATG CCC TAT GGC TGC CTC TTA GAC CAT GTC CGG GAA AAC CGC GGA CGC Met Pro Tyr Gly Cys Leu Leu Asp His Val Arg Glu Asn Arg Gly Arg 805 810 815	2448
CTG GGC TCC CAG GAC CTG CTG AAC TGG TGT ATG CAG ATT GCC AAG GGG Leu Gly Ser Gln Asp Leu Leu Asn Trp Cys Met Gln Ile Ala Lys Gly 820 825 830	2496
ATG AGC TAC CTG GAG GAT GTG CGG CTC GTA CAC AGG GAC TTG GCC GCT Met Ser Tyr Leu Glu Asp Val Arg Leu Val His Arg Asp Leu Ala Ala 835 840 845	2544

Fig. 15 (SEQ ID NO: 9)

CGG AAC GTG CTG GTC AAG AGT CCC AAC CAT GTC AAA ATT ACA GAC TTC	2592
Arg Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe	
850 855 860	
GGG CTG GCT CGG CTG CTG GAC ATT GAC GAG ACA GAG TAC CAT GCA GAT	2640
Gly Leu Ala Arg Leu Leu Asp Ile Asp Glu Thr Glu Tyr His Ala Asp	
865 870 875 880	
GGG GGC AAG GTG CCC ATC AAG TGG ATG GCG CTG GAG TCC ATT CTC CGC	2688
Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu Arg	
885 890 895	
CGG CGG TTC ACC CAC CAG AGT GAT GTG TGG AGT TAT GGT GTG ACT GTG	2736
Arg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val	
900 905 910	
TGG GAG CTG ATG ACT TTT GGG GCC AAA CCT TAC GAT GGG ATC CCA GCC	2784
Trp Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro Ala	
915 920 925	
CGG GAG ATC CCT GAC CTG CTG GAA AAG GGG GAG CGG CTG CCC CAG CCC	2832
Arg Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro	
930 935 940	
CCC ATC TGC ACC ATT GAT GTC TAC ATG ATC ATG GTC AAA TGT TGG ATG	2880
Pro Ile Cys Thr Ile Asp Val Tyr Met Ile Met Val Lys Cys Trp Met	
945 950 955 960	
ATT GAC TCT GAA TGT CGG CCA AGA TTC CGG GAG TTG GTG TCT GAA TTC	2928
Ile Asp Ser Glu Cys Arg Pro Arg Phe Arg Glu Leu Val Ser Glu Phe	
965 970 975	
TCC CGC ATG GCC AGG GAC CCC CAG CGC TTT GTG GTC ATC CAG AAT GAG	2976
Ser Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn Glu	
980 985 990	
GAC TTG GGC CCA GCC AGT CCC TTG GAC AGC ACC TTC TAC CGC TCA CTG	3024
Asp Leu Gly Pro Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu	
995 1000 1005	
CTG GAG GAC GAT GAC ATG GGG GAC CTG GTG GAT GCT GAG GAG TAT CTG	3072
Leu Glu Asp Asp Asp Met Gly Asp Leu Val Asp Ala Glu Glu Tyr Leu	
1010 1015 1020	

Fig. 15 (SEQ ID NO: 9)

GTA CCC CAG CAG GGC TTC TTC TGT CCA GAC CCT GCC CCG GGC GCT GGG Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly Ala Gly 1025 1030 1035 1040	3120
GGC ATG GTC CAC CAC AGG CAC CGC AGC TCA TCT ACC AGG AGT GGC GGT Gly Met Val His His Arg His Arg Ser Ser Ser Thr Arg Ser Gly Gly 1045 1050 1055	3168
GGG GAC CTG ACA CTA GGG CTG GAG CCC TCT GAA GAG GAG GCC CCC AGG Gly Asp Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Ala Pro Arg 1060 1065 1070	3216
TCT CCA CTG GCA CCC TCC GAA GGG GCT GGC TCC GAT GTA TTT GAT GGT Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp Gly 1075 1080 1085	3264
GAC CTG GGA ATG GGG GCA GCC AAG GGG CTG CAA AGC CTC CCC ACA CAT Asp Leu Gly Met Gly Ala Ala Lys Gly Leu Gln Ser Leu Pro Thr His 1090 1095 1100	3312
GAC CCC AGC CCT CTA CAG CGG TAC AGT GAG GAC CCC ACA GTA CCC CTG Asp Pro Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Val Pro Leu 1105 1110 1115 1120	3360
CCC TCT GAG ACT GAT GGC TAC GTT GCC CCC CTG ACC TGC AGC CCC CAG Pro Ser Glu Thr Asp Gly Tyr Val Ala Pro Leu Thr Cys Ser Pro Gln 1125 1130 1135	3408
CCT GAA TAT GTG AAC CAG CCA GAT GTT CGG CCC CAG CCC CCT TCG CCC Pro Glu Tyr Val Asn Gln Pro Asp Val Arg Pro Gln Pro Pro Ser Pro 1140 1145 1150	3456
CGA GAG GGC CCT CTG CCT GCT GCC CGA CCT GCT GGT GCC ACT CTG GAA Arg Glu Gly Pro Leu Pro Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu 1155 1160 1165	3504
AGG CCC AAG ACT CTC TCC CCA GGG AAG AAT GGG GTC GTC AAA GAC GTT Arg Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val 1170 1175 1180	3552
TTT GCC TTT GGG GGT GCC GTG GAG AAC CCC GAG TAC TTG ACA CCC CAG Phe Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Thr Pro Gln 1185 1190 1195 1200	3600

Fig. 15 (SEQ ID NO: 9)

GGA GGA GCT GCC CCT CAG CCC CAC CCT CCT CCT GCC TTC AGC CCA GCC	3648
Gly Gly Ala Ala Pro Gln Pro His Pro Pro Pro Ala Phe Ser Pro Ala	
1205 1210 1215	
TTC GAC AAC CTC TAT TAC TGG GAC CAG GAC CCA CCA GAG CGG GGG GCT	3696
Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala	
1220 1225 1230	
CCA CCC AGC ACC TTC AAA GGG ACA CCT ACG GCA GAG AAC CCA GAG TAC	3744
Pro Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr	
1235 1240 1245	
CTG GGT CTG GAC GTG CCA GTG TGA	3768
Leu Gly Leu Asp Val Pro Val	
1250 1255	

Herceptin Binding by Direct Elisa 10/5/99





Fig. 17

Fig. 18

Comparison of Her2neu ECD-PD Expression in CHO-K1 (S/SF) and Pichia (Non reducing conditions)



Legend : CHO; A, B, C = 2,5µl / 5µl / 10µl

CHO purif; A, B, C = 125ng / 250ng / 500ng

Pichia ; A, B, C = 2,5µl / 5µl / 10µl from a 1/30 dilution of OD 120

Fig. 19 (SEQ ID NO:11)

atggagctgg	eggcctggg	ccgttggggg	ttcctcctcg	ccctcctgtc	ccccggagcc	60
gcgggtaccc	aagtgtgtac	cggtaccgac	atgaagtgtg	gactccctgc	cagtcctgag	120
accacactgg	acatgtctcg	ccacctctac	cagggtgtgc	aggtgggtgca	gggcaatttg	180
gagcttacct	acctgcccgc	caatgccagc	ctctcattcc	tgcaggacat	ccaggaagtc	240
cagggataca	tgctcatcgc	tcacaaccga	gtgaacacg	tcccactgca	gaggttgcg	300
atcgtgagag	ggactcagct	ctttgaggac	aagtatgccc	tggctgtgct	agacaaccga	360
gaccctttgg	acaacgtcac	caccgccgccc	ccaggcagaa	ccccagaagg	gctgccccgag	420
ctgcagcttc	gaagtctcac	agagatcttg	aaggggaggag	ttttgatccg	tgggaaccct	480
cagctctgct	accaggacat	ggttttgtgg	aaggatgtcc	tccgtaagaa	taaccagctg	540
gctcctgtcg	acatggacac	caatcgttcc	cgggcctgtc	caccttgtgc	cccaacctgc	600
aaagacaatc	actgttgggg	tgagagtcc	gaagactgtc	agatcttgac	tggcaccatc	660
tgtactagt	gctgtgccc	gtgcaagggc	cggctgccc	ctgactgttg	ccatgagcag	720
tgtgtgcag	gctgcacggg	tcccagcat	cttgactgcc	tggcctgcct	ccacttcaat	780
catagtggta	tctgtgagct	gcactgccc	gcccctcatca	cctacaacac	agacaccttc	840
gagtccatgc	tcaacctga	gggtcgctac	acctttgggtg	ccagctgtgt	gaccacctgc	900
ccctacaact	acctctccac	ggaagtggga	tcctgcactc	tggctgtgtcc	cccgaacaac	960
caagaggtca	cagctgagga	cggaaacacag	cgggtgtgaga	aatgcagcaa	gcctgtgtgt	1020
ggagtatgct	atgggtctggg	catggagcac	ctccgagggg	cgagggccat	caccagtgtg	1080
aatatccagg	agtttgtgtg	ctgcaagaag	atctttggga	gcctggcatt	tttgccggag	1140
agctttgatg	ggaacccctc	ctccggcggt	gccccactga	agccagagca	tctccaagt	1200
ttcgaaaccc	tggaggagat	cacaggttac	ctatacattt	cagcatggcc	agagagcttc	1260
caagacctca	gtgtcttcca	gaaccttcgg	gtcattcggg	gacggattct	ccatgatggg	1320
gcttactcat	tgacgttgca	aggcctgggg	attcactcac	tggggctacg	ctcactgcgg	1380
gagctgggca	gtggattggc	tctcattcac	cgcaacaccc	atctctgctt	tgtaaacact	1440
gtaccttggg	accagctctt	ccggaacccg	caccaggccc	tactccacag	tgggaaccgg	1500
ccagaagagg	catgtgtgtc	tgagggtctg	gtctgttaact	cactgtgtgc	ccgtgggcac	1560
tgctgggggc	caggggccac	ccagtgtgtc	aactgcagtc	agttcctccg	gggccaggag	1620
tgtgtggagg	agtgccgagt	atggaagggg	ctcccagggg	agtatgtgag	gggcaagcac	1680
tgtctgccat	gccaccccca	gtgtcagcct	caaaacagct	cggagacctg	ctatggatcg	1740
gaggtgacc	agtgtgaggg	ttgtgcccac	tacaaggact	catcttctcg	tgtggctcgc	1800
tgccccagt	gtgtgaagcc	agacctctcc	tacatgccta	tctggaagta	cccggatgag	1860
gagggcatat	gtcagccatg	ccccatcaac	tgcacccact	catgtgtgga	cctggacgaa	1920
cgaggctgcc	cagcagagca	gagagccagc	ccagtgcac	tcatcattgc	aactgtgggtg	1980
ggcgtcctgt	tgttcctgat	catagtgtgtg	gtcattggaa	tcctaataca	acgaaggcca	2040
cagaagatcc	ggaagtatac	catgcgtagg	ctcgtcagg	agaccgagct	gggtggagccg	2100
ctgacgcccc	gtggagctgt	gcccacccag	gctcagatgc	ggatccctaaa	ggagacagag	2160
ctaaggaagc	tgaaggtgct	tgggtcagga	gccttcggca	ctgtctacaa	gggcatctgg	2220
atcccagatg	gggagaacgt	gaaaatcccc	gtggccatca	aggtgttgag	ggaaaacaca	2280
tctcctaaag	ctaacaaga	aatcctagat	gaagcgtacg	tcatggctgg	tgtgggttct	2340
ccatattgtgt	cccgctcct	gggcatctgc	ctgacatcca	cagtgcagct	ggtgacacag	2400
cttatgccct	atggctgcct	tctggaccat	gtccgagaac	accgaggtcg	cttaggctcc	2460
caggacctgc	tcaactgggt	tgttcagatt	gccaagggga	tgagctacct	ggagggaagt	2520
cggcttgttc	acagggacct	agctgcccga	aacgtgctag	tcaagagtcc	caaccacgtc	2580
aagattaccg	acttcgggct	ggcacggctg	ctggacattg	atgagactga	ataccatgca	2640
gatgggggca	aggtgccc	caagtggatg	gcattggaat	ctattctcag	acgccgggtc	2700
actcatcaga	gtgatgtgtg	gagctatggt	gtgactgtgt	gggagctgat	gacctttggg	2760
gccaaacctt	acgatgggat	cccagctcgg	gagatccctg	atttgctgga	gaaggagaa	2820
cgcctacctc	agcctccaat	ctgcaccatc	gacgtctaca	tgatcatggg	caaatgttgg	2880
atgattgact	ccgaatgtcg	cccagagattc	cgggagttgg	tatcagaatt	ctcccgtatg	2940
gcaagggacc	cccagcgctt	tgtgggtcatc	cagaacgagg	acttaggccc	ctccagcccc	3000
atggacagca	ccttctaccg	ttcactgctg	gaggatgatg	acatggggga	gctggctcgat	3060
gctgaagagt	acctgggtacc	ccagcaggga	ttcttctccc	cagaccctgc	cctagggtact	3120
gggagcacag	cccaccgcag	acaccgcagc	tcgtcggcca	ggagtggcgg	tgggtgagctg	3180
acactgggcc	tggagccctc	ggaagaagag	ccccccagat	ctccactggc	tccctccgaa	3240
ggggctggct	ccgatgtgtt	tgatgggtgac	ctggcagtg	gggtaaccaa	aggactgcag	3300
agcctctctc	cacatgacct	cagccctcta	cagcgggtaca	gtgaggatcc	cacattacct	3360

Fig. 19 (SEQ ID NO:11)

ctgccccccg	agactgatgg	ctacgttgct	cccctggcct	gcagccccca	gcccgagtat	3420
gtgaaccagc	cagagggttcg	gcctcagtct	cccttgacct	cagaggggtcc	tccgcctccc	3480
atccgacctg	ctgggtgctac	tctagaaaga	cccaagactc	tctctcctgg	gaaaaatggg	3540
gttgtcaaag	acgttttttc	ctttgggggt	gctgtggaga	accctgaata	cctagcaccc	3600
agagcaggca	ctgcctctca	gccccaccct	tctcctgcct	tcagcccagc	ctttgacaac	3660
ctctattact	gggaccagaa	ctcatcggag	cagggtcctc	caccaagtac	ctttgaaggg	3720
acccccactg	cagagaaccc	tgagtaccta	ggcctggatg	tgccagtatg	a	3771

Fig. 20 (SEQ ID NO:14)

```

Met Glu Leu Ala Ala Trp Cys Arg Trp Gly Phe Leu Leu Ala Leu Leu
 1          5          10          15
Ser Pro Gly Ala Ala Gly Thr Gln Val Cys Thr Gly Thr Asp Met Lys
 20          25          30
Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His
 35          40          45
Leu Tyr Gln Gly Cys Gln Val Gln Gly Asn Leu Glu Leu Thr Tyr
 50          55          60
Leu Pro Ala Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val
 65          70          75          80
Gln Gly Tyr Met Leu Ile Ala His Asn Arg Val Lys His Val Pro Leu
 85          90          95
Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Lys Tyr
100          105          110
Ala Leu Ala Val Leu Asp Asn Arg Asp Pro Leu Asp Asn Val Thr Thr
115          120          125
Ala Ala Pro Gly Arg Thr Pro Glu Gly Leu Arg Glu Leu Gln Leu Arg
130          135          140
Ser Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Arg Gly Asn Pro
145          150          155          160
Gln Leu Cys Tyr Gln Asp Met Val Leu Trp Lys Asp Val Leu Arg Lys
165          170          175
Asn Asn Gln Leu Ala Pro Val Asp Met Asp Thr Asn Arg Ser Arg Ala
180          185          190
Cys Pro Pro Cys Ala Pro Thr Cys Lys Asp Asn His Cys Trp Gly Glu
195          200          205
Ser Pro Glu Asp Cys Gln Ile Leu Thr Gly Thr Ile Cys Thr Ser Gly
210          215          220
Cys Ala Arg Cys Lys Gly Arg Leu Pro Thr Asp Cys Cys His Glu Gln
225          230          235          240
Cys Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys
245          250          255
Leu His Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu
260          265          270
Ile Thr Tyr Asn Thr Asp Thr Phe Glu Ser Met Leu Asn Pro Glu Gly
275          280          285
Arg Tyr Thr Phe Gly Ala Ser Cys Val Thr Thr Cys Pro Tyr Asn Tyr
290          295          300
Leu Ser Thr Glu Val Gly Ser Cys Thr Leu Val Cys Pro Pro Asn Asn
305          310          315          320
Gln Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser
325          330          335
Lys Pro Cys Ala Gly Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg
340          345          350
Gly Ala Arg Ala Ile Thr Ser Asp Asn Ile Gln Glu Phe Ala Gly Cys
355          360          365
Lys Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly
370          375          380
Asn Pro Ser Ser Gly Val Ala Pro Leu Lys Pro Glu His Leu Gln Val
385          390          395          400
Phe Glu Thr Leu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp
405          410          415
Pro Glu Ser Phe Gln Asp Leu Ser Val Phe Gln Asn Leu Arg Val Ile

```

420										425					430					
Arg	Gly	Arg	Ile	Leu	His	Asp	Gly	Ala	Tyr	Ser	Leu	Thr	Leu	Gln	Gly					
435										440					445					
Leu	Gly	Ile	His	Ser	Leu	Gly	Leu	Arg	Ser	Leu	Arg	Glu	Leu	Gly	Ser					
450										455					460					
Gly	Leu	Ala	Leu	Ile	His	Arg	Asn	Thr	His	Leu	Cys	Phe	Val	Asn	Thr					
465	470										475					480				
Val	Pro	Trp	Asp	Gln	Leu	Phe	Arg	Asn	Pro	His	Gln	Ala	Leu	Leu	His					
485										490					495					
Ser	Gly	Asn	Arg	Pro	Glu	Glu	Ala	Cys	Gly	Leu	Glu	Gly	Leu	Val	Cys					
500										505					510					
Asn	Ser	Leu	Cys	Ala	Arg	Gly	His	Cys	Trp	Gly	Pro	Gly	Pro	Thr	Gln					
515										520					525					
Cys	Val	Asn	Cys	Ser	Gln	Phe	Leu	Arg	Gly	Gln	Glu	Cys	Val	Glu	Glu					
530										535					540					
Cys	Arg	Val	Trp	Lys	Gly	Leu	Pro	Arg	Glu	Tyr	Val	Arg	Gly	Lys	His					
545	550										555					560				
Cys	Leu	Pro	Cys	His	Pro	Glu	Cys	Gln	Pro	Gln	Asn	Ser	Ser	Glu	Thr					
565										570					575					
Cys	Tyr	Gly	Ser	Glu	Ala	Asp	Gln	Cys	Glu	Ala	Cys	Ala	His	Tyr	Lys					
580										585					590					
Asp	Ser	Ser	Ser	Cys	Val	Ala	Arg	Cys	Pro	Ser	Gly	Val	Lys	Pro	Asp					
595										600					605					
Leu	Ser	Tyr	Met	Pro	Ile	Trp	Lys	Tyr	Pro	Asp	Glu	Glu	Gly	Ile	Cys					
610										615					620					
Gln	Pro	Cys	Pro	Ile	Asn	Cys	Thr	His	Ser	Cys	Val	Asp	Leu	Asp	Glu					
625	630										635					640				
Arg	Gly	Cys	Pro	Ala	Glu	Gln	Arg	Ala	Ser	Pro	Val	Thr	Phe	Ile	Ile					
645										650					655					
Ala	Thr	Val	Val	Gly	Val	Leu	Leu	Phe	Leu	Ile	Ile	Val	Val	Val	Ile					
660										665					670					
Gly	Ile	Leu	Ile	Lys	Arg	Arg	Arg	Gln	Lys	Ile	Arg	Lys	Tyr	Thr	Met					
675										680					685					
Arg	Arg	Leu	Leu	Gln	Glu	Thr	Glu	Leu	Val	Glu	Pro	Leu	Thr	Pro	Ser					
690										695					700					
Gly	Ala	Val	Pro	Asn	Gln	Ala	Gln	Met	Arg	Ile	Leu	Lys	Glu	Thr	Glu					
705	710										715					720				
Leu	Arg	Lys	Leu	Lys	Val	Leu	Gly	Ser	Gly	Ala	Phe	Gly	Thr	Val	Tyr					
725										730					735					
Lys	Gly	Ile	Trp	Ile	Pro	Asp	Gly	Glu	Asn	Val	Lys	Ile	Pro	Val	Ala					
740										745					750					
Ile	Lys	Val	Leu	Arg	Glu	Asn	Thr	Ser	Pro	Lys	Ala	Asn	Lys	Glu	Ile					
755										760					765					
Leu	Asp	Glu	Ala	Tyr	Val	Met	Ala	Gly	Val	Gly	Ser	Pro	Tyr	Val	Ser					
770										775					780					
Arg	Leu	Leu	Gly	Ile	Cys	Leu	Thr	Ser	Thr	Val	Gln	Leu	Val	Thr	Gln					
785	790										795					800				
Leu	Met	Pro	Tyr	Gly	Cys	Leu	Leu	Asp	His	Val	Arg	Glu	His	Arg	Gly					
805										810					815					
Arg	Leu	Gly	Ser	Gln	Asp	Leu	Leu	Asn	Trp	Cys	Val	Gln	Ile	Ala	Lys					
820										825					830					
Gly	Met	Ser	Tyr	Leu	Glu	Glu	Val	Arg	Leu	Val	His	Arg	Asp	Leu	Ala					
835										840					845					
Ala	Arg	Asn	Val	Leu	Val	Lys	Ser													

Fig. 20 (SEQ ID NO:14)


```

Asp Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu
      885                      890                      895
Arg Arg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr
      900                      905                      910
Val Trp Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro
      915                      920                      925
Ala Arg Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln
      930                      935                      940
Pro Pro Ile Cys Thr Ile Asp Val Tyr Met Ile Met Val Lys Cys Trp
      945                      950                      955                      960
Met Ile Asp Ser Glu Cys Arg Pro Arg Phe Arg Glu Leu Val Ser Glu
      965                      970                      975
Phe Ser Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn
      980                      985                      990
Glu Asp Leu Gly Pro Ser Ser Pro Met Asp Ser Thr Phe Tyr Arg Ser
      995                      1000                      1005
Leu Leu Glu Asp Asp Asp Met Gly Glu Leu Val Asp Ala Glu Glu Tyr
      1010                      1015                      1020
Leu Val Pro Gln Gln Gly Phe Phe Ser Pro Asp Pro Ala Leu Gly Thr
      1025                      1030                      1035                      1040
Gly Ser Thr Ala His Arg Arg His Arg Ser Ser Ser Ala Arg Ser Gly
      1045                      1050                      1055
Gly Gly Glu Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Pro Pro
      1060                      1065                      1070
Arg Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp
      1075                      1080                      1085
Gly Asp Leu Ala Val Gly Val Thr Lys Gly Leu Gln Ser Leu Ser Pro
      1090                      1095                      1100
His Asp Leu Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Leu Pro
      1105                      1110                      1115                      1120
Leu Pro Pro Glu Thr Asp Gly Tyr Val Ala Pro Leu Ala Cys Ser Pro
      1125                      1130                      1135
Gln Pro Glu Tyr Val Asn Gln Pro Glu Val Arg Pro Gln Ser Pro Leu
      1140                      1145                      1150
Thr Pro Glu Gly Pro Pro Pro Pro Ile Arg Pro Ala Gly Ala Thr Leu
      1155                      1160                      1165
Glu Arg Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp
      1170                      1175                      1180
Val Phe Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Ala Pro
      1185                      1190                      1195                      1200
Arg Ala Gly Thr Ala Ser Gln Pro His Pro Ser Pro Ala Phe Ser Pro
      1205                      1210                      1215
Ala Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asn Ser Ser Glu Gln Gly
      1220                      1225                      1230
Pro Pro Pro Ser Thr Phe Glu Gly Thr Pro Thr Ala Glu Asn Pro Glu
      1235                      1240                      1245
Tyr Leu Gly Leu Asp Val Pro Val
      1250                      1255

```

Fig. 21

Expression of mNeu and mECDPD in HEK Cells



Western blot analysis of mNeu and mECDPD recombinants expressed in HEK293 lysates. Delta mNeu has a mutated kinase domain while mECDPD-Tcp0 contains a 6-mer peptide encoding a ribosomal P-peptide epitope. The cNeu3 anti-PD antibody was used for detection of proteins.

Met	Glu	Leu	Ala	Ala	Trp	Cys	Arg	Trp	Gly	Phe	Leu	Leu	Ala	Leu	Leu
				5					10					15	
Ser	Pro	Gly	Ala	Ala	Gly	Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys
			20					25					30		
Leu	Arg	Leu	Pro	Ala	Ser	Pro	Glu	Thr	His	Leu	Asp	Met	Leu	Arg	His
		35					40					45			
Leu	Tyr	Gln	Gly	Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	Glu	Leu	Thr	Tyr
	50					55					60				
Leu	Pro	Ala	Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val
	65				70					75					80
Gln	Gly	Tyr	Met	Leu	Ile	Ala	His	Asn	Arg	Val	Lys	His	Val	Pro	Leu
				85					90					95	
Gln	Arg	Leu	Arg	Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Lys	Tyr
			100					105					110		
Ala	Leu	Ala	Val	Leu	Asp	Asn	Arg	Asp	Pro	Leu	Asp	Asn	Val	Thr	Thr
		115					120					125			
Ala	Ala	Pro	Gly	Arg	Thr	Pro	Glu	Gly	Leu	Arg	Glu	Leu	Gln	Leu	Arg
	130					135					140				
Ser	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Arg	Gly	Asn	Pro
	145				150					155					160
Gln	Leu	Cys	Tyr	Gln	Asp	Met	Val	Leu	Trp	Lys	Asp	Val	Leu	Arg	Lys
				165					170					175	
Asn	Asn	Gln	Leu	Ala	Pro	Val	Asp	Met	Asp	Thr	Asn	Arg	Ser	Arg	Ala
			180					185					190		
Cys	Pro	Pro	Cys	Ala	Pro	Thr	Cys	Lys	Asp	Asn	His	Cys	Trp	Gly	Glu
		195					200					205			
Ser	Pro	Glu	Asp	Cys	Gln	Ile	Leu	Thr	Gly	Thr	Ile	Cys	Thr	Ser	Gly
	210					215					220				
Cys	Ala	Arg	Cys	Lys	Gly	Arg	Leu	Pro	Thr	Asp	Cys	Cys	His	Glu	Gln
	225				230					235					240
Cys	Ala	Ala	Gly	Cys	Thr	Gly	Pro	Lys	His	Ser	Asp	Cys	Leu	Ala	Cys
				245					250					255	
Leu	His	Phe	Asn	His	Ser	Gly	Ile	Cys	Glu	Leu	His	Cys	Pro	Ala	Leu
			260					265					270		
Ile	Thr	Tyr	Asn	Thr	Asp	Thr	Phe	Glu	Ser	Met	Leu	Asn	Pro	Glu	Gly
		275					280					285			
Arg	Tyr	Thr	Phe	Gly	Ala	Ser	Cys	Val	Thr	Thr	Cys	Pro	Tyr	Asn	Tyr
	290					295					300				

Fig. 22 (SEQ ID NO:27)

Leu Ser Thr Glu Val Gly Ser Cys Thr Leu Val Cys Pro Pro Asn Asn
 305 310 315 320
 Gln Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser
 325 330 335
 Lys Pro Cys Ala Gly Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg
 340 345 350
 Gly Ala Arg Ala Ile Thr Ser Asp Asn Ile Gln Glu Phe Ala Gly Cys
 355 360 365
 Lys Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly
 370 375 380
 Asn Pro Ser Ser Gly Val Ala Pro Leu Lys Pro Glu His Leu Gln Val
 385 390 395 400
 Phe Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp
 405 410 415
 Pro Glu Ser Phe Gln Asp Leu Ser Val Phe Gln Asn Leu Arg Val Ile
 420 425 430
 Arg Gly Arg Ile Leu His Asp Gly Ala Tyr Ser Leu Thr Leu Gln Gly
 435 440 445
 Leu Gly Ile His Ser Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser
 450 455 460
 Gly Leu Ala Leu Ile His Arg Asn Thr His Leu Cys Phe Val Asn Thr
 465 470 475 480
 Val Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His
 485 490 495
 Ser Gly Asn Arg Pro Glu Glu Ala Cys Gly Leu Glu Gly Leu Val Cys
 500 505 510
 Asn Ser Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln
 515 520 525
 Cys Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu
 530 535 540
 Cys Arg Val Trp Lys Gly Leu Pro Arg Glu Tyr Val Arg Gly Lys His
 545 550 555 560
 Cys Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Ser Ser Glu Thr
 565 570 575
 Cys Tyr Gly Ser Glu Ala Asp Gln Cys Glu Ala Cys Ala His Tyr Lys
 580 585 590
 Asp Ser Ser Ser Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp
 595 600 605

Fig. 22 (SEQ ID NO:27)

```

Leu Ser Tyr Met Pro Ile Trp Lys Tyr Pro Asp Glu Glu Gly Ile Cys
610                               615           620

Gln Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Glu
625                               630           635           640

Arg Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Val Thr Ser Gln Asn
                               645           650           655

Glu Asp Leu Gly Pro Ser Ser Pro Met Asp Ser Thr Phe Tyr Arg Ser
                               660           665           670

Leu Leu Glu Asp Asp Asp Met Gly Glu Leu Val Asp Ala Glu Glu Tyr
675                               680           685

Leu Val Pro Gln Gln Gly Phe Phe Ser Pro Asp Pro Ala Leu Gly Thr
690                               695           700

Gly Ser Thr Ala His Arg Arg His Arg Ser Ser Ser Ala Arg Ser Gly
705                               710           715           720

Gly Gly Glu Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Pro Pro
725                               730           735

Arg Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp
740                               745           750

Gly Asp Leu Ala Val Gly Val Thr Lys Gly Leu Gln Ser Leu Ser Pro
755                               760           765

His Asp Leu Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Leu Pro
770                               775           780

Leu Pro Pro Glu Thr Asp Gly Tyr Val Ala Pro Leu Ala Cys Ser Pro
785                               790           795           800

Gln Pro Glu Tyr Val Asn Gln Pro Glu Val Arg Pro Gln Ser Pro Leu
805                               810           815

Thr Pro Glu Gly Pro Pro Pro Pro Ile Arg Pro Ala Gly Ala Thr Leu
820                               825           830

Glu Arg Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp
835                               840           845

Val Phe Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Ala Pro
850                               855           860

Arg Ala Gly Thr Ala Ser Gln Pro His Pro Ser Pro Ala Phe Ser Pro
865                               870           875           880

Ala Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asn Ser Ser Glu Gln Gly
885                               890           895

Pro Pro Pro Ser Thr Phe Glu Gly Thr Pro Thr Ala Glu Asn Pro Glu
900                               905           910

Tyr Leu Gly Leu Asp Val Pro Val
915                               920

```

Fig. 23 (SEQ ID NO:28)

```
atggagctgg  eggcctgggtg  ccgttggggg  ttctctctcg  ccctcctgtc  ccccgagacc  60
gcgggtaccc  aagtgtgtac  cggtagccgac  atgaagttgc  gactccctgc  cagtcctgag  120
acccacctgg  acatgcttcg  ccacctctac  cagggtgtgc  aggtgggtgca  gggcaatttg  180
gagcttacct  acctgcccgc  caatgccagc  ctctcattcc  tgcaggacat  ccaggaagtc  240
cagggataca  tgctcatcgc  tcacaaccga  gtgaaacacg  tcccactgca  gaggttgctc  300
atcgtgagag  ggactcagct  ctttgaggac  aagtatgccc  tggctgtgct  agacaaccga  360
gaccctttgg  acaacgtcac  caccgcccgc  ccaggcagaa  ccccgagaag  gctgcgggag  420
ctgcagcttc  gaagtctcac  agagatcttg  aaggaggag  ttttgatccg  tgggaaccct  480
cagctctgct  accaggacat  ggttttgtgg  aaggatgtcc  tccgtaagaa  taaccagctg  540
gctcctgtcg  acatggacac  caatcgttcc  cgggcctgtc  cactctgtgc  cccaacctgc  600
aaagacaatc  actgttgggg  tgagagtcct  gaagactgtc  agatcttgac  tggcaccatc  660
tgtactagt  gctgtgccc  gtgcaagggc  cggctgccc  ctgactgttg  ccctgagcag  720
tgtgtgag  gctgcacggg  tcccaagcat  tctgactgcc  tggcctgcct  ccacttcaat  780
catagtggta  tctgtgagct  gcaactgccc  gccctcatca  cctacaacac  agacaccttc  840
gagtccatgc  tcaacctga  gggctcgtac  acctttgggt  ccagctgtgt  gaccacctgc  900
ccctacaact  acctctccac  ggaagtggga  tctgtcactc  tggctgtgtc  cccgaacaac  960
caagaggtca  cagctgagga  cggaaacacg  cgggtgtgaga  aatgcagcaa  gccctgtgct  1020
ggagtatgct  atgggtctggg  catggagcac  ctccgagggg  cgagggccat  caccagtgc  1080
aatatccagg  agtttgctgg  ctgcaagaag  atctttggga  gcctggcatt  tttgccggag  1140
agctttgatg  ggaacccctc  ctccggcggt  gcccactga  agccagagca  tctccaagt  1200
ttcgaaacc  tggaggagat  cacaggttac  ctatacattt  cagcatggcc  agagagcttc  1260
caagacctca  gtgtcttcca  gaaccttcgg  gtcattcggg  gacggattct  ccatgatggt  1320
gcttactcat  tgacgttgca  aggcctgggg  attcactcac  tggggctacg  ctactgcgg  1380
gagctgggca  gtggattggc  tctcatcac  cgcaacacc  atctctgctt  tgtaaacact  1440
gtaccttggg  accagctctt  ccggaaccgc  caccaggccc  tactccacag  tgggaaccgg  1500
ccagaagagg  catgtggtct  tgagggttg  gtctgtaact  cactgtgtgc  ccgtgggcac  1560
tgtgtggggc  cagggccac  ccagtgtgtc  aactgcagtc  agttcctccg  gggccaggag  1620
tgtgtggagg  agtgccgagt  atggaagggg  ctcccagggg  agtatgtgag  gggcaagcac  1680
tgtctgcat  gccaccccg  gtgtcagcct  caaaacagct  cggagacctg  ctatggatcg  1740
gaggctgacc  agtgtgaggc  ttgtgccac  tacaaggact  catcttctcg  tgtggctcgc  1800
tgccccagt  gtgtgaagcc  agacctctcc  tacatgccta  tctggaagta  cccgatgag  1860
gagggcata  gtcagccatg  ccccatcaac  tgcacccact  catgtgtgga  cctggacgaa  1920
cgaggctgcc  cagcagagca  gagagccagc  ccagtgcgt  ctgagaacga  ggacttaggc  1980
ccctccagcc  ccattctac  cacttctac  cgttcaactg  tggaggatga  tgacatgggg  2040
gagctggctg  atgtgaaga  gtacctggta  cccagcagg  gattcttctc  cccagaccct  2100
gccctaggta  ctgggagcac  agcccaccgc  agacaccgca  gctcgtcggc  caggagtggc  2160
gggtgtgagc  tgacactggg  cctggagccc  tcggaagaag  agccccccag  atctccactg  2220
gtccctccg  aaggggctgg  ctccgatgtg  tttgatgggt  acctggcagt  gggggtaacc  2280
aaaggactgc  agagcctctc  tccacatgac  ctacgcctc  tacagcggt  cagtgaggat  2340
cccacattac  ctctgcccc  cgagactgat  ggctacgttg  ctccctggc  ctgcagcccc  2400
cagcccgagt  atgtgaacca  gccagaggtt  cggcctcagt  ctccctgac  cccagaggg  2460
cctccgcctc  ccacccgacc  tgctgggtgc  actctagaaa  gacccaagac  tctctctcct  2520
gggaaaaatg  ggggtgtcaa  agacgttttt  gcctttgggg  gtgctgtgga  gaacctgaa  2580
tacctagcac  ccagagcagg  cactgcctct  cagccccacc  ctctcctgc  cttcagccca  2640
gcctttgaca  acctctatta  ctgggaccag  aactcatcgg  agcagggtcc  tccaccaagt  2700
acctttgaag  ggacccccac  tgcagagaac  cctgagtacc  taggcctgga  tgtgccagta  2760
tga
```

Met	Glu	Leu	Ala	Ala	Trp	Cys	Arg	Trp	Gly	Phe	Leu	Leu	Ala	Leu	Leu
				5					10					15	
Ser	Pro	Gly	Ala	Ala	Gly	Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys
			20					25					30		
Leu	Arg	Leu	Pro	Ala	Ser	Pro	Glu	Thr	His	Leu	Asp	Met	Leu	Arg	His
		35					40					45			
Leu	Tyr	Gln	Gly	Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	Glu	Leu	Thr	Tyr
	50					55					60				
Leu	Pro	Ala	Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val
65					70					75					80
Gln	Gly	Tyr	Met	Leu	Ile	Ala	His	Asn	Arg	Val	Lys	His	Val	Pro	Leu
				85					90					95	
Gln	Arg	Leu	Arg	Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Lys	Tyr
			100					105					110		
Ala	Leu	Ala	Val	Leu	Asp	Asn	Arg	Asp	Pro	Leu	Asp	Asn	Val	Thr	Thr
		115					120					125			
Ala	Ala	Pro	Gly	Arg	Thr	Pro	Glu	Gly	Leu	Arg	Glu	Leu	Gln	Leu	Arg
		130				135					140				
Ser	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Arg	Gly	Asn	Pro
145					150					155					160
Gln	Leu	Cys	Tyr	Gln	Asp	Met	Val	Leu	Trp	Lys	Asp	Val	Leu	Arg	Lys
				165					170					175	
Asn	Asn	Gln	Leu	Ala	Pro	Val	Asp	Met	Asp	Thr	Asn	Arg	Ser	Arg	Ala
			180					185					190		
Cys	Pro	Pro	Cys	Ala	Pro	Thr	Cys	Lys	Asp	Asn	His	Cys	Trp	Gly	Glu
		195					200					205			
Ser	Pro	Glu	Asp	Cys	Gln	Ile	Leu	Thr	Gly	Thr	Ile	Cys	Thr	Ser	Gly
		210				215					220				
Cys	Ala	Arg	Cys	Lys	Gly	Arg	Leu	Pro	Thr	Asp	Cys	Cys	His	Glu	Gln
225					230					235					240
Cys	Ala	Ala	Gly	Cys	Thr	Gly	Pro	Lys	His	Ser	Asp	Cys	Leu	Ala	Cys
				245					250					255	
Leu	His	Phe	Asn	His	Ser	Gly	Ile	Cys	Glu	Leu	His	Cys	Pro	Ala	Leu
			260					265					270		
Ile	Thr	Tyr	Asn	Thr	Asp	Thr	Phe	Glu	Ser	Met	Leu	Asn	Pro	Glu	Gly
		275					280					285			
Arg	Tyr	Thr	Phe	Gly	Ala	Ser	Cys	Val	Thr	Thr	Cys	Pro	Tyr	Asn	Tyr
	290					295					300				

Fig. 24 (SEQ ID NO:29)

Leu Ser Thr Glu Val Gly Ser Cys Thr Leu Val Cys Pro Pro Asn Asn
 305 310 315 320
 Gln Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser
 325 330 335
 Lys Pro Cys Ala Gly Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg
 340 345 350
 Gly Ala Arg Ala Ile Thr Ser Asp Asn Ile Gln Glu Phe Ala Gly Cys
 355 360 365
 Lys Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly
 370 375 380
 Asn Pro Ser Ser Gly Val Ala Pro Leu Lys Pro Glu His Leu Gln Val
 385 390 395 400
 Phe Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp
 405 410 415
 Pro Glu Ser Phe Gln Asp Leu Ser Val Phe Gln Asn Leu Arg Val Ile
 420 425 430
 Arg Gly Arg Ile Leu His Asp Gly Ala Tyr Ser Leu Thr Leu Gln Gly
 435 440 445
 Leu Gly Ile His Ser Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser
 450 455 460
 Gly Leu Ala Leu Ile His Arg Asn Thr His Leu Cys Phe Val Asn Thr
 465 470 475 480
 Val Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His
 485 490 495
 Ser Gly Asn Arg Pro Glu Glu Ala Cys Gly Leu Glu Gly Leu Val Cys
 500 505 510
 Asn Ser Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln
 515 520 525
 Cys Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu
 530 535 540
 Cys Arg Val Trp Lys Gly Leu Pro Arg Glu Tyr Val Arg Gly Lys His
 545 550 555 560
 Cys Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Ser Ser Glu Thr
 565 570 575
 Cys Tyr Gly Ser Glu Ala Asp Gln Cys Glu Ala Cys Ala His Tyr Lys
 580 585 590
 Asp Ser Ser Ser Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp
 595 600 605

Fig. 24 (SEQ ID NO:29)

Leu Ser Tyr Met Pro Ile Trp Lys Tyr Pro Asp Glu Glu Gly Ile Cys
 610 615 620
 Gln Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Glu
 625 630 635 640
 Arg Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Val Thr Ser Gln Asn
 645 650 655
 Glu Asp Leu Gly Pro Ser Ser Pro Met Asp Ser Thr Phe Tyr Arg Ser
 660 665 670
 Leu Leu Glu Asp Asp Asp Met Gly Glu Leu Val Asp Ala Glu Glu Tyr
 675 680 685
 Leu Val Pro Gln Gln Gly Phe Phe Ser Pro Asp Pro Ala Leu Gly Thr
 690 695 700
 Gly Ser Thr Ala His Arg Arg His Arg Ser Ser Ser Ala Arg Ser Gly
 705 710 715 720
 Gly Gly Glu Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu Glu Pro Pro
 725 730 735
 Arg Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser Asp Val Phe Asp
 740 745 750
 Gly Asp Leu Ala Val Gly Val Thr Lys Gly Leu Gln Ser Leu Ser Pro
 755 760 765
 His Asp Leu Ser Pro Leu Gln Arg Tyr Ser Glu Asp Pro Thr Leu Pro
 770 775 780
 Leu Pro Pro Glu Thr Asp Gly Tyr Val Ala Pro Leu Ala Cys Ser Pro
 785 790 795 800
 Gln Pro Glu Tyr Val Asn Gln Pro Glu Val Arg Pro Gln Ser Pro Leu
 805 810 815
 Thr Pro Glu Gly Pro Pro Pro Pro Ile Arg Pro Ala Gly Ala Thr Leu
 820 825 830
 Glu Arg Pro Lys Thr Leu Ser Pro Gly Lys Asn Gly Val Val Lys Asp
 835 840 845
 Val Phe Ala Phe Gly Gly Ala Val Glu Asn Pro Glu Tyr Leu Ala Pro
 850 855 860
 Arg Ala Gly Thr Ala Ser Gln Pro His Pro Ser Pro Ala Phe Ser Pro
 865 870 875 880
 Ala Phe Asp Asn Leu Tyr Tyr Trp Asp Gln Asn Ser Ser Glu Gln Gly
 885 890 895
 Pro Pro Pro Ser Thr Phe Glu Gly Thr Pro Thr Ala Glu Asn Pro Glu
 900 905 910
 Tyr Leu Gly Leu Asp Val Pro Val Gly Met Gly Ala Leu Phe
 915 920 925

Fig. 25: SEQ ID NO:30

```

atggagctgg cggcctggtg ccgttggggg ttcctctctg cctctctgtc ccccgagacc 60
gcggttacc aagtgtgtac cggtagcgac atgaagttgc gactccctgc cagtcctgag 120
acccacctgg acatgcttcg ccacctctac cagggtgtgc aggtggtgca gggcaatttg 180
gagcttacct acctgcccgc caatgccagc ctctcattcc tgcaggacat ccaggaaagtc 240
cagggataca tgctcatcgc tcacaaccga gtgaaacacg tcccactgca gaggttgccg 300
atcgtgagag ggactcagct ctttgaggac aagtatgccc tggctgtgct agacaaccga 360
gaccttttgg acaacgtcac caccgcccgc ccaggcagaa cccagaaagg gctgcgggag 420
ctgcagcttc gaagtctcac agagatcttg aaggaggagg tttgatccg tgggaaccct 480
cagctctgct accaggacat ggttttgtgg aaggatgtcc tccgtaagaa taaccagctg 540
gctcctgtcg acatggacac caatcgttcc cgggcctgtc cacttctgtc cccaacctgc 600
aaagacaatc actgttgggg tgagagtcct gaagactgtc agatcttgac tggcaccatc 660
tgtactagtg gctgtgcccg gtgcaagggc cggctgccc a ctgactgttg ccatgagcag 720
tgtgctgcag gctgcacggg tccaagcat tctgactgcc tggcctgccc ccacttcaat 780
catagtggta tctgtgagct gcactgccc gcccctcatc cctacaacac agacaccttc 840
gagtcctatc tcaacctga gggctcgctac acctttgggt ccagctgtgt gaccacctgc 900
ccctacaact acctctccac ggaagtggga tcctgcactc tggctctgtc cccgaacaac 960
caagaggtca cagctgagga cggaacacag cgggtgtgaga aatgcagcaa gccctgtgct 1020
ggagtatgct atggtctggg catggagcac ctccgagggg cgagggccat caccagtgc 1080
aatatccagg agtttctggg ctgcaagaag atctttggga gcctggcatt tttgccggag 1140
agctttgatg ggaacctctc ctccggcggt gccccactga agccagagca tctccaagt 1200
ttcgaaaccc tggaggagat cacaggttac ctatacattt cagcatggcc agagagcttc 1260
caagacctca gtgtcttcca gaaccttcgg gtcatctggg gacggattct ccatgatgg 1320
gcttactcat tgacgttgca aggcctgggg attcactcac tggggctacg ctactgccc 1380
gagctgggca gtggattggc tctcattcac cgcaacaccc atctctgctt tgtaaacact 1440
gtaccttggg accagctctt ccggaacctg caccaggccc tactccacag tgggaaccgg 1500
ccagaagagg catgtggtct tgagggcttg gtctgtaact cactgtgtgc ccgtgggcac 1560
tgttgggggc cagggcccac ccagtgtgtc aactgcagtc agttcctccg gggccaggag 1620
tgtgtggagg agtgccagat atggaagggg ctccccaggg agtatgtgag gggcaagcac 1680
tgtctgccat gccaccccgc gtgtcagcct caaacagct cggagacctg ctatggatcg 1740
gaggctgacc agtgtgaggc ttgtgcccac tacaaggact catcttctcg tgtggctcgc 1800
tgccccagtg gtgtgaagcc agacctctcc tacatgccta tctggaagta cccggatgag 1860
gagggcatat gtcagccatg ccccatcaac tgcacccact catgtgtgga cctggacgaa 1920
cgaggtgccc cagcagagca gagagccagc ccagtgcgtc tcagaacga ggacttaggc 1980
ccctccagcc ccattggacag cacctctctac cgttcactgc tggaggatga tgacatgggg 2040
gagctggtcg atgtgaaga gtacctggta cccagcagg gattcttctc cccagaccct 2100
gccctaggtg ctgggagcac agcccaccgc agacaccgca gctcgtcggc caggagtggc 2160
ggtggtgagc tgacctggg cctggagccc tcggaagaag agccccccag atctccactg 2220
gctccctccg aaggggctgg ctccgatgtg tttgatggtg acctggcagt gggggtaacc 2280
aaaggactgc agagcctctc tccacatgac ctacgcccct tacagcggtg cagttaggat 2340
cccacattac ctctgcccc cgagactgat ggctacgttg ctccccctggc ctgcagcccc 2400
cagccccagt atgtgaacca gccagaggtt cggcctcagt ctcccttgac cccagagggt 2460
cctccgcctc ccatecgacc tgctggtgct actctagaaa gacccaagac tctctctcct 2520
gggaaaaaat ggggtgtcaa agacgttttt gcctttgggg gtgctgtgga gaacctgaa 2580
tacctagcac ccagagcagg cactgcctct cagccccacc cttctcctgc cttcagccca 2640
gcctttgaca acctctatta ctgggaccag aactcatcgg agcagggtcc tccaccaagt 2700
acctttgaag ggacccccac tgcaagaac cctgagtacc taggcctgga tgtgccagta 2760
ggaatgggag cgctgttcta a

```


Fig. 26 (SEQ ID NO:31)

Gly Met Gly Ala Leu Phe
5

Fig. 27 (SEQ ID NO:32)

ggaatgggcg cgctgttc 18

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 February 2002 (14.02.2002)

PCT

(10) International Publication Number
WO 02/012341 A3

(51) International Patent Classification?: **C12N 15/12**,
15/62, 15/86, C07K 14/71, 14/82, A61K 31/7088, 31/713,
38/18, 48/00

(21) International Application Number: **PCT/US01/24283**

(22) International Filing Date: **3 August 2001 (03.08.2001)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
09/632,507 **3 August 2000 (03.08.2000) US**

(71) Applicants (for all designated States except US):
**CORIXA CORPORATION [US/US]; 1124 Columbia
St., Suite 200, Seattle, WA 98104 (US). SMITHKLINE
BEECHAM BIOLOGICALS S.A. [BE/BE]; 89, Rue De
L'Institut, B-B-1330 Rixensart (BE).**

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CHEEVER, Mar-
tin, A. [US/US]; 6210 SE 22nd, Mercer Island, WA 98040**

(US). **GHEYSEN, Dirk [US/BE]; 89, Rue de L'Institut,
B-B-1330 Rixensart (BE).**


(74) Agents: **PARENT, Annette, S. et al.; Townsend and
Townsend and Crew LLP, Two Embarcadero Center,
Eighth Floor, San Francisco, CA 94111-3834 (US).**

(81) Designated States (national): **AE, AG, AL, AM, AT (uti-
lity model), AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ (utility model), CZ, DE (uti-
lity model), DE, DK (utility model), DK, DM, DZ, EC, EE
(utility model), EE, ES, FI (utility model), FI, GB, GD, GE,
GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ,
LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN,
MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,
SK (utility model), SK, SL, TJ, TM, TR, TT, TZ, UA, UG,
US, UZ, VN, YU, ZA, ZW.**

(84) Designated States (regional): **ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,**

[Continued on next page]

(54) Title: **HER-2/NEU FUSION PROTEINS**



(57) Abstract: The present invention is generally directed to Her-2/neu fusion proteins, nucleic acid molecules encoding Her-2/neu fusion proteins, viral vectors expressing Her-2/neu fusion proteins, and pharmaceutical compositions (e.g., vaccines) comprising the Her-2/neu fusion proteins and/or nucleic acid molecules encoding the Her-2/neu fusion proteins. The present invention is also directed to methods of treating or preventing cancer by eliciting or enhancing an immune response to the Her-2/neu protein, including for uses in the treatment of malignancies associated with the Her-2/neu oncogene.

CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

(88) Date of publication of the international search report:
17 April 2003

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/24283

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C12N15/12 C12N15/62 C12N15/86 C07K14/71 C07K14/82 A61K31/7088 A61K31/713 A61K38/18 A61K48/00														
According to International Patent Classification (IPC) or to both national classification and IPC														
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N C07K A61K														
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched														
Electronic data base consulted during the International search (name of data base and, where practical, search terms used) EMBL, SEQUENCE SEARCH, EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, CHEM ABS Dat														
C. DOCUMENTS CONSIDERED TO BE RELEVANT <table border="1"> <thead> <tr> <th>Category *</th> <th>Citation of document, with indication, where appropriate, of the relevant passages</th> <th>Relevant to claim No.</th> </tr> </thead> <tbody> <tr> <td>X</td> <td>WO 90 14357 A (GENENTECH INC) 29 November 1990 (1990-11-29) page 11 -page 13; figure 1 ---</td> <td>1-115</td> </tr> <tr> <td>P, X, L</td> <td>WO 00 44899 A (GHEYSEN DIRK ;SMITHKLINE BEECHAM (GB); CORIXA CORP (US); CHEEVER M) 3 August 2000 (2000-08-03) (L: Priority) the whole document ---</td> <td>1-115</td> </tr> <tr> <td>A</td> <td>WO 96 30514 A (DISIS MARY L ;UNIV WASHINGTON (US); CHEEVER MARTIN A (US)) 3 October 1996 (1996-10-03) --- -/--</td> <td></td> </tr> </tbody> </table>			Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.	X	WO 90 14357 A (GENENTECH INC) 29 November 1990 (1990-11-29) page 11 -page 13; figure 1 ---	1-115	P, X, L	WO 00 44899 A (GHEYSEN DIRK ;SMITHKLINE BEECHAM (GB); CORIXA CORP (US); CHEEVER M) 3 August 2000 (2000-08-03) (L: Priority) the whole document ---	1-115	A	WO 96 30514 A (DISIS MARY L ;UNIV WASHINGTON (US); CHEEVER MARTIN A (US)) 3 October 1996 (1996-10-03) --- -/--	
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
X	WO 90 14357 A (GENENTECH INC) 29 November 1990 (1990-11-29) page 11 -page 13; figure 1 ---	1-115												
P, X, L	WO 00 44899 A (GHEYSEN DIRK ;SMITHKLINE BEECHAM (GB); CORIXA CORP (US); CHEEVER M) 3 August 2000 (2000-08-03) (L: Priority) the whole document ---	1-115												
A	WO 96 30514 A (DISIS MARY L ;UNIV WASHINGTON (US); CHEEVER MARTIN A (US)) 3 October 1996 (1996-10-03) --- -/--													
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.														
* Special categories of cited documents : <table border="0"> <tr> <td style="vertical-align: top;"> "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the International filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the International filing date but later than the priority date claimed </td> <td style="vertical-align: top;"> "T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "8" document member of the same patent family </td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the International filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the International filing date but later than the priority date claimed	"T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "8" document member of the same patent family										
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the International filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the International filing date but later than the priority date claimed	"T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "8" document member of the same patent family													
Date of the actual completion of the International search 19 November 2002		Date of mailing of the International search report 28/11/2002												
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016		Authorized officer Devijver, K												

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/24283

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DISIS M L ET AL: "Peptide-based, but not whole protein, vaccines elicit immunity to HER-2/neu, oncogenic self-protein" JOURNAL OF IMMUNOLOGY, THE WILLIAMS AND WILKINS CO. BALTIMORE, US, vol. 156, no. 9, 1 May 1996 (1996-05-01), pages 3151-3158, XP002139506 ISSN: 0022-1767</p> <p>-----</p>	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/24283

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 27-35, 58-66, 89-102, 105, 106 and 108-110 are directed to a method of treatment of the human/animal body (as far as in vivo methods are concerned), the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-66 (completely); 98-115 (in part, as far as applicable)

An isolated protein comprising a Her-2/neu extracellular domain fused to a Her-2/neu phosphorylation domain, wherein the protein is capable of producing an immune response in a warm-blooded animal. A nucleic acid molecule encoding such a protein and a viral vector comprising such a nucleic acid molecule. A pharmaceutical composition comprising such a protein or such a nucleic acid molecule. Methods for eliciting or enhancing an immune response to Her-2/neu protein. An isolated protein comprising a Her-2/neu extracellular domain fused to a fragment of the Her-2/neu phosphorylation domain, wherein the protein is capable of producing an immune response in a warm-blooded animal. A nucleic acid molecule encoding such a protein and a viral vector comprising such a nucleic acid molecule. A pharmaceutical composition comprising such a protein or such a nucleic acid molecule. Methods for eliciting or enhancing an immune response to Her-2/neu protein. Methods for inhibiting the development of a cancer in a patient. Method for removing tumor cells from a biological sample. Method for stimulating and/or expanding T cells specific for a Her-2/neu fusion protein. An isolated T cell population. A method for making such a fusion protein.

2. Claims: 67-97 (completely); 98-115 (in part, as far as applicable)

An isolated protein comprising a Her-2/neu extracellular domain fused to a Her-2/neu intracellular domain, wherein the protein is capable of producing an immune response in a warm-blooded animal. A nucleic acid molecule encoding such a protein and a viral vector comprising such a nucleic acid molecule. A pharmaceutical composition comprising such a protein or such a nucleic acid molecule. Methods for eliciting or enhancing an immune response to Her-2/neu protein. Methods for inhibiting the development of a cancer in a patient. Method for removing tumor cells from a biological sample. Method for stimulating and/or expanding T cells specific for a Her-2/neu fusion protein. An isolated T cell population. A method for making such a fusion protein.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 01/24283

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9014357 A	29-11-1990	AT 155813 T	15-08-1997
		CA 2055441 A1	20-11-1990
		DE 69031120 D1	04-09-1997
		DE 69031120 T2	15-01-1998
		DK 474727 T3	12-01-1998
		EP 0474727 A1	18-03-1992
		ES 2106033 T3	01-11-1997
		WO 9014357 A1	29-11-1990
		US 6015567 A	18-01-2000
		US 6333169 B1	25-12-2001
WO 0044899 A	03-08-2000	AU 2742600 A	18-08-2000
		BR 0007840 A	22-01-2002
		CN 1345374 T	17-04-2002
		CZ 20012587 A3	15-05-2002
		EP 1147190 A1	24-10-2001
		HU 0105303 A2	29-05-2002
		NO 20013701 A	28-09-2001
		TR 200102191 T2	21-12-2001
		WO 0044899 A1	03-08-2000
WO 9630514 A	03-10-1996	US 5801005 A	01-09-1998
		AU 708237 B2	29-07-1999
		AU 5522296 A	16-10-1996
		BR 9607889 A	10-08-1999
		CA 2216601 A1	03-10-1996
		CN 1183117 A	27-05-1998
		CZ 9703096 A3	14-01-1998
		EP 0817846 A1	14-01-1998
		HU 9801826 A2	30-11-1998
		JP 11502710 T	09-03-1999
		NO 974502 A	27-11-1997
		NZ 306616 A	30-03-2001
		WO 9630514 A1	03-10-1996
		US 2002055614 A1	09-05-2002
		US 6075122 A	13-06-2000
		US 5726023 A	10-03-1998
		US 5876712 A	02-03-1999
		US 5846538 A	08-12-1998
		US 5869445 A	09-02-1999

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

